



In vitro antiplasmodial and cytotoxic activities of sesquiterpene lactones from *Vernonia fimbrillifera* Less. (Asteraceae)

Annélie Bordignon^a, Michel Frédérick^a, Allison Ledoux^a, Pierre-Eric Campos^b, Patricia Clerc^b, Thomas Hermann^c, Joëlle Quetin-Leclercq^d and Ewa Ciekiewicz^a

^aLaboratory of Pharmacognosy, CIRM, University of Liège, Liège, Belgium; ^bLaboratoire de Chimie des Substances Naturelles et de Sciences des Aliments (LCSNSA), Faculté des Sciences et Technologies, Université de La Réunion, Saint-Denis, La Réunion; ^cSecteur Nord, Parc National de La Réunion, Saint-Denis, La Réunion; ^dPharmacognosy Research Group, Louvain Drug Research Institute, Université catholique de Louvain, Belgium

ABSTRACT

Due to the *in vitro* antiplasmodial activity of leaf extracts from *Vernonia fimbrillifera* Less. (Asteraceae), a bioactivity-guided fractionation was carried out. Three sesquiterpene lactones were isolated, namely 8-(4'-hydroxymethacrylate)-dehydromelitensin (**1**), onopordopicrin (**2**) and 8 α -[4'-hydroxymethacryloyloxy]-4-*epi*-sonchucarpolide (**3**). Their structures were elucidated by spectroscopic methods (1D and 2D NMR and MS analyses) and by comparison with published data. The isolated compounds exhibited antiplasmodial activity with IC₅₀ values $\leq 5 \mu\text{g/mL}$. Cytotoxicity of the compounds against a human cancer cell line (HeLa) and a mouse lung epithelial cell line (MLE12) was assessed to determine selectivity. Compound **3** displayed promising selective antiplasmodial activity (SI > 10).



1. Introduction

According to the World Health Organization (WHO 2016), malaria caused at least 429 000 deaths globally in 2015. Among these, 70% were children under the age of five and 99% were due to the parasite species *Plasmodium falciparum*. The fight against this major public health problem is threatened by the rapid emergence and spread of parasite strains resistant towards available approved treatments. Urgent efforts are therefore needed to identify new antimalarial drugs. As such, our research groups decided to study Reunion Island as a potential source for the discovery of new antiplasmodial metabolites. This natural patrimony has been identified as a biodiversity hotspot due to its numerous endemic species and its preserved ecosystems (Myers et al. 2000). The National Botanical Conservatory of the Mascarenes indicates that *Vernonia fimbrillifera* Less. is an endemic species from Reunion Island, locally named 'bois de source', which grows in the humid tropical forest (Lavergne and Véra 1989). To the best of our knowledge, neither phytochemical nor pharmacological works have been performed up to now on *V. fimbrillifera*. This study isolated and determined the structure of three antiplasmodial sesquiterpene lactones. Compounds were additionally tested *in vitro* for their cytotoxicity and their selectivity indices were established.

2. Results and discussion

2.1. Chemical composition

An antiplasmodial bioassay-guided fractionation of the CH₂Cl₂ extract from *V. fimbrillifera* leaves was carried out and three compounds (**1–3**) were obtained as active ingredients (Figure 1). Based on their NMR and HR-ESI-MS data and through comparison with those already described in literature, their structures were established as 8-(4'-hydroxymethacrylate)-dehydromelitensin (**1**) (Rustaiyan et al. 1979; Cardona et al. 1989; García et al. 1996), onopordopicrin (**2**) (Lonergan et al. 1992) and 8α-[4'-hydroxymethacryloyloxy]-4-epi-sonchucarpolide (**3**) (Rustaiyan et al. 1986; García et al. 1996).

2.2. Biological activities: antiplasmodial and cytotoxic activities

The three sesquiterpene lactones (**1–3**) isolated from the CH₂Cl₂ extract of *V. fimbrillifera* (leaves) demonstrated *in vitro* antiplasmodial activity with IC₅₀ values of 2.96 ± 0.86, 3.37 ± 0.56 and 3.27 ± 1.32 µg/mL (n = 3, see Table S1), respectively, against the 3D7 strain of *P. falciparum*. This study is the first report of such activities concerning **1** and **3**. The




Figure 1. Chemical structures of compounds 1–3.

antiplasmodial activity of **2** has already been described in *Arctium nemorosum* Lej. (Zimmermann et al. 2012). The cytotoxic activity of the compounds was also assessed in order to determine their selectivity. All tested compounds showed cytotoxic activities (ranging from 10.17 ± 0.88 to 35.48 ± 0.15 $\mu\text{g}/\text{mL}$, $n = 3$) against both mammalian cell lines HeLa and MLE12. As stated in the literature, these compounds have already been found to be cytotoxic (Lonergan et al. 1992; Formisano et al. 2012; Kolli et al. 2012). However, according to WHO guidelines (Pink et al. 2005), compound **3** has a promising selectivity index, being 10 times more potent against *P. falciparum* (Table S1). Several structure–activity relationship studies (Picman 1986; Scotti et al. 2007) have highlighted that α,β -unsaturated ketones seems to be crucial for the biological properties of these compounds due to their particular reactivity (e.g. Michael-type addition) towards free thiol groups. Nevertheless, the higher selectivity of compound **3**, as shown in our work, cannot be easily explained. The presence of an aldehydic moiety on this compound could perhaps contribute to its selectivity, and will be the focus of further studies.

3. Conclusion

To conclude, this study both revealed and confirmed the *in vitro* antiplasmodial and cytotoxic activities of three sesquiterpene lactones isolated from *V. fimbriifera*. To the best of our knowledge, this is the first report of the antiplasmodial activity and the isolation of compounds **1** and **3** from the *Vernonia* genus. These findings justify that plants of this genus are still to be considered as a rich source of compounds (Dogra and Kumar 2015), particularly sesquiterpene lactones, with potential antiplasmodial activities.

Supplementary Material

Experimental, biological activity data and full NMR data are available online.

Acknowledgments

The authors gratefully acknowledge financial support from the European Commission and the Regional Council of Reunion Island: BIOMOL-TCN programme (Activités Thérapeutiques, Cosmétologiques et Nutraceutiques de Molécules Issues de la Biodiversité Terrestre, Marine et Microbienne de la Zone Sud-Ouest de l'Océan Indien), ERDF (European Regional Development Fund 2920). The following reagent was obtained through BEI Resources Repository, NIAID, NIH: *Plasmodium falciparum*, Strain 3D7, MRA-102, contributed by Daniel J. Carucci. The authors wish also to thank the laboratory of Pharmacognosy from the Université catholique de Louvain (MASSMET platform) and particularly Dr. Marie-France Herent for recording MS spectra. The authors would also like to thank Dr. Alexandra Manos-Turvey and Matthew Saive for their editorial assistance.

Disclosure statement

No potential conflict of interest was reported by the authors.

Funding

This research was supported by the Belgian National Fund for Scientific Research (FNRS) [grant number T0190–13].

ORCID

Annélie Bordignon  <http://orcid.org/0000-0003-1954-5994>

References

- Cardona ML, Garcia B, Pedro JR, Sinisterra JF. 1989. Sesquiterpene lactones and an elemene derivative from *Onopordon corymbosum*. *Phytochemistry*. 28:1264–1267.
- Dogra NK, Kumar S. 2015. A review on ethno-medicinal uses and pharmacology of *Vernonia cinerea* Less. *Nat Prod Res*. 29:1102–1117.
- Formisano C, Rigano D, Russo A, Cardile V, Caggia S, Arnold NA, Mari A, Piacente S, Rosselli S, Senatore F, et al. 2012. Phytochemical profile and apoptotic activity of *Onopordum cynarocephalum*. *Planta Med*. 78:1651–1660.
- García B, Skaltsa H, Navarro FI, Pedro JR, Lazari D. 1996. Sesquiterpene lactones and elemene derivatives from *Onopordon myriacanthum*. *Phytochemistry*. 41:1113–1117.
- Kolli EH, León F, Benayache F, Estévez S, Quintana J, Estévez F, Brouard I, Bermejo J, Benayache S. 2012. Cytotoxic sesquiterpene lactones and other constituents of *Centaurea omphalotricha*. *J Braz Chem Soc*. 23:977–983.
- Lavergne R, Véra R. 1989. Etude ethnobotanique des plantes utilisées dans la pharmacopée traditionnelle à La Réunion [Ethnobotanical study of plants used in the traditional pharmacopoeia of Reunion Island]. Paris: Agence de coopération culturelle et technique.
- Loneragan G, Routsis E, Georgiadis T, Agelis G, Hondrelis J, Matsoukas J. 1992. Isolation, NMR studies, and biological activities of onopordopicrin from *Centaurea sonchifolia*. *J Nat Prod*. 55:225–228.
- Myers N, Mittermeier RA, Mittermeier CG, DaFonessa GAB, Kent J. 2000. Biodiversity hotspots for conservation priorities. *Nature*. 403:853–858.
- Picman AK. 1986. Biological activities of sesquiterpene lactones. *Biochem Syst Ecol*. 14:255–281.
- Pink R, Hudson A, Mouriès M-A, Bendig M. 2005. Opportunities and challenges in antiparasitic drug discovery. *Nat Rev Drug Discov*. 4:727–740.
- Rustaiyan A, Nazarians L, Bohlmann F. 1979. Two new elemenolides from *Onopordon leptolepis*. *Phytochemistry*. 18:879–880.
- Rustaiyan A, Ahmadi B, Jakupovic J, Bohlmann F. 1986. Sesquiterpene lactones and eudesmane derivatives from *Onopordon carmanicum*. *Phytochemistry*. 25:1659–1662.
- Scotti MT, Fernandes MB, Ferreira MJP, Emerenciano VP. 2007. Quantitative structure-activity relationship of sesquiterpene lactones with cytotoxic activity. *Bioorganic Med Chem*. 15:2927–2934.
- WHO. 2016. World Malaria report 2016 [Internet]; [cited 2016 Dec 20]. Available from: <https://www.who.int/malaria/publications/world-malaria-report-2016/report/en/>
- Zimmermann S, Thomi S, Kaiser M, Hamburger M, Adams M. 2012. Screening and HPLC-based activity profiling for new antiprotozoal leads from european plants. *Sci Pharm*. 80:205–213.