



HAL
open science

Isothermal kinetics of malondialdehyde content changes in chicken meats

Stéphanie Roux, Matthieu Petit, Elisabeth Baéza, Denis Bastianelli,
Emmanuel Tillard, Elodie Arnaud

► **To cite this version:**

Stéphanie Roux, Matthieu Petit, Elisabeth Baéza, Denis Bastianelli, Emmanuel Tillard, et al.. Isothermal kinetics of malondialdehyde content changes in chicken meats. International Conference on Chemical Reactions in Food, Nov 2012, Prague, Czech Republic. pp.74–74. hal-01193243

HAL Id: hal-01193243

<https://hal.univ-reunion.fr/hal-01193243>

Submitted on 6 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

L-28

ISOTHERMAL KINETICS OF MALONDIALDEHYDE CONTENT CHANGES IN CHICKEN MEATS

Stéphanie Roux^{1*}, Matthieu Petit², Elisabeth Baéza³, Denis Bastianelli⁴, Emmanuel Tillard⁵, Elodie Arnaud⁶

^{1,2,5,6} CIRAD, Saint-Denis, La Réunion, France

³ INRA, Nouzilly, France

⁴ CIRAD, Montpellier, France

* E-mail: stephanie.roux@cirad.fr, Phone: +262 (0)262 92 24 54

Lipid oxidation is a classic parameter of meat quality evaluation, mostly when the composition of the meat is modified, even by modifying the animal's diet or by adding food supplements during post-slaughter meat processing. Malondialdehyde (MDA) is commonly accepted as the major lipid oxidation product in meat and considered as a terminal accumulating compound. It is generally measured together with the other coloured aldehydes resulting from lipid oxidation, with a global methodology called TBARS (thiobarbituric acid reactive substances) index. This index is expressed as equivalent MDA, considering MDA as the preponderant compound by way of a calibration using 1,1,3,3-tetramethoxypropane (TMP) or 1,1,3,3-tetraethoxypropane (TEP). For more precision, MDA is sometimes directly quantified by high performance liquid chromatography (HPLC). The TBARS index or MDA content are mostly measured on fresh meat, but several studies have focused on their changes during meat processing (storage, cooking or other processes). This study set out to characterize changes in MDA content during meat cooking. For this purpose, small samples (1.5 g) of ground chicken thighs were conditioned in plastic bags to form thin films of less than 0.5 mm. The plastic bags were then immersed in a water bath at different temperatures (50, 70 and 100°C) for different immersion times (from 0 to 30 min). Due to the very thin layer of meat, the temperature was considered as constant from the immersion time to the cooling time in an ice bath. MDA content was measured by HPLC. Different meat samples enriched with omega 3 poly-unsaturated fatty acids (transferred in the meat from a dietary flax seeds supplementation) were used to determine the differences in MDA content depending on the vitamin E and xanthophylls' contents in the diets administered to the chickens. The results showed that the appearance kinetics of MDA depended on the heating temperature. At 70°C, three phases were observed in MDA content change: first an increasing phase, then a plateau phase and finally a decreasing phase. At 100°C, the three phases also appeared but the results were more dispersed, probably because of contradictory MDA appearance and disappearance reactions at the same time. Finally, at 50°C, no modification on the MDA content was observed. So, the best temperature for studying MDA content changes during meat cooking was 70°C. It was selected to study differences between the meat sources. The diet composition had no effect on the kinetics of MDA content changes but it greatly affected the MDA contents. The highest MDA contents were determined in meats from chickens reared without a vitamin E dietary supplement.

Keywords: Oxidation, omega 3, chicken meat, malondialdehyde, kinetics

Acknowledgement: Authors want to thanks the "Fond Unique Interministériel" of France, the "Région Réunion" and the "Département de la Réunion" for financing the NUTRIVOL project. They are also grateful to each partner of the project, Crête d'Or Entreprise, Proval, Sanders and Avi-pôle for their participation in the study and to the Lycée Agricole de Saint Joseph for raising the chickens.