



HAL
open science

Microsatellites in the tree *Foetidia mauritiana* (Lecythidaceae) and utility in other *Foetidia* taxa from the Mascarene Islands

Florent Martos, Gérard Lebreton, Eric Rivière, Laurence Humeau,
Marie-Hélène Chevallier

► To cite this version:

Florent Martos, Gérard Lebreton, Eric Rivière, Laurence Humeau, Marie-Hélène Chevallier. Microsatellites in the tree *Foetidia mauritiana* (Lecythidaceae) and utility in other *Foetidia* taxa from the Mascarene Islands. *Applications in Plant Sciences*, 2016, 4 (8), pp.1600034. 10.3732/apps.1600034 . hal-01456701

HAL Id: hal-01456701

<https://hal.univ-reunion.fr/hal-01456701>

Submitted on 28 Jun 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

MICROSATELLITES IN THE TREE *FOETIDIA MAURITIANA* (LECYTHIDACEAE) AND UTILITY IN OTHER *FOETIDIA* TAXA FROM THE MASCARENE ISLANDS¹

FLORENT MARTOS^{2,4}, GÉRARD LEBRETON², ERIC RIVIÈRE², LAURENCE HUMEAU³,
AND MARIE-HÉLÈNE CHEVALLIER²

²CIRAD, UMR PVBMT, F-97410 Saint-Pierre, La Réunion, France; and ³Université de la Réunion, UMR PVBMT, F-97400 Saint-Denis, La Réunion, France

- *Premise of the study:* Polymorphic markers were required for a native tree of the Mascarene Islands, *Foetidia mauritiana* (Lecythidaceae), to investigate the effects of fragmentation of lowland tropical habitats on tree mating systems and on gene flow.
- *Methods and Results:* Using microsatellite enrichment and next-generation sequencing, we identified 13 microsatellite loci (dinucleotide repeats). They were highly polymorphic in 121 trees sampled in the largest three populations on Réunion, revealing 2–17 different alleles per locus. Furthermore, they were found to be polymorphic in conspecific populations on Mauritius and in *F. rodriguesiana* from Rodrigues.
- *Conclusions:* These results indicate the utility of these markers to investigate genetic diversity, mating systems, and gene flow in a genus native to the biodiversity hotspot of Madagascar and the Indian Ocean islands.

Key words: ecological restoration; *Foetidia mauritiana*; island biotas; Lecythidaceae; Madagascar; tropical dry forests.

Trees that belong to the family Lecythidaceae are often used as indicators of disturbance in lowland tropical forests, in particular because they are usually among the most common trees in these rich but fragile ecosystems (Mori et al., 2007). In addition to their ecological significance, some species may also be economically important, such as the Brazil nut tree *Bertholletia excelsa* Bonpl. For these reasons, polymorphic genetic markers have been developed for several species of several subfamilies of Lecythidaceae, mostly in taxa occurring as large trees in the Amazon Basin (e.g., *Bertholletia* Bonpl. [Reis et al., 2009], *Cariniana* Casar. [Guidugli et al., 2009, 2010], and *Lecythis* Loeffl. [Rodrigues et al., 2015]), but also in a few other taxa found in the Old World (e.g., *Barringtonia* J. R. Forst. & G. Forst. [Xie et al., 2015]). However, to our knowledge, polymorphic markers are not yet available for the representatives of Lecythidaceae in the biodiversity hotspot formed by Madagascar and the Indian Ocean islands.

Out of 18 species that make up the genus *Foetidia* Comm. ex Lam. (subfamily Foetidioideae), 17 are endemic to island biotas in Madagascar, the Comoros, and the Mascarene Islands, while one species is found only on the African continent in Tanzania (Prance, 2008; Labat et al., 2011). The endemic species

F. mauritiana Lam. was common in drier areas of Mauritius and Réunion where precipitation is low and temperatures are high, relative to the wet conditions generally found on these two tropical islands. However, for this species as for many indigenous taxa adapted to dry tropical habitats, populations have undergone rapid decline in less than 400 years since human settlement, and those few remaining stands are left in highly fragmented landscapes on both islands. This species is considered endangered in Réunion and Mauritius. There is an urgent need to protect and restore natural communities in tropical dry habitats on the Indian Ocean islands as well as worldwide (Miles et al., 2006). A European Union-supported project, Life+ Corexrun, was launched in 2009 on Réunion; it aims at both reintroducing 48 indigenous plant species (including *F. mauritiana*) and controlling invasions by alien plants within and around semi-dry forest stands.

METHODS AND RESULTS

Genomic DNA of *F. mauritiana* was extracted with the DNeasy Plant Mini Kit (QIAGEN, Hilden, North Rhine-Westphalia, Germany). Production of a microsatellite-enriched library was outsourced to the high-throughput platform set up by Genoscreen (Lille, Nord-Pas-de-Calais-Picardie, France). Following the method described in Malausa et al. (2011), 1 µg of genomic DNA was mechanically fragmented, ligated to standard adapters (Adap-F: GTTTAAGGCCTAGC-TAGCAGAATC and Adap-R: GATTCTGCTAGCTAGGCCTT), and enriched by addition of eight biotin-labeled oligoprobes corresponding to the following microsatellite motifs: (TG)_n, (TC)_n, (AAC)_n, (AAG)_n, (AGG)_n, (ACG)_n, (ACAT)_n, and (ACTC)_n. Enriched DNA was isolated using Dynabeads (Invitrogen, Waltham, Massachusetts, USA) and amplified by PCR with primers corresponding to the library adapters (PCR protocol not communicated by Genoscreen). Sequencing was carried out through 454 GS-FLX Titanium pyrosequencing (Roche Applied Science, Penzberg, Bavaria, Germany). Sequences were analyzed using the bioinformatics program QDD (Megléczy et al., 2010), which detects microsatellite sequences and designs primers in flanking regions. We then selected 13 primer pairs among the microsatellite sequences (dinucleotide

¹Manuscript received 17 March 2016; revision accepted 11 May 2016.

The authors thank S. Dafreville, T. M'sa, and J. Segrestin (laboratory assistance); P. Adolphe, S. Baret, L. Calichiana, M. Félicité, R. Lucas, H. Thomas (assistance on Réunion); and J. T. Genave, R. Parmananda, J.-C. Sevathian, and A. Waterstone (assistance on Mauritius and Rodrigues). This work was funded by the European Regional Development Fund (ERDF), by the Région Réunion, and by the Centre de Coopération International en Recherche Agronomique pour le Développement (CIRAD).

⁴Author for correspondence: florent.martos@cirad.fr

doi:10.3732/apps.1600034

TABLE 1. Characteristics of 13 microsatellite loci developed in *Foetidia mauritiana*.

Locus	Primer sequences (5'–3')	Repeat motif	Allele size range (bp)	T_a (°C)	GenBank accession no.
FmCIR27	F: AAGGAAAAGATGCATGCCAA R: AGACAATTCTAAACAAGATAGGACG	(CA) ₁₄	79–97	57	KU713062
FmCIR29	F: CATGTGGATTCCAAAATGGC R: TTGCAATGATAATTCACCAACC	(AG) ₁₂	87–93	57	KU713063–70
FmCIR31	F: CATGAATAGGTCCCAGGCTC R: TATCTATGCTTGCGTGTGCG	(CA) ₁₃	86–100	57	KU713071
FmCIR32	F: GAAGAGCACAGAAGAACACATCA R: GCCACTTCTATCATCGGGAG	(GA) ₁₂	92–96	57	KU713072
FmCIR43	F: AGCATGACCCCTAAACCCCTAA R: AACATGACTGTGATGGCCTAAG	(AC) ₁₄	119–143	57	KU713073
FmCIR45	F: GTGACTAGCTCACCAAGAGCC R: TTGTCCCTAACGTTTCTTCTC	(GA) ₁₂	132–146	57	KU713074
FmCIR47	F: TTCTTCACTGAGTGTATTTCCATAGG R: TGTAATAAGTTCCTGGACCGAC	(GA) ₁₂	134–156	57	KU713075
FmCIR52	F: TGCTACTCTGTGGTGTGAAAGG R: GCATGAACAGGCAGAACATAA	(AC) ₁₄	144–204	57	KU713076
FmCIR16	F: GAAAAGTCACGGTTCCTCCG R: TTTGGTTCGAGGATGGGTAG	(AC) ₁₅	164–181	57	KU713061
FmCIR57	F: TAAAATCAACAACCTAAAACACGAA R: TGAGATTACCCAGGAGCAGG	(TC) ₁₂	188–196	57	KU713077
FmCIR61	F: GAGCATTGAAGTAGCTGGT R: ATTTGAGCCCTGAACCAATG	(GA) ₁₁	196–228	57	KU713078
FmCIR11	F: TGAAGCTCAAGCAATGGAA R: GGGTCCGGTAGGGTACTGTT	(TC) ₁₃	194–208	57	KU713058–60
FmCIR3	F: CGATTGGCATGGAGAAAG R: GCTCTTGCCCAAGAAGGTC	(AG) ₁₀	286–288	57	KU713056–57

Note: T_a = annealing temperature.

repeats), because they had adequate flanking regions for designing primers (see Table 1 for locus information, primers, and GenBank accession numbers). Eight additional microsatellite loci are provided in this paper, although population testing was not conducted for these markers (Appendix 1).

For biological validation, we selected the main wild populations on Réunion. This included 49 individuals sampled near the Lataniers River (mean elevation 330 m), 45 individuals on the southern slope of the Grande-Chaloupe River (545 m), and 27 individuals near the Tamarins River (270 m), for a total of 121 individuals (sampling authorized by the Parc National de La Réunion, the Office National des Forêts, the Département de La Réunion, and the Conservatoire du Littoral). Because the species is considered critically endangered on Réunion, we harvested no more than 1–2 leaves per individual tree, with the exception of three individuals (one per locality) from which voucher specimens were made (see Appendix 2). Plant genomic DNA was isolated with the DNeasy Plant Mini Kit (QIAGEN). Multiplex PCR was performed in a total volume of 15 μ L containing 7.5 μ L 2 \times Type-it Multiplex PCR Master Mix (QIAGEN), 1.5 μ L

5 \times Q-Solution, 0.2 μ M each primer, and 20–50 ng template DNA. The thermal cycling protocol was as follows: initial denaturation at 95°C for 5 min; 28 cycles of denaturation at 95°C for 30 s, primer annealing at 57°C for 90 s, extension at 72°C for 30 s; and final extension at 60°C for 30 min. PCR products were diluted in HPLC-grade water (1:10), denatured in formamide, and separated on a 16-capillary ABI PRISM 3130xl Genetic Analyzer (Applied Biosystems, Foster City, California, USA); GeneScan 500 LIZ (Applied Biosystems) was used for sizing alleles in the expected range of 80–300 bp.

Allele sizes were estimated using the Microsatellite Plugin version 1.4 implemented in Geneious version 8.1.7 (Biomatters, Auckland, New Zealand). The number of different alleles and observed and expected heterozygosity were calculated for each locus and population in GenAIEx (Peakall and Smouse, 2006, 2012). Hardy–Weinberg exact tests (9999 iterations) and linkage disequilibrium were analyzed in GENEPOP version 4.2 (Rousset, 2008). The presence of null alleles was estimated with MICRO-CHECKER version 2.2.3 (van Oosterhout et al., 2004). The minimum number of microsatellite loci necessary to discriminate

TABLE 2. Genetic properties of the 13 newly developed microsatellites of *Foetidia mauritiana*.^a

Locus	Grande-Chaloupe (n = 45)			Lataniers (n = 49)			Tamarins (n = 27)		
	A	H_o	H_e^b	A	H_o	H_e^b	A	H_o	H_e^b
FmCIR27	8	0.163	0.678***	7	0.408	0.695***	6	0.480	0.762***
FmCIR29	4	0.644	0.621	4	0.521	0.683	4	0.741	0.735
FmCIR31	4	0.489	0.539	4	0.592	0.586	5	0.741	0.651
FmCIR32	3	0.578	0.560	3	0.510	0.574	3	0.667	0.592
FmCIR43	10	0.778	0.752	10	0.918	0.831	10	0.926	0.783
FmCIR45	5	0.727	0.624	4	0.59	0.526	3	0.64	0.506
FmCIR47	2	0.001	0.044*	7	0.167	0.299***	4	0.037	0.372***
FmCIR52	17	0.867	0.886	15	0.857	0.874	11	0.889	0.845
FmCIR16	10	0.432	0.767***	10	0.479	0.808***	11	0.667	0.804*
FmCIR57	5	0.511	0.720	5	0.653	0.690	4	0.593	0.536
FmCIR61	11	0.644	0.675	11	0.714	0.653	9	0.778	0.696
FmCIR11	6	0.578	0.548	6	0.408	0.394	5	0.481	0.514
FmCIR3	2	0.023	0.107***	2	0.020	0.230***	2	0.037	0.324***

Note: A = number of alleles; H_e = expected heterozygosity; H_o = observed heterozygosity; n = number of individuals sampled.

^aAll three populations are located on Réunion; see Appendix 2 for locality and voucher information.

^bAsterisks refer to significant deviations from Hardy–Weinberg equilibrium (* P < 0.05, *** P < 0.01).

TABLE 3. Cross-species amplification (showing number of different alleles and size range) of the 13 newly developed microsatellites of *Foetidia mauritiana*.^a

Locus	<i>F. mauritiana</i>		<i>F. rodriguesiana</i>
	Réunion (<i>n</i> = 121)	Mauritius (<i>n</i> = 28)	Rodrigues (<i>n</i> = 30)
FmCIR27	8 (79–97)	1 (91)	3 (97–101)
FmCIR29	4 (87–93)	6 (85–95)	3 (79–89)
FmCIR31	5 (86–100)	3 (94–98)	—
FmCIR32	3 (92–96)	3 (92–96)	2 (98–102)
FmCIR43	13 (119–143)	5 (121–135)	6 (121–133)
FmCIR45	5 (132–146)	6 (130–146)	5 (122–134)
FmCIR47	7 (134–156)	5 (132–152)	1 (134)
FmCIR52	21 (144–204)	6 (154–168)	10 (146–184)
FmCIR16	14 (164–181)	8 (164–180)	9 (164–182)
FmCIR57	5 (188–196)	2 (190–192)	9 (192–210)
FmCIR61	13 (196–228)	9 (192–222)	17 (200–264)
FmCIR11	7 (194–208)	6 (196–218)	1 (200)
FmCIR3	2 (286–288)	4 (278–288)	1 (286)

Note: — = no amplification; *n* = number of individuals sampled.

^a See Appendix 2 for locality and voucher information.

individuals of *F. mauritiana* was assessed using the package *poppr* in R software (Kamvar et al., 2015).

All microsatellite loci revealed polymorphisms in *F. mauritiana* populations on Réunion. The number of different alleles per locus ranged from two to 17 (Table 2). Four loci showed significant deviation from Hardy–Weinberg equilibrium: FmCIR27, FmCIR47, FmCIR16, and FmCIR3. No significant linkage disequilibrium was detected between pairs of loci. We found that the minimum number of loci necessary to discriminate individuals in the data set was eight (data not shown).

Transferability of the microsatellite loci was tested on 28 individuals of *F. mauritiana* and 30 individuals of *F. rodriguesiana* F. Friedmann sampled across Mauritius and Rodrigues, respectively (sampling authorized by the National Parks and Conservation Service of Mauritius). Conspecific populations on different islands were tested because they are expected to experience strong genetic isolation. *Foetidia rodriguesiana* is morphologically similar to *F. mauritiana* (Prance, 2008). Using the above-mentioned protocol, we found that all loci amplified in *Foetidia* populations found on other islands, with the exception of FmCIR31, which did not amplify in *F. rodriguesiana* (Table 3). Moreover, most loci were polymorphic in the Mauritius and Rodrigues populations.

CONCLUSIONS

We developed 13 polymorphic genetic markers for *Foetidia*, a widespread genus in the Indian Ocean islands biodiversity hotspot. They will aid in designing priority populations for conservation and implementing adaptive conservation plans for the genus. They may also be used to study mating systems and pollen and seed flow between lowland forest fragments.

LITERATURE CITED

GUIDUGLI, M. C., T. DE CAMPOS, A. C. B. DE SOUSA, J. M. FERES, A. M. SEBBENN, M. A. MESTRINER, E. P. B. CONTEL, AND A. L. ALZATE-MARIN.

2009. Development and characterization of 15 microsatellite loci for *Cariniana estrellensis* and transferability to *Cariniana legalis*, two endangered tropical tree species. *Conservation Genetics* 10: 1001–1004.
- GUIDUGLI, M. C., K. A. GUERRIERI, M. A. MESTRINER, E. P. B. CONTEL, C. A. MARTINEZ, AND A. L. ALZATE-MARIN. 2010. Genetic characterization of 12 heterologous microsatellite markers for the giant tropical tree *Cariniana legalis*. *Genetics and Molecular Biology* 33: 131–134.
- KAMVAR, Z. N., J. C. BROOKS, AND N. J. GRÜNWARD. 2015. Novel R tools for analysis of genome-wide population genetic data with emphasis on clonality. *Frontiers in Genetics* 6: 208.
- LABAT, J.-N., E. BIDAULT, AND G. VISCARDI. 2011. A new critically endangered species of *Foetidia* (Lecythidaceae, subfamily Foetidioideae) recently discovered in Mayotte, Comoros archipelago. *Adansonia* 33: 263–269.
- MALAUSSA, T., A. GILLES, E. MEGLÉCZ, H. BLANQUART, S. DUTHOY, C. COSTEDOAT, ET AL. 2011. High-throughput microsatellite isolation through 454 GS-FLX Titanium pyrosequencing of enriched DNA libraries. *Molecular Ecology Resources* 11: 638–644.
- MEGLÉCZ, E., C. COSTEDOAT, V. DUBUT, A. GILLES, T. MALAUSSA, N. PECH, AND J.-F. MARTIN. 2010. QDD: A user-friendly program to select microsatellite markers and design primers from large sequencing projects. *Bioinformatics* 26: 403–404.
- MILES, L., A. C. NEWTON, R. S. DEFRIES, C. RAVILIOUS, I. MAY, S. BLYTH, V. KAPO, AND J. E. GORDON. 2006. A global overview of the conservation status of tropical dry forests. *Journal of Biogeography* 33: 491–505.
- MORI, S. A., C.-H. TSOU, C.-C. WU, B. CRONHOLM, AND A. A. ANDERBERG. 2007. Evolution of Lecythidaceae with an emphasis on the circumscription of neotropical genera: Information from combined *ndhF* and *trnL-F* sequence data. *American Journal of Botany* 94: 289–301.
- PEAKALL, R., AND P. E. SMOUSE. 2006. GenAlEx 6: Genetic analysis in Excel. Population genetic software for teaching and research. *Molecular Ecology Notes* 6: 288–295.
- PEAKALL, R., AND P. E. SMOUSE. 2012. GenAlEx 6.5: Genetic analysis in Excel. Population genetic software for teaching and research—an update. *Bioinformatics (Oxford, England)* 28: 2537–2539.
- PRANCE, G. T. 2008. A revision of *Foetidia* (Lecythidaceae subfamily Foetidioideae). *Brittonia* 60: 336–348.
- REIS, A. M. M., A. C. BRAGA, M. R. LEMES, R. GRIBEL, AND R. G. COLLEVATTI. 2009. Development and characterization of microsatellite markers for the Brazil nut tree *Bertholletia excelsa* Humb. & Bonpl. (Lecythidaceae). *Molecular Ecology Resources* 9: 920–923.
- RODRIGUES, A. B., C. T. FLORENCE, E. MARIANO-NETO, AND F. A. GAJOTTO. 2015. First microsatellite markers for *Lecythis pisonis* (Lecythidaceae), an important resource for Brazilian fauna. *Conservation Genetics Resources* 7: 437–439.
- ROUSSET, F. 2008. GENEPOP'007: A complete reimplementation of the GENEPOP software for Windows and Linux. *Molecular Ecology Resources* 8: 103–106.
- VAN OOSTERHOUT, C., W. F. HUTCHINSON, D. P. M. WILLIS, AND P. SHIPLEY. 2004. MICRO-CHECKER: Software for identifying and correcting genotyping errors in microsatellite data. *Molecular Ecology Resources* 4: 535–538.
- XIE, H., Y. YUAN, X. FANG, Y. LIU, C. YANG, J. JIN, F. TAN, AND Y. HUANG. 2015. Development of EST-SSR markers in *Barringtonia* (Lecythidaceae) and cross-amplification in related species. *Applications in Plant Sciences* 3: 1500080.

APPENDIX 1. Eight additional microsatellite loci identified in *Foetidia mauritiana*.^a

Locus	Primer sequences (5'–3')	Repeat motif	Allele size (bp)
FmCIR7	F: GGTAACAGCTCAAGCCCAA R: TTATTCGGCCAAACAACCTC	(AAC) ₁₂	148
FmCIR37	F: AAGAAAATTCTGCCCGATTG R: CACTGTTGCAAGAGGGTGAG	(CT) ₁₂	113
FmCIR38	F: TTTGGAGATCCTATGTTGAGCA R: AAATTTTCCCAAATTAACCCAA	(TG) ₁₂	116
FmCIR41	F: CTCCTCCCACTGTTTCTCG R: TATGGCAAGGGTTTGGATGT	(CT) ₁₂	127
FmCIR48	F: AAGGATAACTATCAACCTCAAGCA R: ACCCTCAGGTATGTGTCAGTTT	(AC) ₁₂	144
FmCIR49	F: CCATGTTTGCCCATGCAC R: TGGCCGAGATGCATAATGT	(CA) ₁₂	144
FmCIR58	F: TTGTCTCTGTCTAAAGTTTGTGAGG R: TCGCGAAATCTTGACCATC	(AC) ₁₁	190
FmCIR68	F: AGGTCAGTGTCTACCAATACAG R: CCAGAAATCTCCTATCCTCTTGC	(GA) ₁₁	173

^aPolymorphism has not been assessed in these markers.

APPENDIX 2. Voucher information for *Foetidia* populations used in this study.

Species	Voucher specimen no. ^a	Collection locality	Geographic coordinates	N
<i>F. mauritiana</i>	Cir 919	Grande-Chaloupe/Cap Francis, La Possession, Réunion	20°55'24.532"S, 55°23'16.63"E	45
<i>F. mauritiana</i>	Cir 875	Ravine des Lataniers, La Possession, Réunion	20°56'41.467"S, 55°20'56.857"E	49
<i>F. mauritiana</i>	Cir 936	Ravine des Tamarins, Saint-Denis, Réunion	20°53'50.798"S, 55°23'1.33"E	27
<i>F. mauritiana</i>	—	Black River Gorges National Park, Mauritius	20°23'53.467"S, 57°25'32.564"E	11
<i>F. mauritiana</i>	—	Chamarel, Black River, Mauritius	20°25'4.843"S, 57°23'10.731"E	8
<i>F. mauritiana</i>	—	Domaine du Chasseur National Park, Anse Jonchée, Mauritius	20°20'46.128"S, 57°45'23.91"E	7
<i>F. mauritiana</i>	—	Bras d'Eau National Park, Mauritius	20°8'42.781"S, 57°43'35.767"E	2
<i>F. rodriguesiana</i>	—	Anse Quitor Nature Reserve, Rodrigues	19°45'18.9"S, 63°22'10.599"E	5
<i>F. rodriguesiana</i>	—	Graviers, Rodrigues	19°43'57"S, 63°28'38.099"E	5
<i>F. rodriguesiana</i>	—	Mourouk, Rodrigues	19°44'13.7"S, 63°27'40.499"E	5
<i>F. rodriguesiana</i>	—	Mont Malgache, Rodrigues	19°43'40.598"S, 63°27'21.099"E	5
<i>F. rodriguesiana</i>	—	Grande Montagne, Rodrigues	19°42'14.501"S, 63°27'56.699"E	4
<i>F. rodriguesiana</i>	—	Baie Malgache, Rodrigues	19°43'39.299"S, 63°23'24.399"E	1
<i>F. rodriguesiana</i>	—	Rivière Cascade Victoire, Rodrigues	19°43'49.4"S, 63°26'59.798"E	1
<i>F. rodriguesiana</i>	—	Caverne Provert, Rodrigues	19°40'26.4"S, 63°26'25.299"E	1
<i>F. rodriguesiana</i>	—	Crève-coeur, Rodrigues	19°40'43.201"S, 63°25'47.499"E	1
<i>F. rodriguesiana</i>	—	Grande Baie, Rodrigues	19°40'57.202"S, 63°26'57.001"E	1
<i>F. rodriguesiana</i>	—	Solitude, Rodrigues	19°41'37"S, 63°26'15.399"E	1

Note: N = number of individuals.

^aVouchers for the *F. mauritiana* populations collected in Réunion were deposited at Université de la Réunion (REU). Vouchers were not collected for the *F. mauritiana* populations sampled in Mauritius and the *F. rodriguesiana* populations sampled in Rodrigues per agreements with local authorities to collect DNA samples only.