



HAL
open science

Monitoring and mapping forest fires in the south-western Indian Ocean using very high resolution optical imagery

Christophe Révillion, Thibault Catry

► **To cite this version:**

Christophe Révillion, Thibault Catry. Monitoring and mapping forest fires in the south-western Indian Ocean using very high resolution optical imagery. Pléiades Days, Apr 2014, Toulouse, France. , 2014. hal-01475838

HAL Id: hal-01475838

<https://hal.univ-reunion.fr/hal-01475838v1>




Submitted on 24 Feb 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1) The South-western Indian Ocean is characterized by various heterogeneous insular territories. Many of these islands are biodiversity hotspots and in order to preserve this heritage, some of them are protected.

2) In 2013, the monitoring and mapping of the fires of Etang-Saint-Paul (July, La Reunion) and Europa (June) were carried out by the SEAS-OI (Survey of Environment Assisted by Satellites in the Indian Ocean), in collaboration with the ONF (Office National des Forêts) of La Reunion and the SDIS (Service Départemental d'Incendie et de Secours), using high resolution SPOT 5 imagery. These maps allowed the estimation of the fire extent, to about 4 and 70 hectares for Europa and Etang-Saint-Paul respectively, and to support field intervention during the crisis.



The Study areas, Europa and La Reunion (A) and focus on Saint-Paul (B)

© CNES (2013), Distribution Astrium services / Spot images S.A., France, all rights reserved.

Land cover

Anthropogenic

- Urban
- Pasture
- Cropland

Forest or semi-natural


- Forest
- Non forest vegetation
- Herbaceous cover
- Barren land

Wetland and water

- Aquatic vegetation with Typha domingensis
- Aquatic vegetation with Cyperus papyrus
- Mangrove
- Water


Fire and recovery

- Delimitation of the burned area
- burned area
- Barren land area after the fire
- Recovery in progress by the wet vegetation




Example of Typha Domingensis (A) and Cyperus papyrus (B)

3) The monitoring of the post-fire recovery was realized using Pléiades images. Very high spatial resolution images allowed us to understand more precisely how different plant groupings recovered after the fire. On the pond of Saint-Paul vegetation dominated by Cyperus Papyrus (in dark green on map) is almost back to its pre-fire condition in six months, however areas with Typha domingensis began to regenerate but much more slowly.



PLEIADES imagery



© CNES (2013, 2014), Distribution Astrium services / Spot images S.A., France, all rights reserved.

4) This study was part of the operational mission of the SEAS-OI station, gathering local actors in the management of forest fire crisis in the south-western Indian Ocean. In particular, we demonstrate here the interest of the satellite approach in the management of natural disasters on remote territories like Europa island, and in protected area where a rapid response is needed. Thanks to the combination of optical SPOT 5 and PLEIADES high resolution products, SEAS-OI was able to provide operational products for the management of these fires, and the monitoring of .