

Comprehensive proteome profiling in Aedes albopictus to decipher Wolbachia-arbovirus interference phenomenon

Yoann Saucereau, Claire Valiente Moro, Cindy Dieryckx, Jean-William Dupuy, Florence-Hélène Tran, Vincent Girard, Patrick Potier, Patrick Mavingui

To cite this version:

Yoann Saucereau, Claire Valiente Moro, Cindy Dieryckx, Jean-William Dupuy, Florence-Hélène Tran, et al.. Comprehensive proteome profiling in Aedes albopictus to decipher Wolbachia-arbovirus interference phenomenon. BMC Genomics, 2017, 18 (1), pp.635. $10.1186/s12864-017-3985$ -y. hal-01589764

HAL Id: hal-01589764 <https://hal.univ-reunion.fr/hal-01589764>

Submitted on 19 Sep 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

[Distributed under a Creative Commons Attribution 4.0 International License](http://creativecommons.org/licenses/by/4.0/)

RESEARCH ARTICLE Example 2014 12:30 The Contract of Contract ACCESS

Comprehensive proteome profiling in Aedes albopictus to decipher Wolbachia-arbovirus interference phenomenon

Yoann Saucereau^{1,2,3,4}, Claire Valiente Moro^{1,2,3,4}, Cindy Dieryckx⁵, Jean-William Dupuy⁶, Florence-Hélène Tran^{1,2,3,4}^, Vincent Girard⁵, Patrick Potier^{1,2,3,4} and Patrick Mavingui^{1,2,3,4,7*}

Abstract

Background: Aedes albopictus is a vector of arboviruses that cause severe diseases in humans such as Chikungunya, Dengue and Zika fevers. The vector competence of Ae. albopictus varies depending on the mosquito population involved and the virus transmitted. Wolbachia infection status in believed to be among key elements that determine viral transmission efficiency. Little is known about the cellular functions mobilized in Ae. albopictus during co-infection by Wolbachia and a given arbovirus. To decipher this tripartite interaction at the molecular level, we performed a proteome analysis in Ae. albopictus C6/36 cells mono-infected by Wolbachia wAlbB strain or Chikungunya virus (CHIKV), and bi-infected.

Results: We first confirmed significant inhibition of CHIKV by Wolbachia. Using two-dimensional gel electrophoresis followed by nano liquid chromatography coupled with tandem mass spectrometry, we identified 600 unique differentially expressed proteins mostly related to glycolysis, translation and protein metabolism. Wolbachia infection had greater impact on cellular functions than CHIKV infection, inducing either up or down-regulation of proteins associated with metabolic processes such as glycolysis and ATP metabolism, or structural glycoproteins and capsid proteins in the case of bi-infection with CHIKV. CHIKV infection inhibited expression of proteins linked with the processes of transcription, translation, lipid storage and miRNA pathways.

Conclusions: The results of our proteome profiling have provided new insights into the molecular pathways involved in tripartite Ae. albopictus-Wolbachia-CHIKV interaction and may help defining targets for the better implementation of Wolbachia-based strategies for disease transmission control.

Keywords: Aedes albopictus, Wolbachia, Viral inhibition, Proteome, Glycolysis, Metabolism, miRNA

Background

The Asian tiger mosquito Aedes albopictus is a species native to South and East Asia, with a great capacity for invasion. It has been classified by the WHO as the fourth most invasive species in the world [1]. Since the mid-twentieth century, Ae. albopictus has considerably increased its distribution, and is currently present on five continents [2]. Ae. albopictus is involved in the transmission of many human-infecting arboviruses, including Chikungunya virus (CHIKV), Dengue virus (DENV), and

probably Zika virus [3–5]. Historically, Ae. Albopictus has been considered of secondary importance in terms of arbovirosis incidence relative to Aedes aegypti. However, this has changed since the implication of Ae. albopictus in the explosive epidemics of CHIKV on La Reunion Island and neighboring islands in southern Indian Ocean [6, 7], as well as in the CHIKV outbreaks in Italy [8] and successive autochthonous transmissions of both CHIKV and DENV in metropolitan France [9–12]. Efficient transmission of CHIKV has been associated with a mutation in E1 envelope glycoprotein (Ala-226-Val) that increases viral infectivity in Ae. albopictus compared to Ae. aegypti [6, 13]. Advances in technologies of large-scale analysis and the availability of genome sequencing allow

© The Author(s). 2017 **Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License [\(http://creativecommons.org/licenses/by/4.0/](http://creativecommons.org/licenses/by/4.0/)), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver [\(http://creativecommons.org/publicdomain/zero/1.0/](http://creativecommons.org/publicdomain/zero/1.0/)) applies to the data made available in this article, unless otherwise stated.

^{*} Correspondence: patrick.mavingui@cnrs.fr ^ˆDeceased

¹Université de Lyon, Lyon, France

² Université Lyon 1, Villeurbanne, France

Full list of author information is available at the end of the article

the meaning of this host tolerance to be examined at the molecular level by the screening of cell factors possibly mobilized during viral cell invasion. In Ae. aegypti differential trends of proteomic expression were seen in the midgut and salivary glands infected by CHIKV or DENV in comparison to uninfected specimens [14, 15]. Using cellular models, microarrays studies have shown that CHIKV enters Ae. albopictus cells by clathrin-dependent endocytosis [16], activating diverse biological processes, including protein folding and metabolic pathways [17]. Overall, the modulation of the synthesis of some classes of host proteins clearly favors virus survival, replication and transmission [18].

Ae. albopictus is naturally infected by the intracellular bacterium Wolbachia pipientis that are maternally transmitted from mother to offspring. Two distinct Wolbachia strains (wAlbA and wAlbB) are present in variable density in Ae. albopictus tissues [19–21] and they usually induce sterility through the phenomenon known as cytoplasmic incompatibility [22–24]. In Ae. aegypti, naturally devoid of Wolbachia, transinfected females harboring the wAlbB strain have been found to inhibit the transmission of both CHIKV and DENV [25, 26]. In Ae. albopictus dissemination of DENV serotype 2 to salivary glands of Wolbachia-infected Ae. albopictus from La Reunion was considerably diminished in comparison to Wolbachia-uninfected individuals generated by antibiotic treatment [27]. When Ae. albopictus was transinfected with Wolbachia wMel strain derived from Drosophila melanogaster, the transmission of DENV serotype 2 was totally abolished [28]. However, the inhibitory effect of Wolbachia is not universal [29, 30], and one study noted an increase in parasite infection in Anopheles [31], suggesting that variable mechanisms are involved depending on the interacting partners. Investigations into the molecular mechanisms behind Wolbachia interference have suggested that the bacterium may act by modulating expression of insect innate immune genes, including antimicrobial peptides, or more broadly by inducing oxidative and metabolic stresses that will in turn impact the behavior of the infectious agent in the host cells [32, 33]. It is also proposed that Wolbachia and viruses would compete for the host cells' resources [34].

We recently showed that the wAlbB strain was able to block CHIKV infection in Ae. albopictus C6/36 cell lines relative to uninfected cells [35]. This is in line with observations in all studies using cellular models [36, 37], suggesting that viral inhibition is common in such simplified systems, possibly due in part to the proximity of the interacting partners. Thus, cellular models could represent interesting systems to decipher the mechanisms involved in the tripartite interactions between Wolbachia, arboviruses and host cells. Both naturally and artificially Wolbachia-infected Aedes cell lines have

shown changes in the expression of several genes involved in structural, metabolic and stress functions [38, 39]. On the other hand CHIKV was reported to activate cellular functions necessary for infection and persistence [17]. However, no molecular mechanism for the interplay between Wolbachia and CHIKV in Ae. albopictus has been proposed to date. Therefore, in this study we used proteome profiling of Ae. albopictus C6/36 cell lines to discover how Wolbachia-infected cells reacted when challenged with CHIKV. Two-dimensional electrophoresis (2DE) followed by nano liquid chromatography and coupled with tandem mass spectrometry (nanoLC-MS/ MS) showed differentially expressed proteins likely belonging to diverse processes of glycolysis, protein metabolism, protein modification and amino acid metabolism. Overall, the innovative proteomic approach used in this descriptive work provided potential candidates involved in the tripartite interaction between mosquito-CHIKV-Wolbachia. Future investigation will focus on the functional studies to validate the more promising candidates implicated in cellular processes that mediated the interplay between microbes.

Methods

Mosquito cell line and virus

The C6/36 cells infected by Wolbachia wAlbB strain and uninfected cells generated by removing the bacterium through tetracycline treatment [35] were cultured at 28 °C in medium consisting of equal volumes of Mitsuhashi/ Maramorosh (Bioconcept, Switzerland) and Schneider's insect medium (Sigma, France), supplemented with 10% (v/v) of heat-inactivated fetal bovine serum (PAA, USA) and penicillin/streptomycin (50 U/50 μg/mL; Gibco, Invitrogen, France). Cells were continuously passaged in 25 -cm² flasks by scrapping and seeding a new flask with 1:5 of the cell suspension in 5 mL of fresh medium, every 4 days. The Chikungunya virus (CHIKV) 06.21 strain was isolated in C6/36 from newborn serum sample with neonatal encephalopathy during the outbreak in La Reunion Island [6]. Viral stocks were produced on C6/36 cells in 25-cm2 flasks, at Multiplicity Of Infection (MOI) of 0.01. After 3 days at 28 °C, supernatants from infected cells were recovered and virus titration was done using plaque assay on Vero E6 (green monkey kidney) cells [40]. The titer stock virus was estimated to $10⁸$ plaque-forming units (PFU)/mL and stored in aliquots at −80 °C until used.

Cell infection

To assess the impact of cell co-infection by Wolbachia and CHIKV, we compared four modalities of infection; cells uninfected, mono-infected by wAlbB or CHIKV and bi-infected, each with three independent biological replicates. The day prior infection, 5×10^6 cells were transferred in 25 - $cm²$ flask and allowed to attach for 18 h at 28 °C. Infection at MOI 0.1 with CHIKV 06.21

was performed in 0.5 mL new medium with 2% fetal bovine serum, using virus-free medium as control. After 1 h, 5 mL of fresh medium with 10% fetal bovine serum were added and incubation extended. Cells and supernatants were harvested at 24 and 120 h post-infection. For uninfected cells, we applied the same protocol but fetal bovine serum medium did not contain any virus particles. Blue trypan staining used for cell counting and light microscopy employed to monitor cell monolayers did not show apparent necrotic cells along the course of the experiment (not shown). At the two times (24 h and 120 h), cells were scrapped and pelleted by centrifugation and a fraction of these cells was conserved in 1.5 mL tube for genomic DNA and RNA isolations. Each cell pellet was washed once in 10 mL PBS 1× pH 7.4 (Gibco, Invitrogen, France) and then resuspended in lysis buffer composed of urea 7 M (Sigma, France), 2 M thiourea (Fluka, Sigma, France), 4% CHAPS (Sigma, France), 0.5% Triton ×100 (Sigma, France) and TBP 0.08 mM (Sigma, France) in distilled water (Gibco, Invitrogen, France); and incubated on ice for 30 min with regular vortexing. Cell lysates were stored at −80 °C until protein extractions.

DNA and RNA isolation

Genomic DNA isolation was performed using DNeasy blood and tissues Kit (Qiagen, France) following manufacturer's instructions. Cell pellets were resuspended in 180 μL of ATL lysis buffer and incubated for 2 h at 37 °C with 2 mg/mL lysozyme (Euromedex, France). Residual co-extracted RNA was eliminated by adding 100 mg/mL RNase A, for 2 min at room temperature, then isolated DNA was eluted in 30 μL of DNase-free water. To isolate total RNA, cell pellets were crushed in 350 μL RLT lysis buffer of RNeasy Mini Kit (Qiagen, France) using RNasefree piston pellet (Kontes, USA) and following manufacturer's recommendations. Then RNA was eluted in 37 μL of RNase-free water and treated with DNase using the TURBO-DNA free kit (Ambion, USA) in 50 μL final volume following the manufacturer's instructions. DNA and RNA were quantified using a UV-mc² spectrophotometer and diluted to 5 ng/μL, then frozen at −20 °C (DNA) or −80 °C (RNA) until use.

Quantitative analysis of Wolbachia (qPCR) and CHIKV (RT-qPCR)

To monitor the relative density of Wolbachia per cell, qPCR was performed using Wolbachia Surface Protein (*wsp*) gene for the bacterium and *actin* gene for the host cell. Standard curves were drawn on 10-fold serial dilutions from 1×10^8 to 1×10^1 copies/μL of the DNA plasmid pQuantAlb16S containing fragments of the two targeted genes [20, 41]. Amplification reaction was done in a total volume of 20 μL containing 10 ng of template DNA, 1× (10 μL) Fast-SYBR-Green Master Mix (Roche, Suisse), 200 mM of each wsp primers (5'AAGGAACC GAAGTTCATG3′ and 5'AGTTGTGAGTAAAGTCCC3') and 300 mM each actin primers (5'GCAAACGTGG TATCCTGAC3' and 5'GTCAGGAGAACTGGGTGCT3'). Amplification was performed on LC480 LightCycler (Roche, France) and consisted of 10 min at 95 °C, followed by 40 cycles of 15 s at 95 °C, 1 min at 65 °C, and a final elongation at 72 °C for 30 s. To quantify CHIKV RNA copy number, RT-qPCR was done on the envelope E2 gene using a standard curve of 10-fold serial dilution of a synthetic CHIKV RNA transcript [29]. One-step RTqPCR was performed using EXPRESS One-Step SYBR GreenER Kit (Invitrogen, France) in a volume of 20 μL containing 10 ng of RNA template, $1 \times (10 \mu L)$ EXPRESS SYBR GreenER SuperMix Universal, 200 nM of sense Chik/E2/9018/+ and anti-sense Chik/E2/9235/− primers [42] and $1 \times (0.5 \mu L)$ EXPRESS Superscript Mix. Amplification was performed on a LC480 LightCycler (Roche, France) and consisted of 15 min at 50 °C and by 95 °C for 2 min, followed by 40 cycles of 95 °C for 15 s and 63 °C for 1 min. All PCR reactions were done in triplicate. DNA and RNA extracted from C6/36 uninfected were used as negative control.

Protein extraction, 2D–PAGE and densitometric gel analyses

To extract proteins, cell lysates were defreezed on ice and proteins were precipitated with 10% (w/v) trichloroacetic acid (Sigma, France) at 4 °C overnight. Proteins were pelleted by centrifugation at 14,000 g for 15 min at 4 °C and washed three times with glacial acetone (VWR Chemicals, France). Isoelectric focusing (IEF) was performed using the Protean IEF System (Biorad, France) according to the manufacturer's instructions. The rehydration buffer contained 8 M urea (Sigma-Aldrich) and 4% (w/v) CHAPS (Sigma). IEF was performed with 11 cm no-linear strips, pH 3–10 (Biorad), using the Voltage Ramp protocol recommended by the manufacturer (100 V/30 min/rapid, 250 V/30 min/linear, 1000 V/30 min/linear, 7000 V/ 3 h/linear, and finally 32,000 V/h (pH 3–10 IPG)). The second dimension was carried out using the Criterion Dodeca system (Biorad). A minimum of four gels loaded with biological replicates was used for each condition. Criterion any kD TGX gels (Biorad) were run at 10 °C in Laemmli buffer [43] at 100 V for 2 h. Then the 2D–gels were stained with silver nitrate as previously described [44], scanned and analyzed using the software SameSpots v.4.5 (Non-linear Dynamics Progenesis, UK). An ANOVA test of the spot volumes was calculated to compare the different conditions. Variations in spot volumes with $p < 0.02$ and fold-change >2 were considered significant.

Sample preparation and nanoLC-MS/MS analysis

Protein spots were destained in 60 mM potassium ferricyanide and 200 mM sodium thiosulfate mixed 1:1 until all brown color was removed. The spots were washed through successive incubations with water until all yellow color was removed and shrunk in acetonitrile (ACN) for 10 min. After ACN removal, gel pieces were dried at room temperature. Proteins were digested by incubating each gel slice with 10 ng/μL of trypsin (T6567, Sigma-Aldrich) in 40 mM NH4HCO3, 10% ACN, rehydrated at 4 °C for 10 min, and finally incubated overnight at 37 °C. The resulting peptides were extracted from the gel by three steps: a first incubation in 40 mM NH4HCO3, 10% ACN for 15 min at room temperature followed by two incubations in 47.5% ACN, 5% formic acid for 15 min at room temperature. The three collected extractions were pooled with the initial digestion supernatant, dried in a SpeedVac, and resuspended with 25 μL of 0.1% formic acid before nanoLC-MS/MS analysis. Online nanoLC-MS/MS analyses were performed using an Ultimate 3000 RSLC Nano-UPHLC system (Thermo Scientific, USA) coupled to a nanospray Q-Exactive hybrid quadruplole-Orbitrap mass spectrometer (Thermo Scientific, USA). Ten microliters of each peptide extract were loaded on a 300 μm ID \times 5 mm PepMap C18 precolumn (Thermo Scientific, USA) at a flow rate of 20 μ L/min. After 5 min desalting, peptides were online separated on a 75 μ m ID \times 25 cm C18 Acclaim PepMap® RSLC column (Thermo Scientific, USA) with a 4–40% linear gradient of solvent B (0.1% formic acid in 80% ACN) in 48 min. The separation flow rate was set at 300 nL/min. The mass spectrometer operated in positive ion mode at a 1.8 kV needle voltage. Data were acquired using Xcalibur 3.0 software in a data-dependent mode. MS scans (m/z 300–2000) were recorded at a resolution of $R = 70,000$ (@ m/z 200) and an AGC target of 1×10^6 ions collected within 100 ms. Dynamic exclusion was set to 30 s and top 15 ions were selected from fragmentation in HCD mode. MS/MS scans with a target value of 1×10^5 ions were collected with a maximum fill time of 120 ms and a resolution of $R = 35,000$. Additionally, only +2 and +3 charged ions were selected for fragmentation. Others settings were as follows: no sheath and no auxiliary gas flow, heated capillary temperature, 200 °C; normalized HCD collision energy of 25% and an isolation width of 3 m/z.

Database search and results processing

Mascot, MS Amanda and Sequest algorithms through Proteome Discoverer 1.4 Software (Thermo Fisher Scientific Inc., USA) were used for protein identification in batch mode by searching against a merged database from [http://www.uniprot.org/:](http://www.uniprot.org/) Aedes (taxon identifier: [7158], 24,927 entries, release 2015_04) + Wolbachia (taxon identifier: [952], 24,150 entries, release 2015_04) + Chikungunya virus (taxon identifier: [37124], 2041 entries, release 2015_04) + Dengue virus (taxon identifier [12637], 13,782 entries, release 2015_04). Two missed enzyme cleavages were allowed. Mass tolerances in MS and MS/MS were set to 10 ppm and 0.02 Da. Oxidation of methionine, acetylation of lysine and deamination of asparagine and glutamine were searched as dynamic modifications. Carbamidomethylation on cysteine was searched as static modification. Peptide validation was performed using Target Decoy PSM Validator and only "high confidence" peptides were retained corresponding to a 1% False Positive Rate at peptide level. The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium ([http://proteomecentral.](http://proteomecentral.proteomexchange.org) [proteomexchange.org](http://proteomecentral.proteomexchange.org)) via the PRIDE partner repository [45] with the dataset identifier PXD005091.

Bioinformatics and statistical analysis

The continuous response variables (viral and bacterial titers) were log_{10} -transformed. They were analyzed using a multifactorial linear model, with a normal error distribution and an identity link function that included the effect of the time and MOI as ordinal variables, treatment as discrete variable and their interactions. All the statistical analyses were performed using R environment (version 3.1.0). An annotation in GO term was carried out on the proteins identified using Blast2GO (3.2.7) then they were used to detect possible interaction networks using Cytoscape (3.3.0).

Results and discussion

Wolbachia wAlbB affects CHIKV in cellulo

As our previous study of the C6/36 infected with wAlbB showed that presence of the bacterium decreased the viral titer compared to uninfected cells [35], we measured wAlbB and CHIKV densities at 24 and 120 h post infection (p.i.) using qPCR and RT-qPCR, respectively. The density of *Wolbachia* was about 12 *wsp* gene/*actin* ratio (Fig. 1). The percentage of Wolbachia-infected cells ranged from 60 to 70% (not shown) as determined by fluorescent in situ hybridization published protocol [35]. The CHIK RNA copy number was estimated between $10⁷$ to $10⁹$ per ng of total RNA (Fig. 2). Both *Wolbachia*infected and uninfected cells produce infectious viral particles without visible cytopathic effect (not shown). This was expected as *Aedes* cells are permissive to many arboviruses, including CHIKV, that are found non pathogenic to mosquitoes [46, 47]. This is why the C6/ 36 cell line is extensively used to propagate viruses [48].

Statistical analyzes demonstrated that the density of Wolbachia was not affected by the presence of the CHIKV, and was marginally affected upon time $(P = 0.05262)$ (Fig. 1). As expected, the viral titer was

significantly reduced in the presence of Wolbachia $(P < 2.2e-16)$, without reaching complete inhibition. The inhibitory effect decreased with time, being lower at a late time $(P = 0.0007825)$ (Fig. 2). It has been reported that viral inhibition by Wolbachia is density-dependent [28, 37]. At the two time points tested here the Wolbachia density remained stable, around 12 bacteria per cell, and the level of CHIKV inhibition was similar to previous studies [35]. The chronic Wolbachia infection and the permissiveness to viruses make the C6/36 cell line an interesting model for exploratory functional studies. One unfavorable point of this cell line is the lack of siRNA pathway [49], a primary immune response

against viral infection in mosquitoes. However, it has been shown that insects can mobilize other RNA interference pathways to control viral replication. For instance, Aedes aegypti induces miRNA and specific piRNA pathways to control the replication of DENV [50–52]. Similarly, Wolbachia could have an effect on synthesis of small RNAs [53, 54]. Therefore, this cellular model seems suitable for the study of induced host-cell responses following mono- or bi-partite infection by Wolbachia and/or CHIKV as well as the CHIKV replication cycle.

Differential cell proteome profiles upon microbial infection

For the two time points (24 h and 120 h p.i.) and the four modalities (uninfected, mono-infected by either Wolbachia or CHIKV and bi-infected by both microbes), three independent biological replicates were performed. Total proteins were extracted and similar amounts (approximately 150 μg, estimated on a 1D gel) were used for 2DE. For each modality and each replicate, a minimum of 4 and a maximum of 5 gels were used. Typical 2D gels with spots obtained are illustrated in Fig. 3. The global gel analysis using the ProGenesis SameSpots software enabled detection of 906 spots at 24 h and 901 spots at 120 h p.i. ANOVA analysis allowed identifying 58 spots at 24 h and 32 spots at 120 h p.i that were statistically different ($p < 0.02$ and fold change >2) in comparison to uninfected cells. As many of the spots identified at early time point were linked to Wolbachia infection alone, only 30 of the 58 spots were selected for mass spectrometry sequencing, including all 32 spots observed at the late time point.

A protein was considered present in a spot when a minimum of two different peptides were identified by mass spectrometry (Additional file 1: Table S1). Consequently, a total of 495 unique proteins were identified from 948 sequences at 24 h p.i., whereas 105 unique proteins were found among 168 sequences at 120 h p.i. The elevated number of identified sequences in the analysis can be explained by two major reasons; (i) a high number of proximate proteins that have possibly been subjected to post-translational modifications and (ii) protein fragmentation during experimentation that resulted in modified migration patterns. All peptide sequences and observed fold changes are described on the Additional file 1: Table S1. By combining the protein level in each time point and the modality of infection, a total of four major profiles were defined, including monoinfection, dominance, cumulative and interference (Table 1). Accordingly, in the monoinfection profile each microbial partner tends to affect a particular protein or a group of host proteins. The dominant profile indicates a major impact of one microbial partner on the host

protein synthesis (up or down) whereas the other microbial partner showed an opposite profile. The cumulative effect means that the two microbial partners displayed a synergic effect on protein synthesis. Lastly, the interference profile indicates that each microbial partner induces a specific protein pattern but the co-infection displays a totally new trend.

The 2DE combined with mass spectrometry sequencing did not allow quantification of the level of protein accumulation per spot, and one spot can contain several proteins, consequently it was not possible to identify which protein was involved in the variation observed. In addition, the presence of many identical proteins in several spots simultaneously makes the analysis complex.

Therefore we proceeded by annotating proteins in GO terms that were used to construct interacting networks for each protein profile. This procedure allowed comparison of functions shared by all modalities with those belonging specifically to each partner.

Wolbachia infection has a greater effect on cell functions than CHIKV

Among the 89 spots detected, 77% were specifically synthesized in the presence of Wolbachia in both monoinfection (53 spots) and dominance (16 spots) profiles (Table 1). Since the aim of this study was to characterize the impact of coinfection rather than monoinfection, we have chosen to sequence only 27 out of 53 spots, that

Table 1 Profiling of protein accumulation

All profiles were normalized with respect to uninfected modality. Effective observed fold changes are reported on Additional file 1: Table S1. In comparison to uninfected C6/36 cells: Up: A positive difference on protein synthesis has been observed; Down: A negative difference on proteins synthesis has been observed; ø: No difference has been observed

were selected on the basis of a particular fold change as indicated above. Results of sequencing showed that spots linked to Wolbachia in both mono-infection and dominance profiles contained proteins involved in many cellular functions, including processes related to metabolism for acquisition of resources from the host, regulation of anti-oxidation and cellular functional machinery (transcription and translation), as well as active transport and cellular structures (Fig. 4). These proteins were present at the two times at relatively high percentage (64.5% at early time and 35% at late time) of the total proteins, and some of them have been already described in literature as being upregulated by Wolbachia [39]. One example is the Glutathione S-transferase (A0A023EL34) for the regulation of anti-oxidation process [38], which is abundant at early time in the presence of the bacterium. The large number of proteins mobilized in the presence of Wolbachia indicates a strong relationship between the two partners.

In contrast, the presence of CHIKV alone has only limited effect in comparison to uninfected cells. Few differential spots containing proteins at a very low percentage (<5%) were detected, with tendency to be down-regulated. The majority of the proteins detected were related to the ATP transport and binding, glycolysis,

cytoskeleton and stress responses (Fig. 4). For instance, many proteins associated with ATP consumption were significantly reduced in the presence of CHIKV. Moreover, we observed a decrease in expression of the gene encoding A0A023END7 LSD2 (Lipid Storage Droplet-2), suggesting that CHIKV blocks lipid storage, potentially making them available incorporation into the viral envelope. This phenomenon has already been shown in Ae. aegypti mosquitoes infected by either dengue [55] or chikungunya viruses [14, 15]. Another protein A0A023EQG9 negatively impacted encoded a kinase for double-stranded RNA necessary to the establishment of RISC complex in RNA interference phenomenon. Knowing that the C6/36 cell line has a non-functional siRNA mechanism [49], inhibition of the miRNA pathway is consistent with a viral mechanism to escape cellular defenses.

In dominance profiles, Wolbachia exhibited different protein trends in respect to virus, from a neutral level (W_DOM_1 and W_DOM_3), an increased (W_DOM_2) or repressed (W_DOM_4) synthesis (Fig. 5). The W_DOM_1 profile reduced the CHIKV structural polyprotein V5UMV1 at 24 h post infection (hpi)., whereas the W_DOM_4 profile targeted specifically the viral capsid protein (A0A059VQ68) at 120 hpi (Table 1). These results are in agreement with the *Wolbachia* blocking

phenotype observed recently for CHIKV in C6/36 cells [35]. Viral blocking is therefore explained by inhibition of Wolbachia cellular proteolysis machinery, thus limiting the maturation of virion-associated protein structures and reducing viral replication. Overall, this effect appeared more diverse at early stages post-infection, but of greater magnitude at later times (Table 1). At 24 hpi Wolbachia tends to sustain necessary cellular processes, such as oxidizing processes including glutathione peroxidase activity, translation and transcription (Fig. 5). Whereas, this is not the case at 120 hpi, when the bacterium limits processes that will be exploited by the virus, including oxidative stress, transportation and translation.

The viral dominance profile occurred at 120 hpi, when the virus had established chronic and dense infection (Fig. 6). The ATP synthase subunit beta (A0A023ETB9), involved in active trans-membrane ion transport, appeared negatively regulated as well as Glutathione peroxidase (Q16N54), albeit to a lesser extent. In contrast, some structural proteins such as actin (Q0Z987) and those related to heat shock (A0A023EWK8) were oversynthesized, suggesting a role in the production of virions [17]. The presence of ATP synthase subunit beta in both up and down-regulated profiles suggests several isoforms of this protein that Wolbachia modulates by regulating post-translational modifications.

Proteome trends during Wolbachia and CHIKV coinfection

Two different profiles emerged from bacterial and viral coinfection. The first was a cumulative profile in which a synergistic negative effect on protein synthesis was observed (Fig. 7). The processes observed to be affected by bi-infection were those already identified during infection by bacteria and viruses [14, 15, 17]. These proteins all act to maintain cell integrity and are associated with either down-regulation early post infection or upregulation late post infection.

The second pattern was an interference profile (Fig. 8). At 24 hpi, interference seemed to be directed against CHIKV and in favour of Wolbachia. Indeed, despite

Wolbachia neutral (INT_2) or negative (INT_1) effects, cellular processes that were found to be up-regulated were those that may be of benefit to the bacterium, including cell development processes, transcription, translation and various metabolic pathways. At 120 hpi the INT_3 profile showed establishment of a balance between Wolbachia, which decreases metabolic processes, and the virus who in turn activated them for its own benefit. This sum of effect allows maintaining these processes at a steady-state level in cell. INT_4 profile was essentially related to structural proteins that were inhibited by each microbial partner, but during bi-infection where these proteins were not downregulated. INT_5 profile identified ATP synthase subunit beta of Wolbachia (H0U0S7) that was inhibited by the virus. This later profile highlights a particular pattern where the presence of virus inhibited bacterial proteins through the blocking access to resources, thus limiting the potential of the bacterium to affect the virus.

When comparing the peptides detected in this study with those already described in mono-infection models using mass spectrometry approaches, some common proteins were identified. These include enolase (A0A023ETA6), which was found to stimulate transcription of the Sendai virus genome [56], and upregulated in the interference profile (INT_3). If CHIKV seems to enhance the enolase synthesis, as already shown by Lee et al. [17], Wolbachia tends to reduce its production. Consequently, in the biinfection status, this conflictive pattern appears unfavorable to CHIKV replication. Among proteins involved in glycolysis and metabolism, one promising candidate is the disulfide isomerase protein (A0A023EP23) which has been shown to be modulated by CHIKV according to the infected organs [14, 15] and the duration of infection [17]. In our study, this protein is modulated by Wolbachia (W_Up_1), affecting the early CHIKV replication. Similarly, some chaperonins such as the putative calreticulinlike 2 (A0A023EQL3), chaperonin 60 kDa (A0A023EV59), heat shock cognate 70 (Q1HQZ5), alpha and beta tubulin 1 (A0A023ERN1 and A0A023ESE6) have been described to be modulated during CHIKV infection [14, 15, 17], and for which we found to be impacted by bi-infection status. These observations are also operating in glycolysis with for instance triosephosphate isomerase (A0A023EIM8) shown to be important in energy input necessary for viral replication. Indeed, at early time, this protein is overexpressed in Wolbachia-infected cells, inducing a favorable environment for CHIKV. In contrast, at latter time, Wolbachia seems to reduce the expression of triosephosphate isomerase while CHIKV tends to increase its activity (V_DOM_3 profile), suggesting the importance of such protein in this tripartite interaction.

Conclusions

This study highlights complex processes that occur during arbovirus infection of mosquito cells in symbiosis with Wolbachia. Even though these findings were obtained using a cellular model, the observed trends pave the way for future research into the in vivo characteristics of tripartite interaction. In our experimental conditions, the combination of 2DE and nanoLC-MS/MS revealed a balance in protein synthesis mostly in favor of Wolbachia, which may explain the simultaneous inhibition of viral replication that we observed using RTqPCR. At early times post infection, the presence of Wolbachia greatly influences many cellular processes related to management of anti-oxidant activity, protein production, various metabolic pathways linked to the provisioning of resources; likely impacting CHIKV replication. Under such conditions, CHIKV faces a hostile environment for replication and appears to counterbalance this negative impact by blocking some key cellular pathway, including the inhibition of transcription, translation and locking of an miRNA pathway.

At later times post infection, the proteome is clearly altered, and CHIKV activity seems to have taken control of some cellular functions. Consequently, the virus seems to

limit the impact of Wolbachia on its replication cycle by hoarding the majority of resources, and even blocking Wolbachia's access to these resources. This shift partially explains the increased viral titer that is observed at later periods post-infection. Even if Wolbachia no longer controls some of these cellular processes, its presence limits the effect of CHIKV infection on certain cellular functions, thus modulating its replication, particularly early after the infection process. This cellular level interference could explain phenotypes observed in Ae. albopictus in vivo, where Wolbachia limits transmission of dengue virus by reducing the viral titer in salivary glands [27].

Several studies have shown that Wolbachia can modulate the expression of genes involved in immunity that affect arbovirus infection, suggesting that interference acts by pre-immunization of the host [26, 28, 34]. Strikingly,

proteins was related to this function

we do not observe significant modulation of proteins related to immune response upon CHIKV-inhibition by Wolbachia. Even though C6/36 lacks functional siRNA pathways, other immune response mechanisms could have been mobilized. The fact that we did not identify proteins involved in immunity might suggest that other cellular processes can lead to the antiviral profile, corroborating results obtained from other cellular models. For example, in Ae. albopictus Aa23 cells infected with either wAlbB, wMel or wMelPop, whose density varied from 2.5

to 38 bacteria per host cell, no changes were observed in innate immunity related functions [57]. Recently, an elegant work demonstrated that Wolbachia could inhibit viral replication at early stages post infection by affecting RNA translation or transcription, suggesting a likely direct effect [58]. Together these cellular models revealed alternative mechanisms to immunity in Wolbachia-based viral inhibition that need further investigations. An interesting perspective could be the extension of proteome profiling to mosquito organs as

well as testing other Ae. albopictus-transmitted arboviruses, such as Dengue and Zika, with emphasis on functions revealed in this study.

Additional file

[Additional file 1: Table S1.](dx.doi.org/10.1186/s12864-017-3985-y) List of spots and fold change with sequences of peptides. (XLSX 7247 kb)

Abbreviations

ATP: Adenine triphosphate; CHAPS: 3-[(3-cholamidopropyl) dimethylammonio]- 1-propanesulfonate; CHIKV: Chikungunya virus; FBS: Foetal bovine serum; GO: Gene ontology; IEF: Isoelectric focusing; IPG: Immobilized pH gradient; LC-MS/MS: Liquid chromatography coupled to tandem mass spectrometry; MOI: Multiplicity of infection; p.i.: Post-infection; PBS: Phosphate buffered saline; TBP: Tributylphosphin; TCA: Trichloroacetic acid; TET: Tetracycline; wAlb_B: Wolbachia B

Acknowledgements

This paper is dedicated to the memory of Florence-Hélène Tran who tragically left us in August 2016. We acknowledge the contribution of the BSL3 platform of SFR BioSciences Gerland Lyon Sud (UMS3444/US8), the DTAMB platform of the FR41 Bio-Environment and Health (University Lyon 1), and IBio platform of Microbial Ecology laboratory (UMR5557).

Funding

This work was partially supported by the Region Rhône-Alpes and by ERA-NET BiodivERsA with the national funders ANR-13-EBID-0007-01, FWF I-1437, and DFG KL 2087/6–1 as part of the 2012–2013 BiodivERsA call for research proposals.

Availability of data and materials

All data and analysis are fully available within the paper and its additional files as well as at PRIDE database under project accession: PXD005091. Username: reviewer24708@ebi.ac.uk. Password: F8aBJWI5.

Authors' contributions

YS, CVM and PM conceived the project and the sampling design. YS, CD, JWD, FHT, VG and PP performed the experimental work. YS performed the bioinformatics and statistical analyses. YS, CVM and PM wrote the first draft of the manuscript. All the authors contributed to the manuscript revision. All authors read and approved the final manuscript.

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

1 Université de Lyon, Lyon, France. ²Université Lyon 1, Villeurbanne, France.
³CNPS LIMP5557, Ecologio Microbionne, Villeurbanne, France. ⁴INPA. CNRS, UMR5557, Ecologie Microbienne, Villeurbanne, France. ⁴INRA, UMR1418, Villeurbanne, France. ⁵Laboratoire Mixte UMR 5240, Plateforme de Protéomique, CNRS, Lyon, France. ⁶Centre de Génomique Fonctionnelle, Plateforme Protéome, Université Bordeaux, F-33000 Bordeaux, France. ⁷CNRS 9192, INSERM U1187, IRD 249, Unité Mixte Processus Infectieux en Milieu Insulaire Tropical (PIMIT). Plateforme Technologique CYROI, Université de La Réunion, 2 rue Maxime Rivière, 97490 Sainte Clotilde, île de La Réunion, France.

Received: 5 April 2017 Accepted: 1 August 2017 Published online: 18 August 2017

References

- 1. IUCN/SSC Invasive Species Specialist Group. ISSG Database: Ecology of Aedes albopictus [Internet]. Global Invasive Species Database. [cited 2016 Feb 15]. Available from: [http://www.issg.org/database/species/ecology.asp?si=](http://www.issg.org/database/species/ecology.asp?si=109&fr=1&sts=&lang=EN) [109&fr=1&sts=&lang=EN](http://www.issg.org/database/species/ecology.asp?si=109&fr=1&sts=&lang=EN).
- 2. Benedict MQ, Levine RS, Hawley WA, Lounibos LP. Spread of the tiger: global risk of invasion by the mosquito Aedes Albopictus. Vector Borne Zoonotic Dis Larchmt N. 2007;7(1):76–85.
- 3. Gratz NG. Critical review of the vector status of Aedes Albopictus. Med Vet Entomol. 2004 Sep 1;18(3):215–27.
- 4. Chouin-Carneiro T, Vega-Rua A, Vazeille M, Yebakima A, Girod R, Goindin D, et al. Differential susceptibilities of Aedes Aegypti and Aedes Albopictus from the Americas to Zika virus. PLoS Negl Trop Dis. 2016;10(3):e0004543.
- 5. Grard G, Caron M, Mombo IM, Nkoghe D, Mboui Ondo S, Jiolle D, et al. Zika virus in Gabon (Central Africa)–2007: a new threat from Aedes Albopictus? PLoS Negl Trop Dis. 2014;8(2):e2681.
- 6. Schuffenecker I, Iteman I, Michault A, Murri S, Frangeul L, Vaney M-C, et al. Genome Microevolution of Chikungunya Viruses Causing the Indian Ocean Outbreak. PLoS Med [Internet]. 2006;3(7) [cited 2015 Nov 30];. Available from: [http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1463904/.](http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1463904/)
- 7. Gérardin P, Guernier V, Perrau J, Fianu A, Le Roux K, Grivard P, et al. Estimating Chikungunya prevalence in la Réunion Island outbreak by serosurveys: two methods for two critical times of the epidemic. BMC Infect Dis. 2008;8:99.
- 8. Rezza G, Nicoletti L, Angelini R, Romi R, Finarelli AC, Panning M, et al. Infection with chikungunya virus in Italy: an outbreak in a temperate region. Lancet Lond Engl. 2007;370(9602):1840–6.
- 9. Delisle E, Rousseau C, Broche B, Leparc-Goffart I, L'Ambert G, Cochet A, et al. Chikungunya outbreak in Montpellier, France, September to October 2014. Euro Surveill Bull Eur Sur Mal Transm Eur Commun Dis Bull. 2015;20(17). <http://www.eurosurveillance.org/ViewArticle.aspx?ArticleId=21108>.
- 10. Roiz D, Boussès P, Simard F, Paupy C, Fontenille D. Autochthonous Chikungunya transmission and extreme climate events in southern France. PLoS Negl Trop Dis. 2015;9(6):e0003854.
- 11. Succo T, Leparc-Goffart I, Ferré J-B, Roiz D, Broche B, Maquart M, et al. Autochthonous dengue outbreak in Nîmes, South of France, July to September 2015. Euro Surveill Bull Eur Sur Mal Transm Eur Commun Dis Bull. 2016;21(21). [http://www.eurosurveillance.org/ViewArticle.aspx?ArticleId=](http://www.eurosurveillance.org/ViewArticle.aspx?ArticleId=22485) 22485
- 12. Grandadam M, Caro V, Plumet S, Thiberge JM, Souarès Y, Failloux A-B, et al. Chikungunya virus, southeastern France. Emerg Infect Dis. 2011;17(5):910–3.
- 13. Tsetsarkin KA, Vanlandingham DL, McGee CE, Higgs S. A single mutation in Chikungunya virus affects vector specificity and epidemic potential. PLoS Pathog. 2007;3(12):e201-1906.
- 14. Tchankouo-Nguetcheu S, Bourguet E, Lenormand P, Rousselle J-C, Namane A, Choumet V. Infection by chikungunya virus modulates the expression of several proteins in Aedes Aegypti salivary glands. Parasit Vectors. 2012;5:264.
- 15. Tchankouo-Nguetcheu S, Khun H, Pincet L, Roux P, Bahut M, Huerre M, et al. Differential Protein Modulation in Midguts of Aedes aegypti Infected with Chikungunya and Dengue 2 Viruses. PLoS ONE [Internet]. 2010;5(10) [cited 2016 Mar 17]. Available from: [http://www.ncbi.nlm.nih.gov/pmc/articles/](http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2950154/) [PMC2950154/](http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2950154/).
- 16. Lee RCH, Hapuarachchi HC, Chen KC, Hussain KM, Chen H, Low SL, et al. Mosquito cellular factors and functions in mediating the infectious entry of chikungunya virus. PLoS Negl Trop Dis. 2013;7(2):e2050.
- 17. Lee RCH, Chu JJH. Proteomics Profiling of Chikungunya-Infected Aedes albopictus C6/36 Cells Reveal Important Mosquito Cell Factors in Virus Replication. PLoS Negl Trop Dis [Internet]. 2015 Mar 4 [cited 2016 Feb 15];9(3). Available from:<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4349740/>.
- 18. Patramool S, Surasombatpattana P, Luplertlop N, Sévéno M, Choumet V, Thomas F, et al. Proteomic analysis of an Aedes Albopictus cell line infected with dengue serotypes 1 and 3 viruses. Parasit Vectors. 2011;4:138.
- 19. Kittayapong P, Baisley KJ, Baimai V, O'Neill SL. Distribution and diversity of Wolbachia infections in southeast Asian mosquitoes (Diptera: Culicidae). J Med Entomol. 2000;37(3):340–5.
- 20. Zouache K, Voronin D, Tran-Van V, Mousson L, Failloux A-B, Mavingui P. Persistent Wolbachia and Cultivable Bacteria Infection in the Reproductive and Somatic Tissues of the Mosquito Vector Aedes albopictus. PLoS ONE

[Internet]. 2009;4(7) [cited 2015 Dec 1]. Available from: [http://www.ncbi.nlm.](http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2712238/) [nih.gov/pmc/articles/PMC2712238/.](http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2712238/)

- 21. Minard G, Tran FH, Van VT, Goubert C, Bellet C, Lambert G, et al. French invasive Asian tiger mosquito populations harbor reduced bacterial microbiota and genetic diversity compared to Vietnamese autochthonous relatives. Front Microbiol. 2015;6:970.
- 22. Yen JH, Barr AR. New hypothesis of the cause of cytoplasmic incompatibility in Culex Pipiens L. Nature. 1971 Aug 27;232(5313):657–8.
- 23. Dobson SL, Rattanadechakul W, Marsland EJ, Fitness advantage and cytoplasmic incompatibility in Wolbachia single- and superinfected Aedes Albopictus. Heredity. 2004;93(2):135–42.
- 24. Xi Z, Khoo CCH, Dobson SL. Wolbachia establishment and invasion in an Aedes Aegypti laboratory population. Science. 2005;310(5746):326–8.
- 25. Bian G, Xu Y, Lu P, Xie Y, Xi Z. The endosymbiotic bacterium Wolbachia induces resistance to dengue virus in Aedes Aegypti. PLoS Pathog. 2010;6(4):e1000833.
- 26. Pan X, Zhou G, Wu J, Bian G, Lu P, Raikhel AS, et al. Wolbachia induces reactive oxygen species (ROS)-dependent activation of the toll pathway to control dengue virus in the mosquito Aedes Aegypti. Proc Natl Acad Sci U S A. 2012;109(1):E23–31.
- 27. Mousson L, Zouache K, Arias-Goeta C, Raquin V, Mavingui P, Failloux A-B. The native Wolbachia symbionts limit transmission of dengue virus in Aedes Albopictus. PLoS Negl Trop Dis. 2012;6(12):e1989.
- 28. Blagrove MSC, Arias-Goeta C, Failloux A-B, Sinkins SP. Wolbachia strain wMel induces cytoplasmic incompatibility and blocks dengue transmission in Aedes Albopictus. Proc Natl Acad Sci U S A. 2012;109(1):255–60.
- 29. Mousson L, Martin E, Zouache K, Madec Y, Mavingui P, Failloux AB. Wolbachia modulates Chikungunya replication in Aedes Albopictus. Mol Ecol. 2010;19(9):1953–64.
- 30. Hussain M, Torres S, Schnettler E, Funk A, Grundhoff A, Pijlman GP, et al. West Nile virus encodes a microRNA-like small RNA in the 3' untranslated region which up-regulates GATA4 mRNA and facilitates virus replication in mosquito cells. Nucleic Acids Res. 2012;40(5):2210–23.
- 31. Hughes GL, Vega-Rodriguez J, Xue P, Rasgon JL. Wolbachia strain wAlbB enhances infection by the rodent malaria parasite plasmodium berghei in Anopheles Gambiae mosquitoes. Appl Environ Microbiol. 2012;78(5):1491–5.
- 32. Johnson KN. The impact of Wolbachia on virus infection in mosquitoes. Viruses. 2015;7(11):5705–17.
- 33. Weiss B, Aksoy S. Microbiome influences on insect host vector competence. Trends Parasitol. 2011;27(11):514–22.
- 34. Moreira LA, Iturbe-Ormaetxe I, Jeffery JA, Lu G, Pyke AT, Hedges LM, et al. A Wolbachia symbiont in Aedes Aegypti limits infection with dengue, Chikungunya, and plasmodium. Cell. 2009;139(7):1268–78.
- 35. Raquin V, Valiente Moro C, Saucereau Y, Tran F-H, Potier P, Mavingui P. Native Wolbachia from Aedes albopictus Blocks Chikungunya Virus Infection In Cellulo. PLoS ONE [Internet]. 2015;10(4) [cited 2015 Nov 30]. Available from: [http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4414612/.](http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4414612/)
- 36. Frentiu FD, Robinson J, Young PR, McGraw EA, O'Neill SL. Wolbachia-mediated resistance to dengue virus infection and death at the cellular level. PLoS One. 2010;5(10):e13398.
- 37. Lu P, Bian G, Pan X, Xi Z. Wolbachia induces density-dependent inhibition to dengue virus in mosquito cells. PLoS Negl Trop Dis. 2012;6(7):e1754.
- 38. Brennan LJ, Keddie BA, Braig HR, Harris HL. The Endosymbiont Wolbachia pipientis Induces the Expression of Host Antioxidant Proteins in an Aedes albopictus Cell Line. PLoS ONE [Internet]. 2008;3(5) [cited 2016 Mar 17]. Available from:<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2324199/>
- 39. Baldridge GD, Baldridge AS, Witthuhn BA, Higgins L, Markowski TW, Fallon AM. Proteomic profiling of a robust Wolbachia infection in an Aedes Albopictus Mosquito cell line. Mol Microbiol. 2014;94(3):537–56.
- 40. Dubrulle M, Mousson L, Moutailler S, Vazeille M, Failloux A-B. Chikungunya Virus and Aedes Mosquitoes: Saliva Is Infectious as soon as Two Days after Oral Infection. PLoS ONE [Internet]. 2009;4(6) [cited 2015 Nov 30]. Available from: [http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2690823/.](http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2690823/)
- 41. Tortosa P, Courtiol A, Moutailler S, Failloux A-B, Weill M. Chikungunya-Wolbachia interplay in Aedes Albopictus. Insect Mol Biol. 2008;17(6):677–684.
- 42. Vazeille M, Moutailler S, Coudrier D, Rousseaux C, Khun H, Huerre M, et al. Two Chikungunya Isolates from the Outbreak of La Reunion (Indian Ocean) Exhibit Different Patterns of Infection in the Mosquito, Aedes albopictus. PLoS ONE [Internet]. 2007;2(11) [cited 2015 Dec 3]. Available from: [http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2064959/.](http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2064959/)
- 43. Laemmli UK. Cleavage of structural proteins during the assembly of the head of Bacteriophage T4. Nature. 1970;227(5259):680–685.
- 44. Catusse J, Strub J-M, Job C, Van Dorsselaer A, Job D. Proteome-wide characterization of sugarbeet seed vigor and its tissue specific expression. Proc Natl Acad Sci U S A. 2008;105(29):10262–7.
- 45. Vizcaíno JA, Côté RG, Csordas A, Dianes JA, Fabregat A, Foster JM, et al. The PRoteomics IDEntifications (PRIDE) database and associated tools: status in 2013. Nucleic Acids Res. 2013;41(Database issue):D1063–9.
- 46. Paul SD, Singh KR, Bhat UK. A study on the cytopathic effect of arboviruses on cultures from Aedes Albopictus cell line. Indian J Med Res. 1969;57(2):339–48.
- 47. Buckley SM. Susceptibility of the Aedes Albopictus and A. Aegypti cell lines to infection with Arboviruses. Proc Soc Exp Biol Med. 1969;131(2):625–30.
- 48. Igarashi A. Isolation of a Singh's Aedes Albopictus cell clone sensitive to dengue and Chikungunya viruses. J Gen Virol. 1978;40(3):531–44.
- 49. Brackney DE, Scott JC, Sagawa F, Woodward JE, Miller NA, Schilkey FD, et al. C6/36 Aedes Albopictus cells have a dysfunctional antiviral RNA interference response. PLoS Negl Trop Dis. 2010;4(10):e856.
- 50. Miesen P, Ivens A, Buck AH, van Rij RP. Small RNA profiling in dengue virus 2-infected Aedes Mosquito cells reveals viral piRNAs and novel host miRNAs. PLoS Negl Trop Dis. 2016;10(2):e0004452.
- 51. Avila-Bonilla RG, Yocupicio-Monroy M, Marchat LA, De Nova-Ocampo MA, Del Ángel RM, Salas-Benito JS. Analysis of the miRNA profile in C6/36 cells persistently infected with dengue virus type 2. Virus Res. 2017;232:139–51.
- 52. Su J, Li C, Zhang Y, Yan T, Zhu X, Zhao M, et al. Identification of microRNAs expressed in the midgut of Aedes Albopictus during dengue infection. Parasit Vectors. 2017;10(1):63.
- Zhang G, Hussain M, Asgari S. Regulation of arginine methyltransferase 3 by a Wolbachia-induced microRNA in Aedes Aegypti and its effect on Wolbachia and dengue virus replication. Insect Biochem Mol Biol. 2014;53:81–8.
- 54. Zhang G, Hussain M, O'Neill SL, Asgari S. Wolbachia uses a host microRNA to regulate transcripts of a methyltransferase, contributing to dengue virus inhibition in Aedes Aegypti. Proc Natl Acad Sci U S A. 2013;110(25):10276–81.
- 55. Samsa MM, Mondotte JA, Iglesias NG, Assunção-Miranda I, Barbosa-Lima G, Da Poian AT, et al. Dengue virus capsid protein usurps lipid droplets for viral particle formation. PLoS Pathog. 2009;5(10):e1000632.
- 56. Ogino T, Yamadera T, Nonaka T, Imajoh-Ohmi S, Mizumoto K. Enolase, a cellular Glycolytic enzyme, is required for efficient transcription of Sendai virus genome. Biochem Biophys Res Commun. 2001 Jul 13;285(2):447–55.
- 57. Molloy JC, Sinkins SP. Wolbachia do not induce reactive oxygen species-dependent immune pathway activation in Aedes Albopictus. Viruses. 2015;7(8):4624–39.
- 58. Rainey SM, Martinez J, McFarlane M, Juneja P, Sarkies P, Lulla A, et al. Wolbachia blocks viral genome replication early in infection without a transcriptional response by the Endosymbiont or host small RNA pathways. PLoS Pathog. 2016;12(4):e1005536.

Submit your next manuscript to BioMed Central and we will help you at every step:

- **•** We accept pre-submission inquiries
- **•** Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- **•** Thorough peer review
- Inclusion in PubMed and all major indexing services
- **•** Maximum visibility for your research

Submit your manuscript at www.biomedcentral.com/submit

