



HAL
open science

Bacterial diversity of cosmopolitan *Culex pipiens* and invasive *Aedes japonicus* from Germany

Sina Zotzmann, Antje Steinbrink, Kathrin Schleich, Felix Frantzmann,
Chinhda Xoumpholphakdy, Manuela Spaeth, Claire Valiente Moro, Patrick
Mavingui, Sven Klimpel

► **To cite this version:**

Sina Zotzmann, Antje Steinbrink, Kathrin Schleich, Felix Frantzmann, Chinhda Xoumpholphakdy, et al.. Bacterial diversity of cosmopolitan *Culex pipiens* and invasive *Aedes japonicus* from Germany. *Parasitology Research*, 2017, 116 (7), pp.1899-1906. 10.1007/s00436-017-5466-2 . hal-01591781

HAL Id: hal-01591781

<https://hal.univ-reunion.fr/hal-01591781v1>

Submitted on 22 Sep 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Bacterial diversity of cosmopolitan *Culex pipiens* and invasive *Aedes japonicus* from Germany

Sina Zotzmann¹ • Antje Steinbrink¹ • Kathrin Schleich¹ • Felix Frantzmann¹ • Chinhda Xoumpholphakdy¹ • Manuela Spaeth¹ • Claire Valiente Moro² • Patrick Mavingui^{2,3} • Sven Klimpel¹

Abstract Symbiotic bacteria have gained significant attention in recent years. For example, microbiota of some mosquito species seems to influence the development and transmission of pathogens. Furthermore, several attempts using bacteria as a paratransgenic tool have been made in order to assist the control of mosquito-borne diseases. In this study, we examined the bacterial diversity of wild-caught adult *Culex* (*Cx.*) *pipiens* and laboratory-reared adult *Aedes japonicus* (*Ae. japonicus*) in Germany using a culture-independent method. Genomic DNA was extracted from each specimen and submitted to PCR amplification of eubacterial 16S rDNA. After the cloning reaction, 28 bacterial transformants per sample containing the 16S rDNA inserts were selected per each sample for sequencing. The analysed specimens of *Cx. pipiens* as well as of *Ae. japonicus* showed a diverse bacterial community including some common bacterial genera. Blast analysis allowed to identify 21 bacterial genera belonging to 2 phyla among the 23 specimens of *Cx. pipiens*. The 14 analysed *Ae. japonicus* revealed 11 bacterial genera belonging to 3 phyla. In both mosquito species, identified

isolates were mainly Proteobacteria. Only 4 of the bacterial genera were found in both mosquito species, with the most prevalent genera *Sphingomonas* and *Rahnella* in *Cx. pipiens* and in *Ae. japonicus* respectively. Most of the bacterial genera found in our study have been identified in other mosquito species before. Due to the currently scarce data situation, ongoing examinations on the very abundant bacterial genera or species are strongly required to determine their relevance for the biology and adaptiveness of mosquitoes including pathogen-host relationship.

Introduction

Over the last few years, the interest in microorganisms associated with other organisms has increased rapidly, which can be partly explained by the recognition of their impact for vertebrates as well as invertebrates has been recognized. It has now been recognized that host organisms are not dissociable from their microbial partners; together they form the holobiont (Guerreo et al. 2013). For instance, the digestion of xylophagous termites mainly relies on intestinal microorganisms (Hongoh et al. 2005; Köhler et al. 2012). Another benefit for the host may be the provision of colonization resistance in insects, with indigenous microbiota inhibiting the settlement of pathogens in the gastrointestinal tract (Berg 1996). Different studies on *Drosophila*, locusts and also bumblebees found links between different bacterial endosymbionts and the protection of the host organism against viral infections (Dillon et al. 2005; Hedges et al. 2008; Koch and Schmid-Hempel 2011). Other studies showed that endosymbionts can lead to

✉ Sina Zotzmann
s.zotzmann@web.de

¹ Institute for Ecology, Evolution and Diversity, Goethe-University (GU); Senckenberg Biodiversity and Climate Research Centre (BiK-F); Senckenberg Gesellschaft für Naturforschung (SGN), Max-von-Laue-Str. 13, 60438 Frankfurt/M., Germany

² Ecologie Microbienne, Université Claude Bernard Lyon, Bat. André Lwoff, 10 Rue Raphaël Dubois, 69100 Villeurbanne, France

³ Université de La Réunion, UMR PIMIT, INSERM 1187, CNRS 9192, IRD 249. Plateforme Technologique CYROI, Sainte-Clotilde, Réunion, France

sexual aberrations of the host insect, such as feminization, masculinization or cytoplasmic incompatibility (reviewed in Kageyama et al. 2012).

Mosquitoes (Culicidae) are one of the most important vectors for disease agents, the latter causing infectious diseases like malaria, dengue, West-Nile, Chikungunya or yellow fever as well as different forms of encephalitis (Becker et al. 2014; WHO 2016). They are also currently involved in the spreading of Zika virus in the Americas and the Pacific region. Every year, about 700 million people get infected with disease agents transmitted by mosquitoes and over 1 million of these cases are fatal (Caraballo and King 2014). Due to the spread of mosquitoes like *Ae. albopictus* into new geographical areas, some of the mosquito-borne diseases can now be found outside their endemic range (Bonilauri et al. 2008; La Ruche et al. 2010).

In the context of vector and pathogen control strategies, some new approaches using microbiota seem very promising. Several microorganisms show an influence on the development, the pathogen infection rates or the vector competence of mosquitoes (Werren 1997; Kambris et al. 2009; Minard et al. 2013). Natural bacterial endosymbionts of mosquitoes could also serve as a subject of paratransgenesis, i.e., the application of genetically modified microorganisms, which could be used to inhibit pathogens within the vector or pathogen transmission (Coutinho-Abreu et al. 2010).

Overall, studies on bacterial microbiota diversity in mosquitoes are mainly dealing with the medically most relevant vectors such as *Anopheles gambiae*, *Aedes aegypti* and *Aedes albopictus* (Wang et al. 2011; Zouache et al. 2012; Moro et al. 2013). Although a variety of established German mosquito species are also known as potential vectors for disease agents, there are currently no studies about the composition and structure of their associated bacterial communities. *Culex pipiens* (*Cx. pipiens*) is one of the most widespread mosquito species in Europe and North America and a potential vector for West-Nile virus (Fonseca et al. 2004; Diaz-Badillo et al. 2011; Hesson et al. 2014). In addition, pathogens like *Wuchereria bancrofti*, *Dirofilaria repens*, *Dirofilaria immitis* as well as Usutu and Sindbis virus have already been detected in this species (Ramzy et al. 1997; Cancrini et al. 2007; Jöst et al. 2010; Jöst et al. 2011; Yildirim et al. 2011). *Aedes japonicus* (*Ae. japonicus*, Asian bush mosquito) is an invasive mosquito species originally from Japan and Korea which can be found in Germany since 2008 (Tanaka et al. 1979; Schaffner et al. 2009) and is known as vector for West-Nile virus as well, but also for more tropical viruses such as Chikungunya or dengue virus (Sardelis and Turell 2001; Schaffner et al. 2011). *Aedes japonicus* is a mammalophilic mosquito, whereas *Cx. pipiens* is more known as an ornithophilic mosquito. As both species are often found in human settlements, they are of strong interest for human and veterinary medicine (Becker et al. 2011;

Werblow et al. 2014; Melaun et al. 2015). This study is the first study to investigate the bacterial microbiota diversity of the common mosquito species *Cx. pipiens* and the non-indigenous species *Ae. japonicus* in Germany.

Methods

Sampling and morphological identification

The analysed specimens of *Ae. japonicus* were collected during a survey in October 2014 and April and May 2015 at two locations (Backnang and Remshalden-Geradstetten, both urban environments) in Baden-Wuerttemberg, Germany. The sampling location is situated in the south of Germany, approximately 130 km north of the Swiss border. Larvae were caught with small hand nets from buckets in a private garden (Remshalden-Geradstetten) and flower vases on a cemetery (Backnang) and raised in the laboratory. After hatching, adults ($n = 14$) had no access to nutrition sources like sugar or blood, and were killed by freezing them at -20°C for at least 20 min. The adult *Culex* ($n = 23$) were sampled at the Federal Environmental Agency's premises in Berlin-Marienfelde (urban environment) between May and September 2014 with an EVS-Trap, and were stored at -20°C until further analysis. The sampling site is situated in the east of Germany, about 175 km south of the Baltic Sea.

The morphological identification of *Culex* (to genus level) and *Ae. japonicus* (to species level) was carried out with a stereomicroscope using the identification keys of Gutsevich et al. (1974), Tanaka et al. (1979) and Becker et al. (2010).

DNA extraction

Before DNA extraction, all mosquitoes were individually surface disinfected following the protocol of Zouache et al. (2011). The samples were then homogenized in 150 μl of sterile 0.8% sodium chloride solution with a tissue mill (MM400, Retsch GmbH, Germany) and two of 3 mm stainless steel beads (VWR, Germany) for 2 min at 25 Hz. Out of the homogenate, 75 μl was taken for DNA extraction using the peqGold Microspin Tissue DNA Mini kit (Peqlab Biotechnology GmbH, Erlangen, Germany), with additional 50 μl of lysozyme. DNA was eluted with 60 μl elution buffer.

Amplification of *cox1* genes

In order to verify the morphological species results, the cytochrome c oxidase subunit 1 (*cox1*) gene fragment was amplified for mosquito species identification. The reaction was performed with primers LCO 1490/HCO 2198 for all *Aedes* samples and 13 *Culex* individuals (Folmer et al. 1994). For the remaining *Culex* individuals (*Cx.* 6, *Cx.* 15 – *Cx.* 23, $n = 10$), a

different set of primers BC-kumar forward/reverse (Kumar et al. 2007) was used as they did not show any positive results with primers LCO 1490/HCO 2198. All PCR reaction mixtures contained 12.5 µl HotStart Mix Y (Peqlab Biotechnology GmbH, Erlangen, Germany), 1 µl of each primer (10 pmol µl⁻¹), as well as varying amounts of template DNA and ddH₂O, in a total volume of 25 µl. The amplification was performed in a thermocycler (Eppendorf, Germany) with the following cycle parameters for LCO 1490/HCO 2198: 1 cycle of 94 °C, 3 min; 40 cycles of 94 °C, 45 s; 37 °C, 45 s and 72 °C, 60 s followed by terminal extension of 72 °C, 10 min and a final ramping to 8 °C. The cycling parameters for primers used by Kumar et al. (2007) were 1 cycle of 94 °C, 2 min; 40 cycles of 94 °C, 60 s; 59 °C, 60 s and 72 °C, 60 s followed by terminal extension of 72 °C, 5 min and also a final ramping to 8 °C. Quality and yield of DNA was checked by Midori Green (Nippon Genetic EUROPE GmbH) staining and agarose gel-electrophoresis. Positive samples were purified using the peqGOLD Cycle-Pure Kit (Peqlab Biotechnology GmbH, Erlangen, Germany).

Amplification of 16S rDNA genes

The amplification of the bacterial 16S rDNA genes was carried out with eubacterial-specific primers pA/pH (Edwards et al. 1989; Moro et al. 2013) and the same reaction mix as described for the *cox1* amplification was used. PCR reactions were performed as described in Moro et al. (2013) with terminal extension of 72 °C, 12 min. The expected fragment length of about 1500 bp was again checked by Midori Green staining and agarose gel-electrophoresis. Positive samples were used for cloning reactions.

Cloning reaction and plasmid purification

The cloning reaction (one per mosquito) was carried out with the pGEM-T vector system (Promega Corporation, USA) using the standard reaction protocol for ligation and transformation as described in the manufacturers manual. In preparation for the cloning reaction, we produced chemically competent cells out of a stock of 50 µl One Shot® Top 10 chemically competent *E. coli* cells (Invitrogen Life Technologies, USA) following the protocol described in Mülhardt (2009). The cells were portioned into 200 µl aliquots, shock frosted with liquid nitrogen and stored at -80 °C until further use. Clones, which were positive by blue/white screening, were checked for the proper insert length by amplification of the manufacturer recommended primer pair M13 forward/reverse. We used not more than 28 positive colonies from each plate to check for insert length. Each colony was picked with a sterile toothpick, added to a master plate and 20 µl ddH₂O. The solution of bacterial material and water was used for a cell lysis which was performed at 95 °C for 2 min followed by 2 min

incubation on ice (step performed twice) before centrifuged at 4000 rpm for 10 min. The supernatant was used as template DNA for the amplification with M13 forward/reverse in the same reaction mix as described for *cox1* and 16S rDNA. The amplification steps were 1 cycle of 94 °C, 12 min; 35 cycles of 94 °C, 60 s; 50 °C, 60 s and 72 °C, 60 s followed by terminal extension of 72 °C, 10 min and a final ramping to 8 °C. The insert length was checked by Midori green staining and agarose gel-electrophoresis. Clones with inserts of the expected size (about 1500 bp) were purified with the peqGOLD Plasmid Miniprep Kit according to the manufactures protocol without the third and optional washing step.

Sequencing reaction

For subsequent Sanger-sequencing reactions with product specific forward primers, PCR products were purified using the peqGOLD Cycle-Pure Kit (Peqlab Biotechnology GmbH, Erlangen, Germany). Each obtained sequence was edited using BioEdit (Hall 1999) and compared with sequences deposited in GenBank using the BLAST algorithm (Altschul et al. 1997).

Results

Molecular identification of mosquito species

The identification of the female adult *Culex* mosquitoes to species level resulted in 23 members of the *Cx. pipiens* complex. As neither the morphological differentiation nor DNA barcoding is utterly reliable, no distinction was made between *Cx. pipiens* biotype *pipiens* and *Cx. pipiens* biotype *molestus*. All 14 laboratory-reared female individuals of *Ae. japonicus* were determined to species level. The sequences of the molecular species identification are given under accession numbers KX260917-KX260953 in the GenBank database. All of the 23 *Culex* and 14 *Ae. japonicus* mosquitoes were used for subsequent bacterial analyses.

Bacterial diversity in *Culex pipiens* and *Aedes japonicus*

After the cloning reactions not more than 28 colonies per mosquito specimen (positive by blue/white screening, *Cx. pipiens* = 550, *Ae. japonicus* = 384) were picked and checked for the correct insert. Among them, we only used sequences with more than 800 bp and a query cover and identity in GenBank of 98–100%, resulting in a total of 282 analysed clones. Sequence-based identification classified bacteria in 2 phyla, 4 classes, 14 families and 21 genera for *Cx. pipiens* (Table 1). Although fewer samples of *Ae. japonicus* were analysed, a higher rate of positive clones was identified including one additional phylum, the Actinobacteria. Overall,

Table 1 Taxonomic assignments of bacterial clones from *Culex pipiens*. In total, 123 clones were sequenced with eubacterial-specific primer pA for bacterial 16S *rrs* genes. Only results with a sequence length >800 bp and a query cover and identity in NCBI GenBank between 98 and 100% are shown

Phylum	Class	Family	Nearest genus acc. to BLASTn	Nearest species acc. to BLASTn	Ident [%]	Sequences
Firmicutes	Bacilli	Carnobacteriaceae	<i>Carnobacterium</i>	<i>Carnobacterium maltaromaticum</i>	99	1
		Enterococcaceae	<i>Enterococcus</i>	<i>Enterococcus</i> sp.	99	1
		Staphylococcaceae	<i>Staphylococcus</i>	<i>Staphylococcus warneri</i>	99	1
		Streptococcaceae	<i>Streptococcus</i>	<i>Streptococcus thermophilus</i>	99	5
Proteobacteria	α -Proteobacteria	Bradyrhizobiaceae	<i>Afipia</i>	<i>Afipia</i> sp.	100	1
			<i>Bosea</i>	<i>Bosea vestrisii</i>	99	1
		Methylobacteriaceae	<i>Methylobacterium</i>	<i>Methylobacterium</i> spp.	99	2
		Phyllobacteriaceae	<i>Mesorhizobium</i>	<i>Mesorhizobium</i> spp.	99	3
		Rhizobiaceae	<i>Agrobacterium</i>	<i>Agrobacterium tumefaciens</i>	100	1
		Sphingomonadaceae	<i>Sphingomonas</i>	<i>Sphingomonas echinoides</i>	99	39
			<i>Sphingomoas leidyi</i>	99	1	
			<i>Sphingomonas</i> spp.	99	1	
	β -Proteobacteria	Comamonadaceae	<i>Acidovorax</i>	<i>Acidovorax</i> spp.	99	8
			<i>Curvibacter</i>	<i>Curvibacter</i> spp.	99	12
		Ralstoniaceae	<i>Ralstonia</i>	<i>Ralstonia</i> spp.	99	7
	γ -Proteobacteria	Enterobacteriaceae	<i>Escherichia</i>	<i>Escherichia coli</i>	99	1
			<i>Hafnia</i>	<i>Hafnia paralvei</i>	98	1
			<i>Kluyvera</i>	<i>Kluyvera</i> sp.	99	1
			<i>Pantoea</i>	<i>Pantoea coffeiphila</i>	98	6
<i>Rahnella</i>			<i>Rahnella aquatilis</i>	99	4	
<i>Tatumella</i>			<i>Tatumella</i> spp.	99	3	
<i>Tatumella ptyseos</i>			99	1		
<i>Acinetobacter</i>			<i>Acinetobacter</i> sp.	99	1	
Pseudomonaceae		<i>Pseudomonas</i>	<i>Pseudomonas</i> spp.	99	2	
		<i>Pseudomonas psychrotolerans</i>	99	3		
Unknown	Unknown	Unknown	Unknown	Uncultured bacterium clone	99	2
$\Sigma 2$	4	14	21	26		123

the 159 bacterial clones of *Ae. japonicus* could be subdivided into 3 phyla, 5 classes, 9 families and 11 genera (Table 2). The most abundant classes found in *Cx. pipiens* were the α -Proteobacteria (no. of sequences, $n = 48$) with the genus *Sphingomonas* ($n = 40$) and the species *Sphingomonas echinoides* ($n = 39$) as well as the β -Proteobacteria ($n = 42$) with the most abundant genera *Ralstonia* ($n = 22$) and *Curvibacter* ($n = 12$). Within the γ -Proteobacteria only 23 bacterial isolates were found, however, this class had the highest number of genera ($n = 8$). The lowest number of bacterial isolates ($n = 8$) was found within the class Bacilli (Table 1). Similar to *Cx. pipiens*, the phylum Proteobacteria was also the most abundant phylum ($n = 145$) in *Ae. japonicus* with the genus *Rahnella* ($n = 104$) representing the highest number of isolates (Table 2). The genera *Afipia*, *Mesorhizobium*, *Sphingomonas* as well as *Rahnella* were shared between *Cx. pipiens* and *Ae. japonicus*, whereas all other genera ($n = 28$) were present in only one of the mosquito

species. The bacterial sequences obtained were deposited in the GenBank database under accession numbers KX260635-KX260916.

Discussion

In this study, the bacterial microbiota of the common house mosquito *Cx. pipiens* and the non-indigenous species *Ae. japonicus* in Germany were investigated using a culture-independent method.

Most of the sequences (91.9%) obtained in *Cx. pipiens* belonged to the gram-negative phylum Proteobacteria (class: α -, β - and γ -Proteobacteria). Most of the remaining sequences (6.5%) belonged to the gram-positive phylum Firmicutes (class: Bacilli). High numbers of sequences were affiliated with *Sphingomonas echinoides*, a bacterium which was first isolated from plants and exists in seawater as well

Table 2 Taxonomic assignments of bacterial clones from *Aedes japonicus*. In total, 159 clones were sequenced with eubacterial-specific primer pA for bacterial 16S *rrs* genes. Only results with a sequence length >800 bp and a query cover and identity in NCBI GenBank between 98 and 100% are shown

Phylum	Class	Family	Nearest genus acc. to BLASTn	Nearest species acc. to BLASTn	Ident [%]	Sequences			
Actinobacteria	Actinobacteria	Microbacteriaceae	<i>Microbacterium</i>	<i>Microbacterium oxydans</i>	99	9			
				<i>Microbacterium maritypicum</i>	99	3			
		Propionibacteriaceae	<i>Propionibacterium</i>	<i>Propionibacterium acnes</i>	99	1			
Firmicutes	Bacilli	Carnobacteriaceae	<i>Granulicatella</i>	<i>Granulicatella adiacens</i>	99	1			
Proteobacteria	α -Proteobacteria	Bradyrhizobiaceae	<i>Afipia</i>	<i>Afipia</i> sp.	99	1			
				Caulobacteraceae	<i>Caulobacter</i>	<i>Caulobacter</i> spp.	98	2	
		Phyllobacteriaceae	<i>Mesorhizobium</i>	<i>Mesorhizobium</i> spp.	99	17			
				<i>Mesorhizobium plurifarum</i>	99	1			
				<i>Sphingomonas</i>	<i>Sphingomonas echinoides</i>	99	2		
					<i>Sphingomonas melonis</i>	99	1		
		β -Proteobacteria		<i>Aquincola</i>	<i>Aquincola</i> sp.	98	1		
		γ -Proteobacteria	Aeromonadaceae	<i>Aeromonas</i>	<i>Aeromonas salmonicida</i>	100	1		
					Enterobacteriaceae	<i>Ewingella</i>	<i>Ewingella americana</i>	99	15
						<i>Rahnella</i>	<i>Rahnella</i> spp.	99	57
				<i>Rahnella aquatilis</i>	99	47			
Σ 3	5	9	14	15		159			

(Heumann 1960; De Vos et al. 1989; Kim et al. 2006). *Sphingomonas* spp. were already detected in other mosquito species (i.a. Moro et al. 2013; Dhayal et al. 2014; Ngo et al. 2015). Minard et al. (2015) found that the bacterial family Sphingomonadaceae is very prevalent in mosquitoes. These bacteria have the ability to produce hydrolases involved in the degradation of oligosaccharides, so that there might be a link between the bacteria and the provision of plant sugar to the mosquito host (Minard et al. 2015). Demaio et al. (1996) identified eight bacterial species in a culture-dependent study of microbiota of *Cx. pipiens* in the USA. Two of those were also found in our study belonging to the genera *Acinetobacter* and *Pseudomonas* (both γ -Proteobacteria). The differences regarding the total number of genera found in our study ($n = 21$, culture-independent) and the study by Demaio et al. (1996) ($n = 8$, culture-dependent) might be explained by the choice of method which influences the number of species detected, e.g., only a few bacteria grow on culture media (Hugenholtz et al. 1998; Charan et al. 2013).

The most abundant bacterial phylum in *Ae. japonicus* was also the gram-negative Proteobacteria (class: α -, β - and γ -Proteobacteria) with 91.2% of sequences. The remaining sequences belonged to the gram-positive phyla Actinobacteria and Firmicutes (class: Actinobacteria and Bacilli). The most abundant bacterial sequences were affiliated with *Rahnella aquatilis*, a species that is known to occur in different environments, including fresh water and soil (Guo et al. 2012; Martinez et al. 2012) and has been found in mosquitoes, e.g., within larvae of *Anopheles stephensi* (Dinparast Djadid et al. 2011).

Kim et al. (2015) examined the microbiota of larval as well as adult *Ae. japonicus* from two different sampling sites in the USA. In their culture-independent approach, they found that the relative abundance of bacterial taxa varies between habitats as well as between mosquitoes' developmental stages. In their culture-dependent attempt, they detected 10 bacterial genera in larval *Ae. japonicus* and 2 genera in newly emerged adults. Two of the bacterial genera found by Kim et al. (2015) in the USA were identified in our study as well. Those were *Aeromonas* and *Microbacterium* which were found by Kim et al. (2015) in adults (*Aeromonas*) and larvae (*Microbacterium*). Interestingly, another invasive mosquito species (*Aedes albopictus*, Asian tiger mosquito) showed a reduced bacterial microbiota in recently established populations (France) compared to autochthonous populations (Vietnam), possibly due to an altered/reduced availability of plant nutrition sources within the new habitats (Minard et al. 2015). Such comparative analyses on the invasive rock pool mosquito *Ae. japonicus* are still lacking.

Great differences regarding the bacterial diversity between individuals from laboratory breeding and wild-caught specimens can occur as shown by Rani et al. (2009) for *Anopheles stephensi*. In their study, only 7 bacterial taxa were present in laboratory-reared female mosquitoes in contrast to 36 taxa in wild-caught specimens. This might explain differences in bacterial taxa between the examined mosquito species here (21 genera and 11 genera found in *Cx. pipiens* and *Ae. japonicus* respectively). Individuals of *Ae. japonicus* were laboratory-reared to adults from field-caught larvae and were examined shortly after hatching, whereas specimens of *Cx. pipiens* were

wild-caught as adults. Furthermore, it has been shown that metamorphosis in mosquitoes leads to reduced bacterial content and diversity in newly emerged adults (Demaio et al. 1996; Moll et al. 2001; Wang et al. 2011).

The 7 bacteria genera *Afipia*, *Aquicola*, *Bosea*, *Caulobacter*, *Curvibacter*, *Granulicatella* and *Mesorhizobium* were recorded for the first time in mosquitoes, while the remaining 21 genera in our study have been found in other studies before, mainly in species that are medically highly relevant, like *Aedes aegypti*, *Ae. albopictus*, *Anopheles gambiae* and *An. stephensi* (Minard et al. 2013; Dhayal et al. 2014; Ngo et al. 2015). Overall, more than 90% of the analysed isolates were gram-negative bacteria of the phylum Proteobacteria in both mosquito species, *Cx. pipiens* and *Ae. japonicus*. The dominance of this phylum was observed in different mosquito species before (Pidiyar et al. 2004; Osei-Poku et al. 2012; Moro et al. 2013).

The detection of different bacteria species of the same genus could imply a high association of these bacteria genera with mosquitoes, possibly providing a better adaptability to the environment (Moro et al. 2013). However, the research of the natural microbiota of mosquitoes is only at an early stage. Further studies are necessary to better understand and differentiate the importance of different bacteria for mosquitoes, which could also foster further research of paratransgenetic approaches to assist the control of mosquito-borne diseases. There might be general interspecific variations in associated bacterial diversity of mosquito species based on different ecological and biological habits; however, those cannot be sufficiently evaluated due to the currently scarce data situation.

Conclusion

In this study, the microbiota of two mosquito species, *Cx. pipiens* and *Ae. japonicus*, in Germany were investigated. While 21 bacterial taxa have been also reported from other mosquitoes, 7 identified genera have not been recognized before and are therefore first time records. Further analyses are strongly needed to decipher the significance of associated symbiotic bacteria regarding mosquito biology, adaptiveness and the impact on vector-pathogen-interactions. This knowledge could provide new approaches for mosquito control and the decrease of mosquito-borne infectious diseases.

Acknowledgements This present research was funded by the ERA-Net BiodivERsA, with the national funders DFG KL 2087/6-1, FWF I-1437 and ANR-13-EBID-0007-01 as part of the 2012-13 BiodivERsA call for research proposals and by the German Federal Ministry of Food and Agriculture (BMEL) through the Federal Office for Agriculture and Food (BLE), grant number 2819105115 as well as by the Uniscientia Foundation (P 121-2017). We thank Birgit Nagel and Gabriele Elter for their technical assistance in the molecular laboratory. We also thank Dr.

Rüdiger Berghahn and Ronny Schmiediche from the German Federal Environmental Agency (UBA) in Berlin for mosquito sampling.

References

- Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res* 25: 3389–3402
- Becker N, Petrić D, Zgomba M, Boase C, Dahl C, Madon M, Kaiser A (2010) Mosquitoes and their control, 2nd edn. Springer-Verlag, Berlin Heidelberg
- Becker N, Huber K, Pluskota B, Kaiser A (2011) *Ochlerotatus japonicus japonicus*—a newly established neozoan in Germany and a revised list of the German mosquito fauna. *Eur Mosq Bull* 29:88–102
- Becker N, Krüger A, Kuhn C, Plenge-Bönig A, Thomas SM, Schmidt-Chanasit J, Tannich E (2014) Stechmücken als Überträger exotischer Krankheitserreger in Deutschland. *Bundesgesundheitsbl* 57:531–540. doi:10.1007/s00103-013-1918-8
- Berg RD (1996) The indigenous gastrointestinal microflora. *Trends Microbiol* 11:430–435
- Bonilauri P, Bellini R, Calzolari M, Angelini R, Venturi L, Fallacara F, Cordioli P, Angelini P, Venturelli C, Merialdi G, Dottori M (2008) Chikungunya virus in *Aedes albopictus*, Italy. *Emerg Infect Dis* 14: 852–854. doi:10.3201/eid1405.071144
- Cancrini G, Scaramozzino P, Gabrielli S, Di Paolo M, Toma L, Romi R (2007) *Aedes albopictus* and *Culex pipiens* implicated as natural vectors of *Dirofilaria repens* in Central Italy. *J Med Entomol* 44: 1064–1066
- Caraballo H, King K (2014) Emergency department management of mosquito-borne illness: malaria, dengue, and West Nile virus. *Emerg Med Pract* 16:1–23
- Charan SS, Pawar KD, Severson DW, Patole MS, Shouche YS (2013) Comparative analysis of midgut bacterial communities of *Aedes aegypti* mosquito strains varying in vector competence to dengue virus. *Parasitol Res* 112:2627–2637. doi:10.1007/s00436-013-3428-x
- Coutinho-Abreu IV, Zhu KY, Ramalho-Ortigao M (2010) Transgenesis and paratransgenesis to control insect-borne diseases: current status and future challenges. *Parasitol Int* 59:1–8. doi:10.1016/j.parint.2009.10.002
- Demaio J, Pumpuni CB, Kent M, Beier JC (1996) The midgut bacterial flora of wild *Aedes triseriatus*, *Culex pipiens*, and *Psorophora columbiae* mosquitoes. *AmJTrop Med Hyg* 54:219–223
- De Vos P, Van Landschoot A, Segers P, Tytgat R, Gillis M, Bauwens M, Rossau R, Goor M, Pot B, Kersters K, Lizzaraga P, De Ley J (1989) Genotypic relationships and taxonomic localization of unclassified *Pseudomonas* and *Pseudomonas*-like strains by deoxyribonucleic acid: ribosomal ribonucleic acid hybridizations. *Int J Syst Bacteriol* 39:35–49
- Dhayal D, Parasher H, Sharma A, Kumar P, Adak T, Jaiwal R (2014) Diversity of culturable midgut bacteria of Indian malarial vector *Anopheles stephensi*. *IJARM* 2:305–311
- Diaz-Badillo A, Bolling BG, Perez-Ramirez G, Moore CG, Martinez-Munoz JP, Padilla-Viveros AA, Camacho-Nuez M, Diaz-Perez A, Beaty BJ, Lourdes Munoz M (2011) The distribution of potential West Nile virus vectors, *Culex pipiens pipiens* and *Culex pipiens quinquefasciatus* (Diptera: Culicidae), in Mexico City. *Parasite Vectors* 4:70. doi:10.1186/1756-3305-4-70
- Dillon RJ, Vennard CT, Buckling A, Charnley AK (2005) Diversity of locust gut bacteria protects against pathogen invasion. *Ecol Lett* 8: 1291–1298. doi:10.1111/j.1461-0248.2005.00828.x

- Dinparast Djadjid N, Jazayeri H, Raz A, Favia G, Ricci I, Zakeri S (2011) Identification of the midgut microbiota of *An. stephensi* and *An. maculipennis* for their application as a paratransgenetic tool against malaria. *PLoS One* 6:e28484. doi:10.1371/journal.pone.0028484
- Edwards U, Rogall T, Blöcker H, Emde M, Böttger EC (1989) Isolation and direct complete nucleotide determination of entire genes. Characterization of a gene coding for 16S ribosomal RNA. *Nucleic Acids Res* 17:7843–7853
- Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R (1994) DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Mol Mar Biol Biotechnol* 3(5): 294–299
- Fonseca DM, Keyghobadi N, Malcolm CA, Mehmet C, Schaffner F, Mogi M, Fleischer RC, Wilkerson RC (2004) Emerging vectors in the *Culex pipiens* complex. *Science* 303:1535–1538. doi:10.1126/science.1094247
- Guerreo R, Margulis L, Berlanga M (2013) Symbiogenesis: the holobiont as a unit of evolution. *Int Microbiol* 16:133–143. doi:10.2436/20.1501.01.188
- Guo Y, Jiao Z, Li L, Wu D, Crowley DE, Wang Y, Wu W (2012) Draft genome sequence of *Rahnella aquatilis* strain HX2, a plant growth-promoting Rhizobacterium isolated from vineyard soil in Beijing, China. *J Bacteriol* 194:6646–6647. doi:10.1128/JB.01769-12
- Gutsevich AV, Monchadskii AS, Shtakel'berg AA (1974) Fauna of the U.S.S.R. NS: Diptera: mosquitoes: family Culicidae. 2nd ed. Leningrad. Akademiya Nauk SSSR-Zoologicheskii institute, Leningrad
- Hall TA (1999) BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symp Ser* 41:95–98
- Hedges LM, Brownlie JC, O'Neill SL, Johnson KN (2008) *Wolbachia* and virus protection in insects. *Science* 322:702. doi:10.1126/science.1162418
- Hesson JC, Rettich F, Merdić E, Vignjević G, Östman Ö, Schäfer M, Schaffner F, Foussadier R, Besnard G, Medlock J, Scholte EJ, Lundström JO (2014) The arbovirus vector *Culex torrentium* is more prevalent than *Culex pipiens* in northern and central Europe. *Med Vet Entomol* 28:179–186. doi:10.1111/mve.12024
- Heumann W (1960) Versuche zur Rekombination sternbildender Bakterien. *Naturwissenschaften* 47:330–331
- Hongoh Y, Deevong P, Inoue T, Moriya S, Trakulnaleamsai S, Ohkuma M, Vongkhaluang C, Noparatnaraporn N, Kudo T (2005) Intra- and interspecific comparison of bacterial diversity and community structure support coevolution of gut microbiota and termite host. *Appl Environ Microb* 71:6590–6599. doi:10.1128/AEM.71.11.6590-6599.2005
- Hughenholz P, Goebel BM, Pace NR (1998) Impact of culture-independent studies on the emerging phylogenetic view of bacterial diversity. *J Bacteriol* 180:4765–4774
- Jöst H, Bialonski A, Storch V, Günther S, Becker N, Schmidt-Chanasit J (2010) Isolation and phylogenetic analysis of Sindbis viruses from mosquitoes in Germany. *J Clin Microbiol* 48:1900–1903. doi:10.1128/JCM.00037-10
- Jöst H, Bialonski A, Maus D, Sambri V, Eiden M, Groschup MH, Günther S, Becker N, Schmidt-Chanasit J (2011) Isolation of Usutu virus in Germany. *AmJTrop Med Hyg* 85:551–553. doi:10.4269/ajtmh.2011.11-0248
- Kageyama D, Narita S, Watanabe M (2012) Insect sex determination manipulated by their endosymbionts: incidences, mechanisms and implications. *Insects* 3:161–199. doi:10.3390/insects3010161
- Kambris Z, Cook PE, Phuc HK, Sinkins SP (2009) Immune activation by life-shortening *Wolbachia* and reduced filarial competence in mosquitoes. *Science* 326:134–136. doi:10.1126/science.1177531
- Kim C-H, Lampman R, Muturi EJ (2015) Bacterial communities and midgut microbiota associated with mosquito population from waste tires in east-central Illinois. *J Med Entomol* 52:63–75. doi:10.1093/jme/tju011
- Kim HS, Lee OK, Lee SJ, Hwang S, Kim SJ, Yang SH, Park S, Lee EY (2006) Enantioselective epoxide hydrolase activity of a newly isolated microorganism, *Sphingomonas echinoides* EH-983, from seawater. *J Mol Catal B-Enzym* 41:130–135. doi:10.1016/j.molcatb.2006.05.009
- Koch H, Schmid-Hempel P (2011) Socially transmitted gut microbiota protect bumble bees against an intestinal parasite. *P Natl Acad Sci* 108:19288–19292. doi:10.1073/pnas.111047410
- Köhler T, Dietrich C, Scheffrahn RH, Brune A (2012) High-resolution analysis of gut environment and bacterial microbiota reveals functional compartmentation of the gut in wood-feeding higher termites. *Appl Environ Microb* 78:4691–4701. doi:10.1128/AEM.00683-12
- Kumar NP, Rajavel AR, Natarajan R, Jambulingam P (2007) DNA barcodes can distinguish species of Indian mosquitoes (Diptera: Culicidae). *J Med Entomol* 44:1–7. doi:10.1093/jmedent/41.5.01
- La Ruche G, Souares Y, Armengaud A, Peloux-Petiot F, Delaunay P, Despres P, Lenglet A, Jourdain F, Leparc-Goffart I, Charlet F, Ollier L, Mantey K, Mollet T, Fournier JP, Torrents R, Leitmeyer K, Hilairet P, Zeller H, Van Bortel W, Dejour-Salamanca D, Grandadam M, Gastellu-Etchegorry M (2010) First two autochthonous dengue virus infections in metropolitan France, September 2010. *Euro Surveill* 15:19676
- Martinez RJ, Bruce D, Detter C, Goodwin LA, Han J, Han CS, Held B, Land ML, Mikhailova N, Nolan M, Pennacchio L, Pitluck S, Tapia R, Woyke T, Sobecky PA (2012) Complete genome sequence of *Rahnella aquatilis* CIP 78.65. *J Bacteriol* 194:3020–3021. doi:10.1128/JB.00380-12
- Melaun C, Werblow A, Cunze S, Zotzmann S, Koch LK, Mehlhorn H, Dörge DD, Huber K, Tackenberg O, Klimpel S (2015) Modeling of the putative distribution of the arbovirus vector *Ochlerotatus japonicus japonicus* (Diptera: Culicidae) in Germany. *Parsitol Res* 114:1051–1061. doi:10.1007/s00436-014-4274-1
- Minard G, Mavingui P, Moro CV (2013) Diversity and function of bacterial microbiota in the mosquito holobiont. *Parasite Vector* 6:146. doi:10.1186/1756-3305-6-146
- Minard G, Tran FH, van Tran Van CG, Goubert C, Bellet C, Lambert G, Kim KL, Thuy TH, Mavingui P, Moro CV (2015) French invasive Asian tiger mosquito populations harbor reduced bacterial microbiota and genetic diversity compared to Vietnamese autochthonous relatives. *Front Microbiol* 6:970. doi:10.3389/fmicb.2015.00970
- Moll RM, Romoser WS, Modrakowski MC, Moncayo AC, Lerdthusnee K (2001) Meconial peritrophic membranes and the fate of midgut bacteria during mosquito (Diptera: Culicidae) metamorphosis. *J Med Entomol* 38:29–32. doi:10.1603/0022-2585-38.1.29
- Moro CV, Tran FH, Raharimalala FN, Ravelonandro P, Mavingui P (2013) Diversity of culturable bacteria including *Pantoea* in wild mosquito *Aedes albopictus*. *BMC Microbiol* 13:70. doi:10.1186/1471-2180-13-70
- Mühlhardt C (2009) *Der Experimentator Molekularbiologie/Genomics*. 6th edn. Spektrum Akademischer Verlag Heidelberg
- Ngo CT, Aujoulat F, Veas F, Jumas-Bilak E, Manguin S (2015) Bacterial diversity associated with wild caught *Anopheles* mosquito from Dak Nong province, Vietnam using culture and DNA fingerprint. *PLoS One* 10:e0118634. doi:10.1371/journal.pone.0118634
- Osei-Poku J, Mbogo CM, Palmer WJ, Jiggins FM (2012) Deep sequencing reveals extensive variation in the gut microbiota of wild mosquitoes from Kenya. *Mol Ecol* 21:5138–5150. doi:10.1111/j.1365-294X.2012.05759.x
- Pidiyar VJ, Jangid K, Patole MS, Shouche Y (2004) Studies on cultured and uncultured microbiota of wild *Culex quinquefasciatus* mosquito midgut based on 16S ribosomal RNA gene analysis. *AmJTrop Med Hyg* 70:597–603
- Ramzy RMR, Farid HA, Kamal IH, Ibrahim GH, Morsy ZS, Faris R, Weil GJ, Williams SA, Gad AM (1997) A polymerase chain

- reaction-based assay for detection of *Wuchereria bancrofti* in human blood and *Culex pipiens*. *Trans R Soc Trop Med Hyg* 91:156–160
- Rani A, Sharma A, Rajagopal R, Adak T, Bhatnagar RK (2009) Bacterial diversity analysis of larvae and adult midgut microflora using culture-dependent and culture-independent methods in laboratory-reared and field-collected *Anopheles stephensi*—an Asian malaria vector. *BMC Microbiol* 9:96. doi:10.1186/1471-2180-9-96
- Sardelis MR, Turell MJ (2001) *Ochlerotatus j. japonicus* in Frederick County, Maryland: discovery, distribution, and vector competence for West Nile virus. *J Am Mosq Control Assoc* 17:137–141
- Schaffner F, Kaufmann C, Hegglin D, Mathis A (2009) The invasive mosquito *Aedes japonicus* in Central Europe. *Med Vet Entomol* 23:448–451
- Schaffner F, Vazeille M, Kaufmann C, Failloux A-B, Mathis A (2011) Vector competence of *Aedes japonicus* for chikungunya and dengue viruses. *Eur Mosq Bull* 29:141–142
- Tanaka K, Mizusawa K, Saugstad ES (1979) A revision of the adult and larval mosquitoes of Japan (including the Ryukyu Archipelago and the Ogasawara Islands) and Korea (Diptera: Culicidae). *Contrib Amer Ent Inst* 16:419–422
- Wang Y, Gilbreath TM, Kukutla P, Yan G, Xu J (2011) Dynamic gut microbiom across life history of the malaria mosquito *Anopheles gambiae* in Kenya. *PLoS One* 6:e24767. doi:10.1371/journal.pone.0024767
- Werblow A, Klimpel S, Boliu S, Dorresteijn AWC, Sauer J, Melaun C (2014) Population structure and distribution patterns of the sibling mosquito species *Culex pipiens* and *Culex torrentium* (Diptera: Culicidae) reveal different evolutionary paths. *PLoS One* 9:e102158. doi:10.1371/journal.pone.0102158
- Werren JH (1997) Biology of *Wolbachia*. *Annu Rev Entomol* 42:587–609
- World Health Organization (2016) Vector-borne diseases. Available at: http://www.who.int/mediacentre/factsheets/fs_387/en/ (accessed 04.10.2017)
- Yildirim A, Inci A, Duzlu O, Biskin Z, Ica A, Sahin I (2011) *Aedes vexans* and *Culex pipiens* as the potential vectors of *Dirofilaria immitis* in Central Turkey. *Vet Parasitol* 178:1543–1147
- Zouache K, Raharimalala FN, Raquin V, Tran-Van V, Raveloson LHR, Ravelonandro P, Mavingui P (2011) Bacterial diversity of field-caught mosquitoes, *Aedes albopictus* and *Aedes aegypti*, from different geographic regions of Madagascar. *FEMS Microbiol Ecol* 75:377–389. doi:10.1111/j.1574-6941.2010.01012.x
- Zouache K, Michelland RJ, Failloux A-B, Grundmann GL, Mavingui P (2012) Chikungunya virus impacts the diversity of symbiotic bacteria in mosquito vector. *Mol Ecol* 21:2297–2309. doi:10.1111/j.1365-294X.2012.05526.x