



States in some ordered structures and axioms of choice

Martine Barret

► **To cite this version:**

| Martine Barret. States in some ordered structures and axioms of choice. 2018. hal-01785790

HAL Id: hal-01785790

<http://hal.univ-reunion.fr/hal-01785790>

Submitted on 4 May 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

States in some ordered structures and axioms of choice

Martine BARRET

University of la Réunion

Positivity *IX*
Edmonton 2017

July, 17-21

Axiom of Choice

AC : Given an infinite family $(A_i)_{i \in I}$ of non-empty sets, the product $\prod_{i \in I} A_i$ is non-empty.

We work in **ZF**, set theory **without the Axiom of Choice**.

We consider the **Hahn-Banach axiom (HB)**, a weak form of the Axiom of Choice. Remark : in **ZF**, **HB** is not provable and **HB** does not imply **AC** (see Howard and Rubin's book, [3]).

Theorem 1 : in **ZF**, **HB** implies the following statement **S_g**

S_g : For every abelian ordered group G with a positive order unit e and every subgroup H of G such that $e \in H$, every e -state on H can be extended into a e -state on G .

Remark : the converse statement **S_g** \Rightarrow **HB** also holds in **ZF**.

Hahn-Banach Axiom **HB** : a weak form of the **AC**

HB : Given a real vector space E , a sublinear mapping $p : E \rightarrow \mathbb{R}$ (i.e. a subadditive mapping such that for all $t \in \mathbb{R}^+$ and for all $x \in E, p(tx) = tp(x)$), a vector subspace S of E and a linear mapping $f : S \rightarrow \mathbb{R}$ such that $f \leq p|_S$, there exists a linear mapping $g : E \rightarrow \mathbb{R}$ extending f such that $g \leq p$.

Corollary 1 : in **ZF**, **HB** implies the classical following statement

Given a real normed vector space $(E, \| \cdot \|)$ and $a \in E \setminus \{0\}$, there exists a linear form $\varphi : E \rightarrow \mathbb{R}$ continuous of norm 1 such that $\varphi(a) = \|a\|$.

Proof : apply **HB** to the sublinear mapping $p := \| \cdot \|$, the vector

subspace $S := \text{Vect}(a)$ and the linear form

$$f : S \rightarrow \mathbb{R}$$
$$\lambda a \mapsto \lambda \|a\| .$$

1 On partially ordered groups :

Let G be an abelian ordered group. A non-zero element e of G is an **order unit** of G if : $\forall x \in G \exists k \in \mathbb{Z} -ke \leq x \leq ke$.

2 On partially ordered vector spaces :

Let E be an ordered vector space over \mathbb{R} .

- An element $e \in E \setminus \{0\}$ is an *order unit* of E if it is an order unit of the ordered group $(E, +)$. For an order unit e of E : $e \in E^+$ or $-e \in E^+$.
- Given a positive order unit $e \in E^+$ we associate a *semi-norm* $\| \cdot \|_e$ defined by :

$$\forall x \in E \quad \|x\|_e := \inf\{t \in \mathbb{R}^+, -te \leq x \leq te\}$$

- The semi-norms associated to two positive order units are equivalent and then, they define the same topology on E .

1 On ordered groups :

Let G be an abelian ordered group. A group morphism $f : G \rightarrow \mathbb{R}$ is *positive* if it is *increasing* i.e. $\forall x, y \in G, (x \leq y \Rightarrow f(x) \leq f(y))$.

2 On ordered vector spaces :

Lemma 1 (Characterisation)

Let $f : E \rightarrow \mathbb{R}$ be a linear form on an ordered vector space E with an order unit $e \in E^+$. Then :

f is positive (i.e. increasing) if and only if f is continuous of norm $f(e)$.

Proof :

- Assume that f is positive. Let $x \in E$: there exists $s \in \mathbb{R}_+^*$ such that $-se \leq x \leq se$, so $|f(x)| \leq s|f(e)|$ and then f is continuous and of norm $f(e)$.
- Now assume that f is continuous of norm $f(e)$. Let $x \in E^+$, show that $f(x) \geq 0$:
 - If $x \leq e$ then $0 \leq e - x \leq e$ so $f(e - x) \leq \|f\| \cdot \|e - x\|_e \leq f(e)$ and finally $f(x) \geq 0$.
 - If $x \not\leq e$, there exists $s \in \mathbb{R}_+^*$ such that $-se \leq x \leq se$ then, apply the previous case to $\frac{1}{s}x$.

Concurrent relations (Luxemburg, [4])

Let X and Y be two sets. Given a binary relation \mathcal{R} on $X \times Y$, for every $x \in X$, we define $\mathcal{R}(x) := \{y \in Y \mid x\mathcal{R}y\}$.

- The relation \mathcal{R} is *concurrent* if for every finite subset $F := \{x_1, \dots, x_n\}$ of X , the intersection $\mathcal{R}(x_1) \cap \dots \cap \mathcal{R}(x_n)$ is non-empty.
- If \mathcal{R} is a concurrent relation on $X \times Y$, we can define the *filter* \mathcal{F} on Y generated by the sets $\mathcal{R}(x)$, $x \in X$:

$$\mathcal{F} := \{A \subseteq Y \mid \exists x_1, \dots, x_n \in X \ \mathcal{R}(x_1) \cap \dots \cap \mathcal{R}(x_n) \subseteq A\}$$

Definition

Let E be a real vector space. Consider a set T and \mathcal{F} a filter over T . Denote by Z the following vector subspace of the vector space E^T :

$$Z := \{(x_t)_{t \in T} \in E^T \mid \{t \in T \mid x_t = 0\} \in \mathcal{F}\}$$

The **reduced power** E^T/\mathcal{F} of E by the filter \mathcal{F} is the quotient vector space E^T/Z . We denote by \bar{z} the class of an element $z \in E^T$ and we

consider the canonical embedding :

$$\begin{array}{ccc} \text{can} : & E & \rightarrow & E^T/\mathcal{F} \\ & x & \mapsto & \overline{(x)_{t \in T}} \end{array}$$

Remarks : if E is an ordered vector space :

- The vector space E^T endowed with the product order is an ordered vector space and the vector subspace Z is **order-convex** *i.e.* for every $v, w \in E^T$, $[v, w] := \{x \in E^T, v \leq x \leq w\} \subseteq Z$.

Thus, the reduced power E^T/\mathcal{F} is also an ordered vector space.

- Moreover, if E has an order unit e , then the set :

$$\mathcal{L}_0(E^T/\mathcal{F}) := \{z \in E^T/\mathcal{F} \mid \exists \alpha, \beta \in \mathbb{R} \quad \alpha \text{can}(e) \leq z \leq \beta \text{can}(e)\}$$

is an ordered vector space with order unit $\text{can}(e)$.

A group morphism $f : G \rightarrow \mathbb{R}$ on an abelian ordered group G with positive order unit e is a e -state if f is positive and $f(e) = 1$.

We want to prove the following result :

Theorem 1 : in **ZF**, **HB** implies the following statement S_g

S_g : For every abelian ordered group G with a positive order unit e and every subgroup H of G such that $e \in H$, every e -state on H can be extended into a e -state on G .

The proof is in two steps : first, extending to “one dimension” and then extending to G .

Step 1 : extending to one dimension, in ZF

Let G be an abelian ordered group, H be a subgroup of G and $f : H \rightarrow \mathbb{R}$ a positive group morphism on H .

Extending to one dimension : If H is **cofinal** (i.e. for every $x \in G$, there exists $y \in H$ such that $x \leq y$) and if $x \in G$, we consider :

- $p_H(x) = \sup \left\{ \frac{f(y)}{m} \mid m \in \mathbb{N}^*, y \in H, y \leq mx \right\} \in \mathbb{R}$
- $r_H(x) = \inf \left\{ \frac{f(z)}{n} \mid n \in \mathbb{N}^*, z \in H, nx \leq z \right\} \in \mathbb{R}$

Remark : $p_H(x) \leq r_H(x)$.

Lemma 2 (Goodearl [2], extending to one dimension)

Let G be an abelian ordered group, H be a cofinal subgroup of G and $f : H \rightarrow \mathbb{R}$ a positive group morphism on H . Let $x \in G$:

- 1 *For every positive group morphism $g : H + \mathbb{Z}x \rightarrow \mathbb{R}$ extending f , we have : $p_H(x) \leq g(x) \leq r_H(x)$.*
- 2 *For every $t \in [p_H(x), r_H(x)]$, it is possible to extend f to a positive group morphism $g : H + \mathbb{Z}x \rightarrow \mathbb{R}$ such that $g(x) = t$.*

Corollary 2 : extending to a finite number of dimensions

Let G be an abelian ordered group, H be a cofinal subgroup of G and $f : H \rightarrow \mathbb{R}$ a positive group morphism on H . Let $x_1, \dots, x_n \in G$. There exists a positive group morphism $g : H + \mathbb{Z}x_1 + \dots + \mathbb{Z}x_n$ extending f such that $p_H \leq g \leq r_H$.

Proof : apply the preceding Lemma and remark that if H_1 is a subgroup of G such that $H \subseteq H_1$, $p_H \leq p_{H_1} \leq r_{H_1} \leq r_H$.

Step 2 : Proof of Theorem 1 *i.e.* $\mathbf{HB} \Rightarrow \mathbf{S}_g$

Extending to \mathbf{G} : Let G be an abelian ordered group with positive order unit e , H a subgroup of G such that $e \in H$ (then H is cofinal) and $f : H \rightarrow \mathbb{R}$ a e -state on H .

1. Concurrent relation :

- Denote by $\mathcal{P}_{fin}(G)$ the set of finite subsets of G and $T := \{g \in \mathbb{R}^G \mid p_H \leq g \leq r_H\}$.
- Let \mathcal{R}_f be the binary relation defined by $\forall (F, g) \in \mathcal{P}_{fin}(G) \times T$:

$$\mathcal{R}_f(F, g) : \begin{cases} \forall a, b \in F (a + b \in F \Rightarrow g(a + b) = g(a) + g(b)) \\ \forall a \in F (-a \in F \Rightarrow g(-a) = -g(a)) \\ \forall a, b \in F (a \leq b \Rightarrow g(a) \leq g(b)) \\ \forall a \in F (a \in H \Rightarrow g(a) = f(a)) \\ \forall a \in F p_H(a) \leq g(a) \leq r_H(a) \end{cases}$$

- Using Corollary 2, we prove that if $F \in \mathcal{P}_{fin}(G)$ there exists $g \in T$ extending f ; then $\mathcal{R}_f(F) \neq \emptyset$.
- \mathcal{R}_f is concurrent because if $F_1, \dots, F_n \in \mathcal{P}_{fin}(G)$ then $\emptyset \neq \mathcal{R}_f(F_1 \cup \dots \cup F_n) \subseteq \mathcal{R}_f(F_1) \cap \dots \cap \mathcal{R}_f(F_n)$.
- Thus, we consider the filter \mathcal{F} on T generated by the sets $\mathcal{R}_f(F)$, $F \in \mathcal{P}_{fin}(G)$.

Step 2 : Proof of Theorem 1 *i.e.* $\mathbf{HB} \Rightarrow \mathbf{S}_g$ (cont'd)

2. Reduced power of \mathbb{R} :

- Consider the reduced power \mathbb{R}^T/\mathcal{F} (if $z \in \mathbb{R}^T$, we note \bar{z} the class of z in \mathbb{R}^T/\mathcal{F}) and $\text{can} : \mathbb{R} \rightarrow \mathbb{R}^T/\mathcal{F}$ the canonical embedding.

- Consider $\varphi : G \rightarrow \mathbb{R}^T/\mathcal{F}$
 $x \mapsto \frac{\mathbb{R}^T/\mathcal{F}}{(g(x))_{g \in T}}$: φ is a positive group morphism.

- For all $x \in G$, $\varphi(x) \in \mathcal{L}_0(\mathbb{R}^T/\mathcal{F})$ because :

$$\forall g \in T \quad r_H \leq g \leq p_H$$

Then :

$$\forall x \in G \quad r_H(x) \text{can}(1) \leq \varphi(x) \leq p_H(x) \text{can}(1)$$

First positive group morphism

$$\varphi : G \rightarrow \mathcal{L}_0(\mathbb{R}^T/\mathcal{F})$$

3. Use of \mathbf{HB} :

- **Normed vector space $\mathcal{L}_0(\mathbb{R}^T/\mathcal{F})/N$:**

- $\mathcal{L}_0(\mathbb{R}^T/\mathcal{F})$ is an ordered vector space with an order unit $e_1 := \text{can}(1)$.
- Thus it is endowed with a semi-norm $\| \cdot \|_{e_1}$.
- Let N be the vector subspace $N := \{x \in \mathcal{L}_0(\mathbb{R}^T/\mathcal{F}) \mid \|x\|_{e_1} = 0\}$.
- The quotient vector space $\mathcal{L}_0(\mathbb{R}^T/\mathcal{F})/N$ is endowed with the associated quotient norm.

- Apply **HB** (Corollary 1) to $\mathcal{L}_0(\mathbb{R}^T/\mathcal{F})/N$: there exists a linear form $\psi : \mathcal{L}_0(\mathbb{R}^T/\mathcal{F})/N \rightarrow \mathbb{R}$ continuous of norm 1 such that $\psi(e_1 + N) = \|e_1 + N\| = 1$.

Then, $\Gamma : \mathcal{L}_0(\mathbb{R}^T/\mathcal{F}) \rightarrow \mathbb{R}$ is continuous of norm 1
 $z \mapsto \psi(z + N)$

and $\Gamma(e_1) = 1$: with Lemma 1, Γ is a e_1 -state.

- $\Gamma \circ \text{can} = \text{Id}_{\mathbb{R}}$.

Second positive group morphism

$$\Gamma : \mathcal{L}_0(\mathbb{R}^T/\mathcal{F}) \rightarrow \mathbb{R}$$

4. Existence of state on G :

Extension of f

$$\tilde{f} := \Gamma \circ \varphi : G \rightarrow \mathbb{R}$$

- \tilde{f} is a e-state.
- \tilde{f} extends f because if $x \in H$ then :
 - $\tilde{f}(x) = \Gamma \circ \varphi(x) = \Gamma(\overline{(g(x))_{g \in T}})$.
 - But $\overline{(g(x))_{g \in T}} = \text{can}(f(x))$ because $\mathcal{R}_f(\{x\}) \subseteq \{g \in T \mid g(x) = f(x)\} \in \mathcal{F}$.
 - Then $\tilde{f}(x) = \Gamma(\text{can}(f(x))) = f(x)$ because $\Gamma \circ \text{can} = \text{Id}_{\mathbb{R}}$

We worked on several structures : abelian ordered group with positive order unit, real vector spaces with positive order unit, or unital C^* -algebras.

Given an abelian ordered group G (resp. a real ordered vector space E) with a positive order unit e , a *pure state* on G (resp. on E) is an extreme point of the convex set of e -states on G (resp. on E).


Question

Which consequence of axiom of choice do we need to prove the existence of states or pure states on ordered groups or ordered vector spaces with order unit ?

Consider the two other following weak forms of the Axiom of Choice :


- **KM** (*Krein-Milman axiom*) : Let K be a non-empty compact convex subset of a topological locally convex Hausdorff real vector space X . Then K has an extreme point.
- **T2** (*Tychonov's axiom*) : For every family $(X_i)_{i \in I}$ of compact Hausdorff spaces, the product $\prod_{i \in I} X_i$ is compact.

We have the following diagram :



We obtained the following results :

Diagram : states and axioms of choice



Thank you for listening.



G. Buskes and ACM van Rooij.

A Note on the Gelfand-Naimark-Segal Theorem.

In *Proceedings of the Royal Irish Academy. Section A: Mathematical and Physical Sciences*, pages 143–144. JSTOR, 1993.



K. R. Goodearl.

Partially ordered abelian groups with interpolation, volume 20 of *Mathematical Surveys and Monographs*.

American Mathematical Society, Providence, RI, 1986.



Paul Howard and Jean E. Rubin.

Consequences of the axiom of choice, volume 59 of *Mathematical Surveys and Monographs*.

American Mathematical Society, Providence, RI, 1998.

With 1 IBM-PC floppy disk (3.5 inch; WD).



W. A. J. Luxemburg.

Reduced powers of the real number system and equivalents of the Hahn-Banach extension theorem.

In *Applications of Model Theory to Algebra, Analysis, and Probability (Internat. Sympos., Pasadena, Calif., 1967)*, pages 123–137. Holt, Rinehart and Winston, New York, 1969.