

Marine Biodiversity of La Reunion Island: Echinoderms

Chantal Conand, Sonia Ribes-Beaudemoulin, Florence Trentin, Thierry Mulochau, Emilie Boissin

► **To cite this version:**

Chantal Conand, Sonia Ribes-Beaudemoulin, Florence Trentin, Thierry Mulochau, Emilie Boissin. Marine Biodiversity of La Reunion Island: Echinoderms. Western Indian Ocean Journal of Marine Science, Western Indian Ocean Marine Science Association, 2018, 17 (1), pp.111-124. hal-01906874

HAL Id: hal-01906874

<https://hal.univ-reunion.fr/hal-01906874>

Submitted on 27 Oct 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Marine Biodiversity of La Reunion Island: Echinoderms

Chantal Conand^{1,2}, Sonia Ribes-Beaudemoulin³, Florence Trentin⁴, Thierry Mulochau⁵, Emilie Boissin⁶

¹ UMR ENTROPIE, La Reunion University, Saint Denis, France

² Muséum National Histoire Naturelle, Paris, France

³ Conservatrice Museum de La Reunion, France

⁴ Vie Océane, La Reunion, France

⁵ BIORECIF La Reunion, France

⁶ EPHE-UPVD-CNRS, USR3278-CRIOBE, Laboratoire d'Excellence «CORAIL», Perpignan, France

* Corresponding author: chantal.conand@univ-reunion.fr

Abstract

The inventories of marine species of La Reunion (South West Indian Ocean) are still incomplete for several phylum including the Echinodermata. The present knowledge for the five classes is reported here, in the habitats below (1) and above (2) 50 m depth. The overall diversity for (1) is 136 species and for (2), 48 species. On the whole, as in the broader western Indian Ocean, the Holothuroidea and the Ophiuroidea are the dominant classes. Several new species have been described and many still need further research.

Keywords: sea stars, brittle stars, sea urchins, sea cucumbers, feather stars, SWIO

Introduction

The South West Indian Ocean (SWIO) is recognized as one of the marine biodiversity hotspots in the world, because of high species richness and endemism (Roberts *et al.*, 2002). While fishes and corals have long been well documented from this area, other groups such as Echinoderms needed further attention. Echinoderms are a very distinct five-rayed phylum, very important in marine communities where they play critical roles (Purcell *et al.*, 2016). The five extant classes, namely Asteroidea (sea stars), Ophiuroidea (brittle stars), Echinoidea (sea urchins), Holothuroidea (sea cucumbers) and Crinoidea (feather stars) account for more than 7 000 described species for the world's oceans (Byrne and O' Hara, 2017).

Previous studies had been conducted in several countries or islands of the SWIO region, on one class, as for example for the holothurians (Conand and Muthiga, 2007; Muthiga and Conand, 2014; Conand, 2008; Eriksson *et al.*, 2015; Samyn *et al.*, 2006; Samyn and

Tallon, 2005; FAO, 2013), for the ophiuroids (Stohr *et al.*, 2008; Hoareau *et al.*, 2013; Boissin *et al.*, 2016; Boissin *et al.*, 2017), and the asterids (Jangoux and Aziz, 1988; O'Loughlin and McKenzie, 2013). Other studies have focused on the whole phylum in the region (Clark and Rowe, 1971; Rowe and Richmond, 2011), or groups of islands, such as the French programme BIORECIE on the Îles Eparses (or Scattered islands) (Conand *et al.*, 2010; Mulochau *et al.*, 2014; Conand *et al.*, 2015).

The current knowledge on the diversity of the Echinoderms from La Reunion are reported in this paper. This young volcanic island (21.115°S and 55.536°E), is situated 800 km east of Madagascar. Its marine biodiversity needs to be evaluated given the increases of natural and man-induced threats on the narrow fringing reefs, and the high density of human population. The Echinoderms have been studied during the last forty years and several publications report the ecology of reef populations for several abundant species of the different classes.

The whole phylum has also been focused on in different programmes conducted during this period, such as BIOTAS (ANR-06-BDIV-002) (Hoareau *et al.* 2013; Boissin *et al.*, 2016; Boissin *et al.*, 2017), and BIOLAVE, on the underwater lava flows of the volcano Piton de la Fournaise (Bollard *et al.*, 2013). Recently, the book 'Oursins, Etoiles de mer & autres échinodermes' (2016) has been published in French by the authors of the present paper, with many photos and details. It is important to present this updated inventory to the scientific audience and therefore assist in identifying new species and allow comparisons with other islands or countries in the SWIO.

Methods

The data from the previous publications on the different classes have been gathered and updated.

The first inventory in English is presented here, for the five Echinoderm classes, taking into account the littoral zone, to about 50 m depth, explored by free or scuba diving (1), and deeper zones on the reef-slopes to about 1 000 m (2) which were earlier explored during different cruises by dredging, such as 'MD32' with the 'Marion Dufresne' (1982), and the La Reunion Aquarium team with the 'Explorer' (2011-2014). The species validity has been checked using Miller *et al.* (2017), Stöhr *et al.* (2016) and WoRMS (2018).

Results

Echinoderm diversity

The overall diversity for the Echinoderms of La Reunion is presented in Table 1.

The total number of species presently identified is 184; 4 specimens are identified at the genus level.

The total number of littoral species (1) is 136, while the number of the deep species (2) is 48.

Diversity by classes

Asteroidea diversity

Table 2 presents the list of the Asteroidea collected and/or observed. The forty-six species belong to six orders and sixteen families. A few specimens from the family Ophidiasteridae, *Leiaster* and *Ophidiaster*, remain undetermined.

Twenty-two species are from littoral zones (Table 2 (1)) collected on the coral reefs or from other sandy and rocky biotopes, but are mostly from sparse individuals. During the BIOLAVE programme, 6 species were collected, including the recently described *Aquilonastra conandae* (Bollard *et al.*, 2013). An *Anthenoides* aff. *marleyi* has been found very well conserved in the stomach of a tiger shark *Galeocerdo cuvier* caught at 25 m depth; it was probably recently ingested as it was intact, while the other preys were partly digested.

A few species have been studied in more detail:

1) *Acanthaster planci* now *mauritiensis* (see discussion). Following observations by divers, a study was undertaken in 1998 based on the results of an inquiry distributed to volunteers. From the 352 forms returned during two years, the observations by Emeras *et al.* (2004) concluded that: 1) the distribution was mostly on the western coast, with 17% on reef flats, 83% on outer slopes, rarely seen under 20 meters; 2) the population densities were low (less than 4 individuals per 30 minutes dive) and declining from year 2000; 3) the modal size of the individuals was large (40 cm diameter). Only anecdotal observations have been made on the species in La Reunion since that study.

Table 1. Echinodermata from La Réunion. Number of species in each class in the littoral zone (1) and the deeper zone (2). 'Ni' is number of genera with unidentified species.

	No. spp Littoral (1)	Ni spp Littoral	No. spp Deep (2)	Ni spp Deep	Total + Ni
Asteroidea	22	1	24	1	46 +2
Ophiuroidea	38	1	16		54 +1
Echinoidea	34		7		41
Holothuroidea	38	1			38 +1
Crinoidea	4		1		4 +1
Total	136	3	48	1	184 +4

Table 2. Asteroidea from La Réunion.

Order	Family	Species	Depth
Brisingida	Brisingidae	<i>Brisinga aff. panopla</i> Fisher, 1906	2
Forcipulatida	Asteriidae	<i>Coronaster volsellatus</i> (Sladen, 1889)	1
	Zoroasteridae	<i>Zoroaster carinatus</i> Alcock, 1893	2
Notomyotida	Benthopectinidae	<i>Cheiraster Reunionensis</i> Jangoux & Aziz, 1988	2
		<i>Astropecten bengalensis</i> (Döderlein, 1917)	2
		<i>Astropecten exilis</i> Mortensen, 1933	2
		<i>Astropecten granulatus natalensis</i> John, 1948	2
		<i>Astropecten hemprichi</i> Müller & Troschel, 1842	2
Paxillosida	Astropectinidae	<i>Astropecten leptus</i> H.L. Clark, 1926	2
		<i>Astropecten longibrachius</i> Jangoux & Aziz, 1988	2
		<i>Astropecten polyacanthus</i> (Müller & Troschel, 1842)	1
		<i>Ctenophoraster diploctenius</i> Fisher, 1913	2
		<i>Persephonaster exquisitus</i> Jangoux & Aziz, 1988	2
		<i>Luidia avicularia</i> Fisher, 1913	2
Valvatida	Acanthasterdidae	<i>Acanthaster brevispinus</i> Fisher, 1917	2
		<i>Acanthaster mauritiensis</i> de Loriol, 1885*	1
		<i>Aquilonastra conandae</i> (O'Loughlin & Rowe, 2006)	1
		<i>Aquilonastra richmondi</i> (O'Loughlin & Rowe, 2006)	1
	Asterinidae	<i>Aquilonastra samyni</i> (O'Loughlin & Rowe, 2006)	1
		<i>Aquilonastra aff. watersi</i> O'Loughlin & Rowe, 2006	1
	Asteropseidae	<i>Tegulaster leptalacantha</i> (H.L. Clark, 1946)	2
		<i>Asteropsis carinifera</i> (Lamarck, 1816)	1
		<i>Valvaster striatus</i> (Lamarck, 1816)	1
		<i>Anthenoides cristatus</i> (Sladen, 1889)	2
Goniasteridae	<i>Anthenoides aff. marleyi</i> Mortensen, 1925	1	
	<i>Fromia indica</i> (Perrier, 1869)	1	
	<i>Fromia milleporella</i> (Lamarck, 1816)	1	
	<i>Fromia monilis</i> (Perrier, 1869)	1	
	<i>Mediaster ornatus</i> Fisher, 1906	2	
	<i>Ogmaster capella</i> (Müller & Troschel, 1842)	2	
	<i>Plinthaster doederleini</i> (Koehler, 1909)	2	
	<i>Stellaster equestris</i> (Retzius, 1805)	2	
Leilasteridae	<i>Leilaster spinulosus</i> Aziz & Jangoux, 1985	2	
Mithrodiidae	<i>Mithrodia clavigera</i> (Lamarck, 1816)	1	

Order	Family	Species	Depth
Valvatida	Ophidiasteridae	<i>Cistina columbiae</i> (Gray, 1840)	1
		<i>Dactylosaster cylindricus</i> (Lamarck, 1816)	1
		<i>Ferdina flavescens</i> Gray, 1840	1
		<i>Linckia guildingi</i> Gray, 1840	1
		<i>Linckia multifora</i> (Lamarck, 1816)	1
	Oreasteridae	<i>Nardoa variolata</i> (Retzius, 1805)	1
		<i>Astrosarkus idipi</i> Mah, 2003	1
		<i>Culcita schmideliana</i> (Retzius, 1805)	1
	Poraniidae	<i>Marginaster paucispinus</i> Fisher, 1913	2
		<i>Calyptraster gracilis</i> Jangoux & Aziz, 1988	2
Velatida	Pterasteridae	<i>Euretaster cribrosus</i> (von Martens, 1867)	2
		<i>Hymenaster aff. bartschi</i> Fisher, 1916	2

**Acanthaster mauritiensis* replaces *Acanthaster planci* (see text)

2) *Acanthaster brevispinus* has been observed and filmed in 2001, as a dense population at 40 m depth at the foot of the reef-slope of St Gilles. Despite several dives at the site afterwards, it has not been seen again, suggesting a rapid migration.

3) Several species of Asterinidae have been collected on La Reunion reef flats. A first study was undertaken to determine the modalities of the asexual reproduction by fission and regeneration of an abundant species from the outer reef flat (Kojadinovic *et al.*, 2004) which was later described as *Aquilonastra conandae* (O'Loughlin and Rowe, 2006) and is endemic to the island. These authors identified three other species of *Aquilonastra* from La Reunion also present in the WIO. The sexual reproduction of *Aquilonastra conandae* shows a seasonal gametogenesis with large oocytes suggesting direct development (Ooka *et al.*, 2010).

Half of the species come from deeper zones (Table 2 (2)).

Several species were collected during the MD2 cruise in 1982 and identified by Jangoux and Aziz (1988). They described several new species, as *Cheiraster Reunionensis*, *Astropecten longibrachius*, *Persephonaster exquisitus*, *Leilaster spinulosus*, and *Calyptraster gracilis*.

Echinoidea diversity

Table 3 presents the list of the Echinoidea collected

and/or observed. The thirty-four species belong to eight orders and sixteen families. Most have been collected in littoral zones on coral reefs and sandy bays (Table 3 (1)). During the BIOLAVE programme, 10 species were identified (Bollard *et al.*, 2013).

The biology of some abundant populations important in ecosystem functioning has been studied in detail.

1) *Echinometra mathaei* and *Echinometra mathaei oblonga* present dense populations which play an important role as grazers in the carbonate budget. In La Reunion, they are major eroders on the outer reef flat at one site, with $\text{CaCO}_3 = 8\text{ kg m}^{-2}$. Y-1 (Conand *et al.*, 1997b). A further comparison with other sites of different eutrophication levels shows large differences in urchin sizes and densities and therefore in bioerosion rates, and a clear gradient from the back-reef to the outer reef on the non-degraded sites (Conand *et al.*, 1998; Peyrot-Clausade *et al.*, 2000).

2) *Tripneustes gratilla* is one of the most common herbivorous sea urchin on La Reunion reefs with densities up to 5 individuals. m^{-2} (Naim *et al.*, 1997). It was selected for studying carbon and nitrogen cycling. Its feeding rhythms showed two periods of maximum ingestion (before dawn and after sunset) and a minimum near midday (Lison de Loma *et al.*, 1999). Its diet was mainly algae, with a strong selectivity for *Turbinaria ornata* and avoidance for other species, but with differences between sites (Lison de Loma *et al.*, 2002).

Table 3. Echinoidea from La Réunion.

Order	Family	Species	Depth	
Cidaroida	Cidaridae	<i>Eucidaris metularia</i> (Lamarck, 1816)	1	
		<i>Phyllacanthus imperialis</i> (Lamarck, 1816)	1	
		<i>Stereocidaris indica</i> Döderlein, 1901	2	
		<i>Stylocidaris badia</i> (H.L. Clark, 1925)	2	
Echinothurioida	Echinothuriidae	<i>Asthenosoma marisrrubri</i> Weinberg and de Ridder 1998 *	1	
		<i>Astropyga radiata</i> (Leske, 1778)	1	
		<i>Diadema setosum</i> (Leske, 1778)	1	
		<i>Diadema savignyi</i> (Michelin, 1845)	1	
		<i>Diadema paucispinum</i> (Agassiz, 1863)	1	
		<i>Echinothrix calamaris</i> (Pallas, 1774)	1	
		<i>Echinothrix diadema</i> (Linnaeus, 1758)	1	
		<i>Lissodiadema lorioli</i> Mortensen, 1903	1	
Camarodonta	Echinometridae	<i>Colobocentrotus (Podophora) atratus</i> (Linnæus, 1758)	1	
		<i>Echinometra mathaei</i> (Blainville, 1825)	1	
		<i>Echinometra oblonga</i> (Blainville, 1825)	1	
		<i>Echinostrephus molaris</i> (Blainville, 1825)	1	
		<i>Heterocentrotus mamillatus</i> (Linnaeus, 1758)	1	
		<i>Heterocentrotus trigonarius</i> (Lamarck, 1816)	1	
		<i>Nudechinus verruculatus</i> (Lütken, 1864)	1	
Stomopneustoida	Toxopneustidae	<i>Pseudoboletia maculata</i> Troschel, 1869	1	
		<i>Pseudoboletia indiana</i> (Michelin, 1862)	1	
		<i>Toxopneustes pilleolus</i> (Lamarck, 1816)	1	
		<i>Tripneustes gratilla</i> (Linnæus, 1758)	1	
		<i>Stomopneustes variolaris</i> (Lamarck, 1816)	1	
Spatangoida	Calymnidae	<i>Sternopatagus sibogae</i> de Meijere, 1904	2	
		Brissidae	<i>Brissopsis luzonica</i> (Gray, 1851)	1
			<i>Brissus latecarinatus</i> (Leske, 1778)	1
	<i>Metalia dicrana</i> H.L. Clark, 1917		1	
	Loveniidae	<i>Metalia spatagus</i> (Linnæus, 1758)	1	
		<i>Metalia sternalis</i> (Lamarck, 1816)	1	
		<i>Lovenia elongata</i> (Gray, 1845)	2	
		<i>Maretia planulata</i> (Lamarck, 1816)	1	
	Echinoneoida	Schizasteridae	<i>Schizaster gibberulus</i> L. Agassiz & Desor, 1847	1
			<i>Echinoneus cyclostomus</i> Leske, 1778	1

Order	Family	Species	Depth
		<i>Sculpsitechinus auritus</i> (Leske, 1778)	2
	Astriclypeidae	<i>Echinodiscus bisperforatus</i> Leske, 1778	1
	Clypasteridae	<i>Clypeaster reticulatus</i> (Linnaeus, 1758)	1
Clypasteroida	Echinocyamidae	<i>Echinocyamus megapetalus</i> H.L. Clark, 1914	1
	Laganidae	<i>Jacksonaster depressum</i> (L. Agassiz, 1841)	1
		<i>Laganum decagonale</i> (Blainville, 1827)	2
	Periscomidae	<i>Pericosmus macronesius</i> Koehler, 1914	2

* *Asthenosoma varium* (Grube 1868) has been replaced by *A. marisrubri* after checking by the authors

3) *Colobocentrotus (Podophora) atratus* dense populations on the wave swept intertidal basaltic rocks (Santos and Flammang, 2008); the annual reproduction is during the warm season but its recruitment in this extreme environment needs more studies (Conand, 2001).

4) A few Echinothuriid *Asthenosoma* were collected, they were first listed as *A. varium* Grube, 1868, but are probably *A. marisrubri* (Weinberg and De Ridder, 1998), a species described as endemic to the Red Sea; one specimen was found at 80m depth near St Gilles, one juvenile came from BIOLAVE; it has also been collected near Madagascar during the MIRIKY cruise. These observations extend its distribution in the WIO significantly.

A few species came from local dredging at depths of over 100 meters (Table 3 (2), but no data from MD32 has yet been published.

Ophiuroidea diversity

Table 4 presents the list of the Ophiuroidea collected and/or observed. The fifty-four species belong to seven orders and thirteen families.

In La Reunion, this class was first been studied by Guille & Ribes (1981) who reported 21 species associated with scleractinian corals from La Saline on the west coast. In 1984, 20 species collected from deep water by the 'Marion Dufresne' cruise in 1982 were reported by Vadon & Guille (1984). The only species in common between these two studies was *Ophirolepis irregularis*. More recently, non-focal sampling and the description of *Ophiocanops multispina* Stöhr, Conand & Boissin, 2008 raised the known fauna to 45 species, 26 of which were recorded from shallow waters (Stöhr *et al.*, 2008). From the BIOLAVE programme, 13 species were identified from 8 genera

including juveniles, which made the ophiuroids the more diverse class of echinoderms in Reunion (Bollard *et al.*, 2013). Recently, Boissin *et al.* (2016) presented the results of extensive sampling in shallow water reef ophiuroids and a DNA barcoding study of SWIO brittle-stars revealed that up to 20% of ophiuroid biodiversity might still be unknown (Boissin *et al.*, 2017). The new classification of higher taxa in Ophiuroidea by O'Hara *et al.* (2018) has been followed in this paper. Noticeably, regarding the superorder Ophintegrida, *Ophiopeza* is in a new family Ophiopezidae that belongs together with Ophiocomidae, Ophiodermatidae and Ophiomyxidae to the order Ophiacanthida, suborder Ophiodermatina. Ophiotrichidae, Ophiactidae and Amphiuridae belong to the order Amphilepidida, suborder Gnathophiurina, while Ophionereididae and Ophiolepididae belong to the suborder Ophionereidina. Regarding the superorder, Euryophiurida, Gorgonocephalidae and Euryalidae belong to the order Euryalida, while Ophiuridae belongs to Ophiurida.

The littoral species are presented in Table 4 (1) and the deeper species in Table 4 (2).

Holothuroidea diversity

Table 5 presents the list of the littoral Holothuroidea collected and/or observed. The thirty-eight species belong to four orders and five families. Several specimens from the genera *Holothuria (Stauropora)*, *Stichopus*, *Leptosynapta* and *Polylectana* are not yet determined to species level.

Several programmes, first supported by The Regional Council of La Reunion (Conand & Mangion, 2002; Conand *et al.*, 2003), then WIOMSA (Conand and Frouin, 2007), BIOLAVE where only 5 species were identified (Bollard *et al.*, 2013), and BIOTAS (Conand *et al.*,

Table 4. Ophiuroidea from La Réunion.

Order	Family	Species	Depth
Euryalida	Euryalidae	<i>Asterostegus tuberculatus</i> Mortensen, 1933	2
	Gorgonocephalidae	<i>Astroboa nuda</i> (Lyman, 1874)	1
Ophiurida	Ophiuridae	<i>Amphiophiura bullata convexe</i> (Lyman, 1878)	2
		<i>Amphiophiura paupera</i> (Koehler, 1897)	2
		<i>Amphiophiura sculptilis</i> (Lyman, 1878)	2
		<i>Anthophiura ingolfi</i> Fasmer, 1930	2
		<i>Ophiomastus platydiscus</i> H.L. Clark, 1939	2
		<i>Ophiophyllum borbonica</i> Vadon & Guille, 1984	2
		<i>Ophioplinthus abyssorum</i> (Lyman, 1883)	2
		<i>Ophiotypa simplex</i> Koehler, 1897	2
		<i>Ophiura aequalis</i> (Lyman, 1878)	2
		<i>Ophiura irrorata</i> (Lyman, 1878)	2
		<i>Ophiura irrorata loveni</i> (Lyman, 1878)	2
		<i>Ophiura kinbergi</i> Ljungman, 1866	2
		<i>Perlophiura profundissima</i> Belyaev & Litvinova, 1972	2
		<i>Uriopha ios</i> Paterson, 1980	2
Amphilepidida	Amphiuridae	<i>Amphioplus</i> sp.	1
		<i>Amphipholis squamata</i> (Delle-Chiaje, 1828)	1
		<i>Amphiura crispa</i> Mortensen, 1940	1
		<i>Ophiocentrus aspera</i> (Koehler, 1905)	2
		<i>Ophiactis lymani</i> Ljungman, 1872	1
	Ophiactidae	<i>Ophiactis modesta</i> Brock, 1888	1
		<i>Ophiactis picteti</i> (De Loriol, 1893)	1
		<i>Ophiactis quadripina</i> H.L. Clark, 1915	1
		<i>Ophiactis savignyi</i> (Müller & Troschel, 1842)	1
		<i>Ophionereis porrecta</i> Lyman, 1860 sp1	1
Ophionereididae	<i>Ophionereis porrecta</i> Lyman, 1860 sp2	1	
	<i>Macrophiothrix</i> aff. <i>belli</i> (Döderlein, 1896)	1	
	<i>Macrophiothrix longipeda</i> (Lamarck, 1816)	1	
	<i>Macrophiothrix</i> aff. <i>paucispina</i> Hoggett, 1991	1	
	<i>Macrophiothrix propinqua</i> (Lyman, 1861)	1	
	<i>Macrophiothrix robillardi</i> (De Loriol, 1893)	1	
	<i>Ophiothela</i> aff. <i>danae</i> Verrill, 1869	1	
<i>Ophiothrix foveolata</i> Marktanner-Turneretscher, 1887	1		
<i>Ophiothrix trilineata trilineata</i> Lütken, 1869	1		

Order	Family	Species	Depth
Amphilepidida	Ophiolepididae	<i>Ophiolepis cincta</i> Müller & Troschel, 1842 complex sp1	1
		<i>Ophiolepis cincta</i> Müller & Troschel, 1842 complex sp2	1
		<i>Ophiolepis irregularis</i> Brock, 1888	1
		<i>Ophiolepis superba</i> H.L. Clark, 1915	1
		<i>Ophiomusium luetkeni</i> Lyman, 1878	2
		<i>Ophiomusium lymani</i> Wyville-Thomson, 1873	2
		<i>Ophiomusium scalare</i> Lyman, 1878	2
		<i>Ophioplocus imbricatus</i> Müller & Troschel, 1842	1
		<i>Ophiosphalma fimbriatum</i> (Koehler, 1922)	2
		<i>Ophiosphalma planum</i> (Lyman, 1878)	2
		<i>Ophiacantha funebris</i> (Koehler, 1930)	2
		<i>Ophiacantha pentagona</i> Koehler, 1897	2
		<i>Neoplax ophiodes</i> Bell, 1884	1
		<i>Ophiocanops multispina</i> Stohr Conand et Boissin, 2008	1
<i>Ophiomyxa compacta</i> (Koehler, 1905)	2		
<i>Ophiocoma brevipes</i> Peters, 1851	1		
<i>Ophiocoma cynthiae</i> Benavides-Serrato & O'Hara, 2008	1		
<i>Ophiocoma dentata</i> Müller & Troschel, 1842	1		
<i>Ophiocoma doederleini</i> De Loriol, 1899	1		
<i>Ophiocoma erinaceus</i> Müller & Troschel, 1842	1		
<i>Ophiocoma krohi</i> Stöhr Boissin & Hoareau, 2013	1		
<i>Ophiocoma pica</i> Müller & Troschel, 1842	1		
<i>Ophiocoma pusilla</i> (Brock, 1888)	1		
<i>Ophiocoma scolopendrina</i> (Lamarck, 1816)	1		
<i>Ophiocomella sexradia</i> (Duncan, 1887)	1		
<i>Ophiopsila pantherina</i> Koehler, 1898	1		
<i>Ophiarachnella aff. gorgonia</i> (Müller & Troschel, 1842)	1		
<i>Ophiarachnella septemspinosa</i> (Müller et Troschel, 1842)	1		
<i>Ophioconis cupida</i> Koehler, 1905	1		
<i>Ophiopeza fallax fallax</i> Peters, 1951	1		
<i>Ophiopeza spinosa</i> (Ljungman, 1867)	1		

Table 5. Holothuroidea from La Réunion.

Order	Family	Species	Depth
Aspidochirotida * othuriida *	Holothuriidae	<i>Actinopyga capillata</i> Rowe & Massin, 2006	1
		<i>Actinopyga</i> aff. <i>echinites</i> (Jaeger, 1833)	1
		<i>Actinopyga mauritiana</i> (Quoy & Gaimard, 1834)	1
		<i>Actinopyga</i> aff. <i>obesa</i> (Selenka, 1867)	1
		<i>Bohadschia subrubra</i> (Quoy & Gaimard, 1834)	1
		<i>Bohadschia vitiensis</i> (Semper, 1867)	1
		<i>Holothuria (Cystipus) inhabilis</i> Selenka, 1867	1
		<i>Holothuria (Lessonothuria) insignis</i> Ludwig, 1875	1
		<i>Holothuria (Halodeima) atra</i> Jaeger, 1833	1
		<i>Holothuria (Lessonothuria) lineata</i> Ludwig, 1875	1
		<i>Holothuria (Lessonothuria) pardalis</i> Selenka, 1867	1
		<i>Holothuria (Lessonothuria) verrucosa</i> Selenka, 1867	1
		<i>Holothuria (Mertensiothuria) hilla</i> Lesson, 1830	1
		<i>Holothuria (Mertensiothuria) leucospilota</i> (Brandt, 1835)	1
		<i>Holothuria (Microthele) aff. fuscogilva</i> Cherbonnier, 1980	1
		<i>Holothuria (Microthele) nobilis</i> (Selenka, 1867)	1
		<i>Holothuria (Platyperona) difficilis</i> Semper, 1868	1
		<i>Holothuria (Semperothuria) cinerascens</i> (Brandt, 1835)	1
		<i>Holothuria (Semperothuria) flavomaculata</i> Semper, 1868	1
		<i>Holothuria (Stauropora) fuscocinerea</i> Jaeger, 1833	1
		<i>Holothuria (Stauropora) pervicax</i> Selenka, 1867	1
		<i>Holothuria (Theelothuria) turriscelsa</i> Cherbonnier, 1980	1
		<i>Holothuria (Thymiosycia) arenicola</i> Semper, 1868	1
<i>Holothuria (Thymiosycia) impatiens</i> (Forskål, 1775) complex sp.1	1		
<i>Holothuria (Thymiosycia) impatiens</i> (Forskål, 1775) complex sp.2	1		
<i>Labidodemas pertinax</i> (Ludwig, 1875)	1		
<i>Pearsonothuria graeffei</i> (Semper, 1868)	1		
Synallactida *	Stichopodidae	<i>Stichopus chloronotus</i> Brandt, 1835	1
		<i>Stichopus herrmanni</i> Semper, 1868	1
		<i>Stichopus monotuberculatus</i> (Quoy & Gaimard, 1834)	1
		<i>Stichopus</i> sp	1
		<i>Thelenota ananas</i> (Jaeger, 1833)	1
Apodida	Chiridotidae	<i>Chiridota stuhlmanni</i> Lampert, 1896	1
		<i>Polycheira rufescens</i> (Brandt, 1835)	1

Order	Family	Species	Depth
Apodida	Synaptidae	<i>Euapta godeffroyi</i> (Semper, 1868)	1
		<i>Opheodesoma grisea</i> (Semper, 1867)	1
		<i>Synapta maculata</i> (Chamisso and Eysenhardt, 1821)	1
Dendrochirotida	Sclerodactylidae	<i>Afrocucumis africana</i> (Semper, 1867)	1
		<i>Ohshimella ehrenbergi</i> (Selenka, 1868)	1

* Recent revision by Miller *et al.* 2017

2010) have allowed the biodiversity in La Reunion shallow environments to be recorded. One new species *Actinopyga capillata* has been described from these collections (Rowe and Massin, 2006).

Deeper habitats have not been studied yet.

During recent decades, several studies have also been conducted on the reproductive biology, the ecology and the genetics of several common holothurian species.

The **reproductive biology** has been detailed for several species with dense populations which display seasonal sexual reproduction as well as asexual scission. *Holothuria atra* is the most frequent and abundant Holothuriidae species; the biometry and reproduction have been analyzed at several sites (Conand, 1996; Jaquemet *et al.*, 1999; Conand, 2004). *H. leucospilota* is another abundant black littoral species (Conand *et al.*, 1997a; Gaudron *et al.*, 2008).

The **ecological role** through feeding and bioturbation of these two species has been analysed by Mangion *et al.* (2004). The species were more abundant in eutrophic areas where the mixed populations were able to rework 82 kg dry weight.m⁻².Y⁻¹. *Stichopus chloronotus* is the most frequent and abundant Stichopodiidae on La Reunion reefs. The population parameters

and the reproductive strategies have been studied (Conand *et al.*, 1998; Hoareau and Conand, 2001; Conand *et al.*, 2002.)

The population characteristics of these species and *Actinopyga aff. echinites*, another abundant species, have been summarized in view of regional management efforts of their populations (Kohler *et al.*, 2009).

A first inventory of **diversity** had been prepared for the Regional Council (Conand *et al.*, 2003) and more recent programmes have allowed completion of the inventory for the littoral areas (Conand *et al.*, 2010). Small and cryptic species still need more investigation and deeper populations are not known.

The first genetic data have been collected through a collaboration with Australia on the fissiparous species *S. chloronotus* and *H. atra* (Uthicke *et al.*, 2001; Uthicke and Conand, 2005). The barcoding has been undertaken and established for several commercial species (Uthicke *et al.*, 2010); it will be important in the future for international regulations of CITES or other regulations.

Crinoidea diversity

Table 6 presents the list of the **Crinoidea** collected and/or observed. The five species belong to the order Comatulidae and three families.

Table 6. Crinoidea from La Réunion.

Order	Family	Species	Depth
Comatulida	Guillecrinidae	<i>Guillecrinus Reunionensis</i> Roux, 1985	2
	Colobometridae	<i>Cenometra aff. emendatrix madagascarensis</i> AM Clark, 1972	1
	Mariametridae	<i>Stephanometra indica</i> (Smith, 1876)	1
		<i>Lamprometra palmata</i> (Muller, 1841)	1
	Tropiometridae	<i>Tropiometra aff. carinata</i> (Lamarck, 1816)	1

The littoral species are not very diverse, with only 4 species from 3 families. During the programme BIO-LAVE, 2 of these species were found again on the lava flows (Bollard *et al.*, 2013).

For the deeper species, the MD32 cruise was interesting, as the new species (and new gender and family) *Guillecrinus Réunionensis* was described by Roux (1985) and other specimens are still under study.

Discussion

The different classes show the same proportions as reported by Rowe and Richmond (2011) for the whole WIO, with the dominance of the Holothuroidea and Ophiuroidea. The high diversity of Ophiuroidea comes from many deep species collected during the MD32 cruise. The Holothuroidea, given the increasing commercial value of several species (Conand, 2008), have received much attention, with several local studies in the SWIO. Illegal fisheries, not reported to occur in La Reunion, are an important problem, which needs new tools to be documented and managed (Conand, 2017, 2018). The systematics of the Crinoidea is presently being revised, using integrative taxonomy/morphology coupled with genetics. The Crown-of-Thorns (COT) populations (*Acanthaster*, Asteroidea) deserve special monitoring in the context of climate change and numerous recent outbreaks (Pratchett *et al.*, 2017). The species has previously been reported as *A. planci*, but a recent large-scale study by Haszprunar *et al.* (2017) proposes *A. mauritiensis* de Loriol, 1885 for the WIO. *A. mauritiensis* is therefore used before changes are introduced in the main database WoRMS and Asteroidea (Mah, 2018).

Despite the small size of the reef and deeper habitats of La Reunion, a few new species have been collected and identified: 1 Asteroidea *Aquilonastra conandae*, 1 Holothuroidea *Actinopyga capillata*, 2 littoral Ophiuroidea *Ophiocanops multispina*, and *Ophiocoma krohi*, and several specimens still need further studies.

Many species in each class are new records for La Reunion.

Additional sampling over time in the different habitats will be necessary to follow this diversity and to complete this inventory. Since recent inventories coupled with DNA barcoding in the SWIO have shown that biodiversity might be underestimated in the ophiuroids by 20% (but also in other phyla: e.g. hydroid diversity could be underestimated by two thirds (Boissin *et al.*

2018), further inventories on cryptic habitats and specimens of small size will certainly add to these lists. It is also necessary to conserve and protect these echinoderm species that play such critical ecological roles in coral reefs (Purcell *et al.*, 2016).

Acknowledgements

Thanks to many colleagues for their continuous help in identification, particularly M. Jangoux, C. Mah, S. Stohr, T. O'Hara, C. de Ridder, S. Coppard, F. Michonneau, M. Eléaume.

References

- Boissin E, Hoareau T, Paulay G, Bruggemann H (2016) Shallow-water reef ophiuroids (Echinodermata: Ophiuroidea) of Reunion (Mascarene Islands), with biogeographic considerations. *Zootaxa* 4098: 273-297 [https://doi.org/10.11646/zootaxa.4098.2]
- Boissin E, Hoareau T, Paulay G, Bruggemann H (2017) DNA barcoding of reef brittle stars (Ophiuroidea, Echinodermata) from the southwestern Indian Ocean evolutionary hot spot of biodiversity. *Ecology and Evolution* 7:11197-11203
- Boissin E, Hoareau T, Postaire B, Gravier-Bonnet N, Bourmaud C (2018) Cryptic diversity, low connectivity and suspected human-mediated dispersal among 17 widespread Indo-Pacific hydroid species of the South-western Indian Ocean. *Journal of Biogeography* (in press)
- Bollard S, Pinault M, Quod JP, Boissin E, Hemery L, Conand C (2013) Biodiversity of echinoderms on underwater lava flows with different ages, from the Piton de La Fournaise (Reunion Island, Indian Ocean). *Cahiers De Biologie Marine* 54: 491-497
- Byrne M, O'Hara TD (eds) (2017) Australian Echinoderms: Biology, Ecology & Evolution. CSIRO Publishing, Canberra, 612 pp
- Clark AM, Rowe FWE (1971) Monograph of shallow-water Indo-West Pacific echinoderms. Trustees of the British Museum (Natural History), London, 234 pp
- Conand C (1996) Asexual reproduction by fission in *Holothuria atra*: Variability of some parameters in populations from the tropical Indo-Pacific. *Oceanologica Acta* 19 (3): 209-216
- Conand C (2001) Biology of a poorly known echinoid, *Colobocentrotus atratus*, inhabiting wave swept intertidal basaltic rocks. In: Barker (ed) Echinoderm 2000, Swets & Zeitlinger, pp 431
- Conand C (2004) Monitoring a fissiparous population of *Holothuria atra* on a fringing reef on Reunion Island (Indian Ocean). S.P. C. Bêche-de-mer Information Bulletin 20: 22-25

- Conand C (2008) Population status, fisheries and trade of sea cucumbers in Africa and Indian Ocean. In: Toral-Granda V, Lovatelli A, Vasconcellos M (eds) Sea cucumbers. A global review on fishery and trade. FAO Fisheries Technical Paper No. 516, Rome, FAO, pp 153-205
- Conand C (2017) Recent trends in the world sea cucumbers fisheries and markets. *Revista Biologia Tropical* 65 (S1): 1-10
- Conand C (2018) Tropical sea cucumber fisheries: changes during the last decade. *Marine Pollution Bulletin* 133: 590-594
- Conand C, Morel C, Mussard R (1997a) A new case of asexual reproduction in holothurians: Fission in *Holothuria leucospilota* populations on Reunion island in the indian ocean. S.P.C. Bêche-de-mer Information Bulletin 9
- Conand C, Chabanet P, Cuet P, Letourneur Y (1997b) The carbonate budget of a fringing reef in La Reunion. International Coral Reefs Congress, Panama, 1: 953-958
- Conand C, Heeb M, Peyrot-Clausade M, Fontaine MF (1998) Evaluations of bioerosion by two types of the sea urchin *Echinometra mathaei*, on several sites of a fringing reef in La Reunion Island (Indian Ocean) and comparison with other sites. In: Mooi R, Telford M (eds). Echinoderms: San Francisco. A.A. Balkema, Rotterdam: 609-615
- Conand C, Uthicke S, Hoareau T (2002) Sexual and asexual reproduction of the holothurian *Stichopus chloronotus* (Echinodermata): a comparison between La Reunion (Indian Ocean) and east Australia (Pacific Ocean). *Invertebrate Reproduction Development*, 41 (1-3): 235-242
- Conand C, Mangion P (2002) Holothurians from La Reunion fringing reefs: diversity, distribution, abundance and population structure. S.P.C. Bêche-de-mer Information Bulletin 17: 27-33
- Conand C, Chabanet P, Gravier-Bonnet N (2003) Biodiversité du milieu récifal réunionnais : Echinodermes, Poissons, Hydraires. Rapport Laboratoire Ecomar au Conseil Régional de La Reunion
- Conand C, Frouin P (2007) Sea cucumbers in La Reunion. In: Conand C, Muthiga N (eds) Commercial Sea Cucumbers: A Review for the Western Indian Ocean. WIOMSA Book Series No. 5, pp 21-29
- Conand C, Muthiga N (2007) Conclusion and recommendations. In: Conand C and Muthiga N (eds) Commercial Sea Cucumbers: A Review for the Western Indian Ocean. WIOMSA Book Series No. 5, pp 57-63
- Conand C, Michonneau F, Paulay G, Bruggemann H (2010) Diversity of the holothuroid fauna (Echinodermata) in La Reunion (Western Indian Ocean). *Western Indian Ocean Journal Marine Science* 9 (2): 145-151
- Conand C, Mulochau T, Stohr S, Eléaume M, Chabanet P (2015) Inventory of echinoderms in the Iles Eparses (Europa, Glorieuses, Juan de Nova), Mozambique Channel, France, *Acta Oecologica* [<http://dx.doi.org/10.1016/j.actao.2015.06.007>]
- Conand C, Ribes Beaudemoulin S, Trentin F, Mulochau T, Boissin E (2016) Oursins, Etoiles de mer & autres échinodermes, Collection Biodiversité de La Reunion. Editions du Cyclone, 168 pp
- Emeras J, Falquet MP, Conand C (2004) *Acanthaster planci* on La Reunion reefs (Western Indian Ocean). *Reef Encounter* 32: 26-27
- Eriksson H, Conand C, Lovatelli A, Muthiga N, Purcell SW (2015) Governance structures and sustainability in Indian Ocean sea cucumber fisheries. *Marine Policy* 56: 16-22
- FAO (2013) Report on the FAO Workshop on Sea Cucumber Fisheries: An Ecosystem Approach to Management in the Indian Ocean (SCEAM Indian Ocean) (FAO Fisheries and Aquaculture Report. No. 1038). FAO, Rome.
- Gaudron S, Kohler S, Conand C (2008) Reproduction of the sea cucumber *Holothuria leucospilota* in the fringing reef of Reunion Island (Western Indian Ocean): biological and ecological aspects. *Invertebrate Reproduction Development* 51 (1): 19-31
- Guille A, Ribes S (1981) Echinodermes associés aux Scléractiniaires d'un récif frangeant de l'île de La Reunion (océan Indien). *Bulletin du Muséum national d'Histoire Naturelle* 3 (1): 73-92
- Hoareau TB, Conand C (2001) Sexual reproduction of *Stichopus chloronotus*, a fissiparous sea cucumber, on Reunion Island, Indian Ocean. S.P.C. Bêche-de-mer Information Bull. 15: 4-12
- Hoareau TB, Boissin E, Paulay G, Bruggemann JH (2013) The Southwestern Indian Ocean as a potential marine evolutionary hotspot: perspectives from comparative phylogeography of reef brittle-stars. *Journal of Biogeography* 40: 2167-2179
- Haszprunar G, Vogler C, Wörheide G (2017) Persistent Gaps of Knowledge for Naming and Distinguishing Multiple Species of Crown-of-Thorns-Seastar in the *Acanthaster planci* Species Complex. *Diversity* (9) 22 [doi: 10.3390/d9020022]
- Jangoux M, Aziz A (1988) Les astéries (Echinodermata) récoltées autour de l'île de la Reunion par le N.O. Marion-Dufresne en 1982. *Bulletin du Muséum national d'histoire naturelle. Section A, Zoologie, biologie et écologie animales* 4 (10): 631-650

- Jaquemet S, Rousset V, Conand C (1999) Asexual reproduction parameters and the influence of fission on a *Holothuria atra* sea cucumber population from a fringing reef on Reunion Island (Indian Ocean). *Bêche-de-mer Information Bulletin* 11: 12-18
- Kohler S, Gaudron S, Conand C (2009) Reproductive biology of *Actinopyga echinites* and other sea cucumbers from Reunion Island (Western Indian Ocean): a contribution for a regional management of the fishery. *Western Indian Ocean Journal Marine Science* 8 (1): 97-111
- Kojadinovic J, Falquet MP, Mangion P, Conand C (2004) Distribution, abundance, and asexual reproduction of *Asterina burtoni* (Asteroidea: Echinodermata) from Reunion reefs (Western Indian Ocean). In: München – Heinzeller, Nebelsick (eds) *Echinoderms*. Taylor & Francis Group, London, pp 225-230
- Lison de Loma T, Harmelin-Vivien M.L, Conand C (1999) Diel feeding rhythm of the sea urchin *Tripneustes gratilla* (L.) on a coral reef at La Reunion, Indian Ocean. In: Candia Carnevali, Bonasoro (eds) *Echinoderm Research 1998*. Balkema, Rotterdam, pp 87-92
- Lison De Loma T, Conand C, Harmelin-Vivien ML, Ballesteros E (2002) Food selectivity of *Tripneustes gratilla* (L.) (Echinodermata: Echinoidea) in oligotrophic and nutrient-enriched coral reefs at La Reunion (Indian Ocean). *Bulletin Marine Science* 70: 927-938
- Mah C. (2018) Asteroid systematics in the 21st century: new questions, phylogeny, taxonomy & natural history. International Echinoderm Conference, Nagoya (plenary lecture, abstract)
- Mangion P, Taddei D, Frouin P, Conand C (2004) Feeding rate and impact of sediment reworking by two deposit feeders *Holothuria leucospilota* and *Holothuria atra* on fringing reef (Reunion Island, Indian Ocean). In: München – Heinzeller, Nebelsick (eds). Taylor & Francis Group, London, pp 311-317
- Mulochau T, Conand C, Stöhr S, Eleaume M, Chabanet P (2014) First Inventory of Echinodermata at Juan de Nova (Iles Eparses, France) in the Mozambique Channel. *Western Indian Ocean Journal Marine Science* 13 (1): 23-30
- Muthiga N, Conand C (eds) (2014) *Sea cucumbers, a Poorly Understood but Important Coastal Resource: A Regional Analysis to Improve Management*. WIOMSA Book Series No. 14, 74 pp
- Miller AK, Kerr AM, Paulay G, Reich M, Wilson NG, Carvajal JI, Rouse GW (2017) Molecular phylogeny of extant Holothuroidea (Echinodermata). *Molecular Phylogenetics and Evolution* 111: 110-131
- Naim O, Cuet P, Letourneur Y (1997) Experimental shift in benthic community structure. 8th International Coral Reef Symposium 2: 1873-1878
- O'Hara TD, Stöhr S, Hugall AF, Thuy B, Martynov A (2018) Morphological diagnoses of higher taxa in Ophiuroidea (Echinodermata) in support of a new classification. *European Journal of Taxonomy* 416: 1–35 [<https://doi.org/10.5852/ejt.2018.416>]
- O'Loughlin M, Rowe F (2006) A systematic revision of the asterinid genus *Aquilonastra* O'Loughlin, 2004 (Echinodermata: Asteroidea) *Memoirs of Museum Victoria* 63 (2): 257-287
- O'Loughlin M, Mackenzie M (2013) Asterinid seastars from the Mozambique Channel (Echinodermata: Asteroidea: Asterinidae). *Zootaxa* 3613 (2): 176-180
- Ooka S, Komatsu M, Conand C (2010) Sexual reproduction of the small fissiparous seastar *Aquilonastra sp.* (Asteroidea: Asterinidae) in La Reunion Island. In: Harris LG, Boetger SA, Walker CW, Lesser MP (eds) *Echinoderms*. Durham, Taylor & Francis, London, pp 467-472
- Peyrot-Clausade M, Chabanet P, Conand C, Fontaine MF, Letourneur Y, Harmelin-Vivien M (2000) Carbonate budget of two Indo-Pacific reefs: sea-urchin and fish bioerosion in La Reunion and Moorea. *Bulletin Marine Science* 66 (2): 477-485
- Pratchett MS, Ciemon F, Caballes JC, Wilmes J, Matthews S, Melin C, Sweatman H, Nadler L, Brodie J, Thompson C, Hoey J, Bos A, Byrne M, Messmer V, Fortunato S, Chen CC, Buck A, Babcock R, (2017) Thirty Years of Research on Crown-of-Thorns Starfish (1986–2016): Scientific Advances and Emerging Opportunities. *Diversity* 9, 41; doi:10.3390/d9040041
- Purcell SW, Conand C, Uthicke S, Byrne M (2016) Ecological roles of exploited sea cucumbers. *Oceanography and Marine Biology: An Annual Review* 54: 367-386
- Roberts CM, McClean CJ, Veron JEN (2002) Marine biodiversity hotspots and conservation priorities for tropical reefs. *Science* 295: 1280-1284
- Roux M (1985) Découverte d'un représentant actuel des crinoïdes pédonculés (Echinodermes) dans l'étage bathyal de l'île de La Reunion. *Compte Rendus Academie Sciences Paris* 301 III, 10: 503-506
- Rowe FEW, Massin C (2006) On a new species of *Actinopyga* Bronn, 1860 (Echinodermata, Holothuroidea) from the Indo-West Pacific. *Zoosystema* 28 (4): 955-961
- Rowe FEW, Richmond M (2011) Phylum Echinodermata. In: Richmond (ed) *A field guide of the Seashores of Eastern Africa and the Western Indian Ocean Islands*. Sida/Wiomsa, 464 pp

- Samyn Y, Tallon I (2005) Zoogeography of the shallow-water holothuroids of the western Indian Ocean. *Journal Biogeography* 32: 1523-1538
- Samyn Y, Van den Spiegel D, Massin C (2006) Taxonomie des holothuries des Comores, vol. 1. *AbcTaxa*, 130 pp
- Santos R, Flammang P (2008) Estimation of the attachment strength of the shingle sea urchin, *Colobocentrotus atratus*, and comparison with three sympatric echinoids. *Marine Biology* [DOI 10.1007/s00227-007-0895-6]
- Stöhr S, Conand C, Boissin E (2008) Brittle stars (Echinodermata: Ophiuroidea) from La Reunion and the systematic position of *Ophiocanops* Koehler, 1922. *Zoological Journal of the Linnean Society* 153: 545-560 [http://dx.doi.org/10.1111/j.1096-3642.2008.00401.x]
- Stöhr S, Boissin E, Hoareau TB (2013) Taxonomic revision and phylogeny of the *Ophiocoma brevipes* group, with description of a new subgenus (*Breviturma*) and a new species. *European Journal of Taxonomy* 68: 1-26 [http://dx.doi.org/10.5852/ejt.2013.68]
- Stöhr S, O'Hara T, Thuy B (eds) (2016) World Ophiuroidea database. <http://www.marinespecies.org/ophiuroidea>
- Uthicke S (2017) Thirty Years of Research on Crown-of-Thorns Starfish (1986-2016): Scientific Advances and Emerging Opportunities *Diversity* 9, 41 [doi:10.3390/d9040041]
- Uthicke S, Conand C, Benzie J (2001) Population genetics of the fissiparous holothurians *Stichopus chloronotus* and *Holothuria atra* (Aspidochirotida): a comparison between Torres Strait and La Reunion. *Marine Biology* 139: 257-265
- Uthicke S, Conand C (2005) Amplified fragment length polymorphism (AFLP) analysis indicates importance of both asexual and sexual reproduction in the fissiparous holothurian *Stichopus chloronotus* (Aspidochirotida) in the Indian and Pacific Ocean. *Coral reefs* 24 (1): 103-111
- Uthicke S, Byrne M, Conand C (2010) Genetic barcoding of commercial Beche-de-Mer species (Echinodermata: Holothuroidea). *Molecular Ecology Resources* 10: 634-646
- Vadon C, Guille A 1984. Les Ophiuridae (Ophiuroidea, Echinodermata) de la campagne MD 32 du 'Marion- Dufresne' autour de l'île de La Reunion. *Bulletin Muséum national d'Histoire Naturelle 4e Series* 6: 583-615
- Weinberg S, de Ridder C (1998) *Asthenosoma marisrubri* n. sp. (Echinodermata, Echinoidea) from the Red Sea. *Beaufortia* 48: 27-46
- WoRMS Editorial Board (2018) World Register of Marine Species. <http://www.marinespecies.org> at VLIZ. [doi:10.14284/170]