



**HAL**  
open science

# Error analysis for the finite element approximation of the Brinkmann-Darcy-Forchheimer model for porous media with mixed boundary conditions

Pierre-Henri Cocquet, Michaël Rakotobe, Delphine Ramalingom, Alain Bastide

► **To cite this version:**

Pierre-Henri Cocquet, Michaël Rakotobe, Delphine Ramalingom, Alain Bastide. Error analysis for the finite element approximation of the Brinkmann-Darcy-Forchheimer model for porous media with mixed boundary conditions. *Journal of Computational and Applied Mathematics*, 2020, 10.1016/j.cam.2020.113008 . hal-02561058v2

**HAL Id: hal-02561058**

**<https://hal.univ-reunion.fr/hal-02561058v2>**

Submitted on 25 Mar 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Error analysis for the finite element approximation of the Darcy-Brinkman-Forchheimer model for porous media with mixed boundary conditions

Pierre-Henri Cocquet<sup>(1),(2)</sup>, Michaël Rakotobe<sup>(1)</sup>, Delphine Ramalingom<sup>(1)</sup>,  
Alain Bastide<sup>(1)</sup>

(1) *Physique et Ingénierie Mathématique pour l'Énergie et l'Environnement (PIMENT),  
Université de la Réunion, 2 rue Joseph Wetzell, 97490 Sainte-Clotilde.*

(2) *Laboratoire des Sciences de l'Ingénieur Appliquées à la Mécanique et au Génie  
Electrique (SIAME), E2S-UPPA, Université de Pau et des Pays de l'Adour, 64000 Pau,  
France*

---

## Abstract

This paper deals with the finite element approximation of the Darcy-Brinkman-Forchheimer equation, involving a porous media with spatially-varying porosity, with mixed boundary condition such as inhomogeneous Dirichlet and traction boundary conditions. We first prove that the considered problem has a unique solution if the source terms are small enough. The convergence of a Taylor-Hood finite element approximation using a finite element interpolation of the porosity is then proved under similar smallness assumptions. Some optimal error estimates are obtained if the solution to the Darcy-Brinkman-Forchheimer model are smooth enough. We end this paper by providing a fixed-point method to solve the discrete non-linear problems and with some numerical experiments to make more precise the smallness assumptions on the source terms and to illustrate the theoretical convergence results.

## Keywords:

Finite element, error analysis, Porous media, Darcy-Brinkman-Forchheimer equations.

---

## 1. Introduction

The Darcy-Brinkman-Forchheimer (DBF) model for porous media is obtained from the incompressible Navier-Stokes equation in a porous media through volume averaging. The latter are then completed with closure models for the unknown terms arising in the volume averaged equations. We refer to [46, 16, 37, 48, 49] for the physical modeling of fluid flows in porous media based on volume averaging of the Navier-Stokes equations and the derivation of the DBF model.

Regarding the mathematical study of the steady-state DBF equation, Kaloni & Guo [27] studied the case where the convective non-linear terms vanishes

and show that this problem has a solution for inhomogeneous Dirichlet boundary conditions which is unique for small enough source terms. The existence and uniqueness of solution to the full non-linear problem with inhomogeneous Dirichlet boundary condition can be found in [42, 41]. It is worth noting that the results from [27, 42, 41] have been obtained with similar techniques. The latter are based on the study of a finite dimensional approximation of the non-linear variational problem whose well-posedness comes from [18, p. 597, Lemma IX.3.1] (see also [45, p. 164, Lemma 1.4]) and next proving that the limit satisfies the variational formulation of the DBF model. The existence of solution for homogeneous Dirichlet boundary conditions when considering a generalized Forchheimer term have been obtained in [47] using the Leray-Schauder theorem and the uniqueness again holds for source terms that are small enough. We emphasize that the need of small data to ensure well-posedness of some non-linear PDE is quite classical and is also required when dealing with incompressible Navier-Stokes equation with homogeneous Dirichlet conditions (see e.g. [45, 18]) or fluid-porous media interface problems [14, 20].

In this paper, we are interested in the finite element approximation of the Darcy-Brinkman-Forchheimer equation with mixed boundary conditions since such setting is involved in many physical applications such as those from [46, 1] (see also [42, p. 39, Section 2.4] and [41, Section 3]). As a result, we first need to study the existence and uniqueness of solution to the DBF problem in this setting. Since the model we are interested in is derived from Navier-Stokes equations, we recall below some works dealing with the existence and uniqueness of solution to incompressible steady-state Navier-Stokes equations in the case of mixed boundary conditions. The regularity of the solutions for three dimensional Lipschitz domain with homogeneous mixed boundary conditions involving Dirichlet on some part of the boundary and, on the other part, a vanishing normal trace together with either zero tangential part of the normal stress tensor or the curl of the velocity can be found in [15]. In the case of polyhedral domains with more general inhomogeneous mixed boundary conditions and non-divergence-free velocity fields, one can find existence, uniqueness and regularity results in [29] (see also [28] for similar results for Stokes flows). Finally, the existence and uniqueness of solution to Navier-Stokes equations with mixed inhomogeneous conditions have been considered in [36] where the inhomogeneous Dirichlet boundary conditions have been handled thanks to the introduction of an additional variable. The latter thus yields a non-linear saddle-point problem very similar to the standard weak-formulation of the incompressible Navier-Stokes equation but with a continuous bilinear form  $b$  that is different from the usual one since it does not only involves the divergence of the velocity but also a surface term taking into account the inhomogeneous Dirichlet condition. Regarding the previously mentioned results, even if the steady state incompressible Navier-Stokes system and the DBF model have a similar structure, we cannot deduce from them the existence and uniqueness of a solution to the DBF problem with mixed boundary condition. A first part of the present paper is then going to be dedicated to prove existence and uniqueness of solution to the DBF problem with mixed boundary conditions.

Regarding the convergence of a finite element approximation to the DBF problem, there is actually a large literature on the analysis of mixed finite element method applied to the (generalized) Darcy-Forchheimer model (see e.g. [34, 25, 40, 26, 38]). Nevertheless, compared to the Darcy-Forchheimer model, the DBF equation also involves a nonlinear convective term and a Laplacian of the velocity and seems, to the best of our knowledge, to have been less studied. We can still refer to [35] where the convergence of finite difference method on a staggered grid applied to the unsteady DBF coupled to a solute transport equation have been obtained. For the steady-state DBF, the case of inhomogeneous Dirichlet boundary condition have been studied in [42] where optimal error estimates have been obtained for smooth enough solution. We would like to emphasize that all the aforementioned convergence results are obtained either in the case of a homogeneous porous media or without considering a discrete version of the porosity field in the discrete problem. Note however that having a finite element approximation of the porosity has its own interest since the latter is usually used in the numerical computations. Another advantage of having a finite element formulation with discrete parameters is when one deals with so-called topology optimization problems (see e.g. [4, 43, 5, 39]) or parametric optimization problem [6, Chapter VI]. Indeed, using discretization of these parameters are needed to define discrete optimization problem and to prove the convergence of discrete optimal solution toward the continuous optimal solutions.

In this paper, we are interested in proving the existence and uniqueness of solution to the Darcy-Brinkman-Forchheimer model with a spatially-varying porosity as well as the convergence of a finite element approximation involving a discrete porosity. The paper is thus organized as follow: we begin to prove an existence and uniqueness result for a DBF model with mixed boundary conditions. A finite element method using a discrete porosity is then investigated for which we prove the convergence as well as optimal error estimates for smooth enough solutions. We study next the convergence of a fixed point iteration used to solve the discrete non-linear problem. Since all our results are obtained by assuming that the source term are small enough, we provide some numerical simulations to get more precise estimates on how small these source term have to be in order for the discrete problem to be solve with the fixed point method. This paper ends with some numerical optimal error estimates.

## 2. Steady-state Darcy-Brinkman-Forchheimer model for porous media

We consider a viscous flow inside a porous medium embedded in a computational domain  $\Omega \subset \mathbb{R}^d$  ( $d = 2, 3$ ). The domain is assumed to be a bounded open set with Lipschitz boundary with outward unitary normal  $\vec{n}$ . We assume the porous media has a spatially varying porosity  $\varepsilon : x \in \Omega \mapsto \varepsilon(x) \in (0, 1]$  and that it is modeled by the Darcy-Brinkman-Forchheimer equation. The latter

can be found in e.g. [21, 30, 47] and in dimensionless form reads as follows

$$\begin{cases} -\operatorname{div} (2\operatorname{Re}^{-1}\varepsilon S(\vec{u}) - \varepsilon \vec{u} \otimes \vec{u}) + \varepsilon \nabla p + \alpha(\varepsilon)\vec{u} + \beta(\varepsilon)\vec{u}|\vec{u}| & = \varepsilon \vec{f}, & \text{in } \Omega, \\ \operatorname{div}(\varepsilon \vec{u}) & = 0, & \text{in } \Omega, \end{cases} \quad (1)$$

where  $\operatorname{Re}$  is the Reynolds number and  $\vec{f}$  is an external force field (e.g. gravity). The tensor  $S(\vec{u})$  is the symmetric part of the Jacobian matrix of the velocity field  $\vec{u}$ . We emphasize that one always has  $\alpha(1) = \beta(1) = 0$  and thus the standard Navier-Stokes equation is recovered for  $\varepsilon = 1$ .

**Remark 1.** *Using the formula*

$$\operatorname{div}(\vec{u} \otimes \vec{v}) = \operatorname{div}(\vec{u})\vec{v} + (\vec{u} \cdot \nabla)\vec{v},$$

together with the incompressibility condition  $\operatorname{div}(\varepsilon \vec{u}) = 0$ , the non-linear term can be written as

$$\operatorname{div}(\varepsilon \vec{u} \otimes \vec{u}) = \operatorname{div}(\varepsilon \vec{u})\vec{u} + \varepsilon (\vec{u} \cdot \nabla)\vec{u} = \varepsilon (\vec{u} \cdot \nabla)\vec{u}.$$

We consider the following set of boundary conditions

$$\begin{cases} \vec{u} = 0 & \text{on } \Gamma_w, \\ \vec{u} = \vec{u}_{\text{in}} & \text{on } \Gamma_{\text{in}}, \\ \varepsilon (2\operatorname{Re}^{-1}S(\vec{u}) - p)\vec{n} = 0 & \text{on } \Gamma_{\text{out}}, \end{cases} \quad (2)$$

where we set  $\partial\Omega = \Gamma_w \cup \Gamma_{\text{in}} \cup \Gamma_{\text{out}}$  where each part correspond respectively to the walls, the inlet and the outlet. We also assume throughout this paper that

$$|\Gamma_w| > 0, |\Gamma_{\text{in}}| > 0, |\Gamma_{\text{out}}| > 0 \text{ and } \overline{\Gamma_{\text{in}}} \cap \overline{\Gamma_{\text{out}}} = \emptyset.$$

The porosity, the Darcy and Forchheimer terms satisfy the next set of assumptions

$$\begin{aligned} \varepsilon &\in L^\infty(\Omega) \cap W^{1,r}(\Omega) \text{ with } r > d \text{ and } 0 < \varepsilon_0 \leq \varepsilon(x) \leq 1 \text{ a.e. in } \Omega \\ s &\in [\varepsilon_0, 1[ \mapsto \alpha(s) \in \mathbb{R}^+ \text{ and } s \in [\varepsilon_0, 1[ \mapsto \beta(s) \in \mathbb{R}^+ \text{ are differentiable.} \\ s &\in [\varepsilon_0, 1[ \mapsto \alpha'(s) \in \mathbb{R}^+ \text{ and } s \in [\varepsilon_0, 1[ \mapsto \beta'(s) \in \mathbb{R}^+ \text{ are bounded.} \end{aligned} \quad (3)$$

We emphasize that (3) are satisfied by many example of Darcy and Forchheimer coefficients one may find in the literature (see e.g. [3, 46, 16, 41]).

### 2.1. Well-posedness of the Darcy-Brinkman-Forchheimer problem

We work with inhomogeneous Dirichlet boundary conditions that needs some special functional spaces to be handled properly. For any bounded open set  $\mathcal{O}$  with Lipschitz boundary, we note  $\Gamma_c \subset \partial\mathcal{O}$  a part of the boundary. The trace space  $H_{00}^{1/2}(\Gamma_c)$  is defined in [32, Theorem 11.7] (see also e.g. [31, 33] where some properties are recalled) and can be obtained as the completion of smooth function with compact support in  $\Gamma_c$  with respect to the norm

$$\|\mu\|_{H^{1/2}(\Gamma_c)}^2 := \|\mu\|_{L^2(\Gamma_c)}^2 + \int_{\Gamma_c} \int_{\Gamma_c} \frac{|\mu(x) - \mu(y)|^2}{|x - y|^d} dx dy.$$

Denoting by  $E_0\mu$  the extension by 0 outside  $\Gamma_c$ , we have that any  $\mu \in H_{00}^{1/2}(\Gamma_c)$  satisfy  $E_0\mu \in H^{1/2}(\partial\mathcal{O})$  with

$$\|E_0(\mu)\|_{H^{1/2}(\partial\mathcal{O})} \leq C \|\mu\|_{H_{00}^{1/2}(\Gamma_c)},$$

for a generic constant  $C > 0$ . As a result, we have the equivalent definition of this trace space

$$H_{00}^{1/2}(\Gamma_c) := \left\{ \mu \in H^{1/2}(\Gamma_c) \mid E_0\mu \in H^{1/2}(\partial\mathcal{O}) \right\}.$$

Moreover, the linear application  $E_0 : \mu \in H_{00}^{1/2}(\Gamma_c) \mapsto E_0(\mu) \in H^{1/2}(\partial\mathcal{O})$  is continuous. We finally emphasize that if  $H_{00}^{1/2}(\Gamma_c)$  is endowed with the next norm

$$\|\mu\|_{H_{00}^{1/2}(\Gamma_c)}^2 := \|\mu\|_{H^{1/2}(\Gamma_c)}^2 + \int_{\Gamma_c} \frac{|\mu(s)|^2}{\text{dist}(s, \partial\Gamma_c)} ds,$$

then it is a Banach space.

We are now in position to give the weak formulation to Problem (1,2). We introduce the following Hilbert spaces

$$\mathbf{X}_1 := \{ \vec{v} \in H^1(\Omega)^d \mid \vec{v}|_{\Gamma_w=0} \}, \quad \mathbf{X} := \{ \vec{v} \in H^1(\Omega)^d \mid \vec{v}|_{\Gamma_w \cup \Gamma_{\text{in}}} = 0 \}.$$

Using Korn inequality, we get that

$$\|\vec{u}\|_{\mathbf{X}} := \|S(\vec{u})\|_{L^2(\Omega)} := \sqrt{\int_{\Omega} S(\vec{u}) : S(\vec{u}) dx},$$

is a norm on either  $\mathbf{X}_1$  or  $\mathbf{X}$  where  $A:B = \text{trace}(AB)$ . We also denote by  $C_K > 0$  the constant such that

$$\|\vec{u}\|_{L^2(\Omega)} \leq C_K \|\vec{u}\|_{\mathbf{X}},$$

and emphasize it only depends on  $\Omega$ . Thanks to the assumptions on  $\Gamma_{\text{out}}$  and  $\Gamma_{\text{in}}$ , any  $\vec{u} \in \mathbf{X}_1$  satisfy  $\vec{u}|_{\Gamma_{\text{in}}} \in H_{00}^{1/2}(\Gamma_{\text{in}})^d$  and we thus consider a source term  $\vec{u}_{\text{in}} \in H_{00}^{1/2}(\Gamma_{\text{in}})^d$  as inlet velocity. Lemma 17 ensures that Problem (1,2) is then equivalent to the following variational formulation

$$\begin{aligned} & \text{Find } (\vec{u}, p) \in \mathbf{X}_1 \times L^2(\Omega) \text{ such that} \\ & \vec{u}|_{\Gamma_{\text{in}}} = \vec{u}_{\text{in}} \text{ and} \\ & \begin{cases} a(\varepsilon; \vec{u}, \vec{v}) + c(\varepsilon; \vec{u}, \vec{u}, \vec{v}) + b(\varepsilon; \vec{v}, p) &= \langle \vec{F}, \vec{v} \rangle_{\mathbf{X}' \times \mathbf{X}}, \quad \forall \vec{v} \in \mathbf{X}, \\ b(\varepsilon; \vec{u}, q) &= 0, \quad \forall q \in L^2(\Omega), \end{cases} \end{aligned} \quad (4)$$

where  $\vec{F} = \varepsilon \vec{f}$  and

$$\begin{aligned} a(\varepsilon; \vec{u}, \vec{v}) &= 2\text{Re}^{-1} \int_{\Omega} \varepsilon S(\vec{u}) : S(\vec{v}) dx + \int_{\Omega} \alpha(\varepsilon) \vec{u} \cdot \vec{v} dx, \\ c(\varepsilon; \vec{u}, \vec{v}, \vec{w}) &= \int_{\Omega} \varepsilon (\vec{u} \cdot \nabla) \vec{v} \cdot \vec{w} + \beta(\varepsilon) |\vec{u}| \vec{v} \cdot \vec{w} dx, \\ b(\varepsilon; \vec{u}, q) &= - \int_{\Omega} q \text{div}(\varepsilon \vec{u}) dx. \end{aligned}$$

To deal with the inhomogeneous Dirichlet condition at the inlet, we introduce an extension  $\vec{V}$  of  $\vec{u}_{\text{in}}$  such that  $\text{div}(\varepsilon \vec{V}) = 0$  whose existence is provided by Lemma 16. From this,  $\vec{u} = \vec{w} + \vec{V}$  where  $\vec{w} \in \mathbf{X}$  satisfy the variational formulation

$$\begin{aligned} & \text{Find } (\vec{w}, p) \in \mathbf{X} \times L^2(\Omega) \text{ such that} \\ & \begin{cases} a(\varepsilon; \vec{w}, \vec{v}) + b(\varepsilon; \vec{v}, p) = \langle G(\varepsilon; \vec{w}), \vec{v} \rangle_{\mathbf{X}' \times \mathbf{X}}, & \forall \vec{v} \in \mathbf{X}, \\ b(\varepsilon; \vec{w}, q) = 0, & \forall q \in L^2(\Omega), \end{cases} \end{aligned} \quad (5)$$

where the non-linear term is defined as

$$\langle G(\varepsilon; \vec{w}), \vec{v} \rangle_{\mathbf{X}' \times \mathbf{X}} = \left\langle \vec{F}, \vec{v} \right\rangle_{\mathbf{X}' \times \mathbf{X}} - c(\varepsilon; \vec{w} + \vec{V}, \vec{w} + \vec{V}, \vec{v}) - a(\varepsilon; \vec{V}, \vec{v}). \quad (6)$$

We are going to study the well-posedness to Problem (5) with a fixed-point approach. Therefore, we begin to study the linear problem

$$\begin{aligned} & \text{Find } (\vec{w}, p) \in \mathbf{X} \times L^2(\Omega) \text{ such that} \\ & \begin{cases} a(\varepsilon; \vec{w}, \vec{v}) + b(\varepsilon; \vec{v}, p) = \left\langle \vec{F}, \vec{v} \right\rangle_{\mathbf{X}' \times \mathbf{X}}, & \forall \vec{v} \in \mathbf{X}, \\ b(\varepsilon; \vec{w}, q) = 0, & \forall q \in L^2(\Omega), \end{cases} \end{aligned} \quad (7)$$

where  $\vec{F} \in \mathbf{X}'$  is some source term. Problem (7) is a standard linear saddle-point problem whose well-posedness has been studied in e.g. [12, II.1, Proposition 1.3], [17, p. 474, Theorem A.56] or [19, p. 59, Theorem 4.1]). Since the bilinear form  $a(\varepsilon; \cdot, \cdot)$  is continuous and coercive, namely

$$\begin{aligned} |a(\varepsilon; \vec{u}, \vec{v})| & \leq \left( C_K^2 \|\alpha(\varepsilon)\|_{L^\infty(\Omega)} + 2\text{Re}^{-1} \right) \|\vec{u}\|_{\mathbf{X}} \|\vec{v}\|_{\mathbf{X}}, \\ a(\varepsilon; \vec{u}, \vec{u}) & \geq 2\text{Re}^{-1} \varepsilon_0 \|\vec{u}\|_{\mathbf{X}}^2, \end{aligned} \quad (8)$$

it only remains to prove the bilinear form  $b$  satisfy an inf-sup condition. The latter is obtained in the next lemma.

**Lemma 2.** *Let  $\gamma > 0$  be the inf-sup constant when  $\varepsilon(x) = 1$  for all  $x \in \Omega$ . Assume that  $\varepsilon \in L^\infty(\Omega) \cap W^{1,r(d)}(\Omega)$ , with  $r(2) > 2$  and  $r(3) = 3$  and that*

$$\forall x \in \Omega, \quad 0 < \varepsilon_0 \leq \varepsilon(x) \leq 1.$$

*Then there exists a constant  $\gamma(\varepsilon) > 0$  such that*

$$\inf_{q \in L^2(\Omega) \setminus \{0\}} \sup_{\vec{u} \in \mathbf{X} \setminus \{\vec{0}\}} \frac{b(\varepsilon; \vec{u}, q)}{\|\vec{u}\|_{\mathbf{X}} \|q\|_{L^2(\Omega)}} \geq \gamma(\varepsilon),$$

*where there is a generic constant  $C > 0$  such that  $\gamma(\varepsilon) > 0$  is given by*

$$\gamma(\varepsilon) = \gamma \frac{1}{C \left( \varepsilon_0^{-1} + \varepsilon_0^{-2} \|\nabla \varepsilon\|_{L^3(\Omega)} \right)}.$$

PROOF. Adapting techniques from [8, p. 6, Eq. (2.13)] to the boundary conditions considered in this paper, the following inf-sup condition can be obtained

$$\inf_{q \in L^2(\Omega) \setminus \{0\}} \sup_{\vec{u} \in \mathbf{X}} \frac{b(1; \vec{u}, q)}{\|\vec{u}\|_{\mathbf{X}} \|q\|_{L^2(\Omega)}} \geq \gamma > 0.$$

From [19, p. 58, Lemma 4.1], the inf-sup condition is equivalent to the statement: for any  $q \in L^2(\Omega)$  there exists a  $\vec{v} = \vec{v}(q) \in \mathbf{X}$  such that

$$b(1; \vec{v}, q) \geq C_1 \|q\|_{L^2(\Omega)}^2 \quad \text{and} \quad \|\vec{v}\|_{\mathbf{X}} \leq C_2 \|q\|_{L^2(\Omega)}, \quad (9)$$

where, in that case,  $\gamma = C_1/C_2$ . Since the application  $\vec{u} \in \mathbf{X} \mapsto \varepsilon \vec{u} \in \mathbf{X}$  is an isomorphism (see [7, p. 3, Lemma 2.1]), we have some  $\vec{u} \in \mathbf{X}$  such that  $\vec{v} = \varepsilon \vec{u}$  and (9) becomes

$$b(1; \varepsilon \vec{u}, q) = b(\varepsilon; \vec{u}, q) \geq C_1 \|q\|_{L^2(\Omega)}^2 \quad \text{and} \quad \|\varepsilon \vec{u}\|_{\mathbf{X}} \leq C_2 \|q\|_{L^2(\Omega)}.$$

To conclude, note that

$$\begin{aligned} \|\vec{u}\|_{\mathbf{X}} &= \left\| \frac{\varepsilon \vec{u}}{\varepsilon} \right\|_{\mathbf{X}} \leq C \left( \varepsilon_0^{-1} + \varepsilon_0^{-2} \|\nabla \varepsilon\|_{L^3(\Omega)} \right) \|\varepsilon \vec{u}\|_{\mathbf{X}} \\ &\leq C_2 \|q\|_{L^2(\Omega)} C \left( \varepsilon_0^{-1} + \varepsilon_0^{-2} \|\nabla \varepsilon\|_{L^3(\Omega)} \right) \end{aligned}$$

from which we get that for any  $q \in L^2(\Omega)$ , there exists a  $\vec{u} = \vec{u}(q) \in \mathbf{X}$  so that

$$b(\varepsilon; \vec{u}, q) \geq C_1 \|q\|_{L^2(\Omega)}^2 \quad \text{and} \quad \|\vec{u}\|_{\mathbf{X}} \leq \widetilde{C}_2 \|q\|_{L^2(\Omega)}.$$

The desired inf-sup condition then follows with

$$\gamma(\varepsilon) = \frac{C_1}{\widetilde{C}_2} = \gamma \frac{1}{C \left( \varepsilon_0^{-1} + \varepsilon_0^{-2} \|\nabla \varepsilon\|_{L^3(\Omega)} \right)}.$$

From (8) and Lemma 2, we can apply [12, II.1, Proposition 1.3] (see also [17, p. 474, Theorem A.56] or [19, p. 59, Theorem 4.1]) to get the existence and uniqueness of solution to (7).

**Theorem 3.** *Problem (7) has a unique solution  $(\vec{w}, p) \in \mathbf{X} \times L^2(\Omega)$  that satisfies*

$$\begin{aligned} \|\vec{w}\|_{\mathbf{X}} &\leq \frac{1}{\alpha_0} \left\| \vec{F} \right\|_{\mathbf{X}'}, \\ \|p\|_{L^2(\Omega)} &\leq \frac{1}{\gamma(\varepsilon)} \left( 1 + \frac{\|a\|}{\alpha_0} \right) \left\| \vec{F} \right\|_{\mathbf{X}'}, \end{aligned}$$

where  $\alpha_0$  (respectively  $\|a\|$ ) is the coercivity (respectively the continuity) constant of  $a(\varepsilon; \cdot, \cdot)$ .



Let  $\vec{u}, \vec{v}, \vec{w} \in \mathbf{X}$ . From the continuous Sobolev's embedding  $H^1(\Omega) \subset L^4(\Omega)$  together with Hölder inequality, we get

$$|c(\varepsilon; \vec{u}, \vec{v}, \vec{w})| \leq C \left(1 + \|\beta(\varepsilon)\|_{L^\infty(\Omega)}\right) \|\vec{u}\|_{\mathbf{X}} \|\vec{v}\|_{\mathbf{X}} \|\vec{w}\|_{\mathbf{X}} = C_{\text{NL}} \|\vec{u}\|_{\mathbf{X}} \|\vec{v}\|_{\mathbf{X}} \|\vec{w}\|_{\mathbf{X}}, \quad (10)$$

where  $C$  is a positive constant that depends only on  $\Omega$ . The following result gives some properties of the non-linear term (6) which are needed to prove the well-posedness of (7).

**Lemma 4.** *The nonlinear function  $G(\varepsilon; \cdot) : \mathbf{X} \rightarrow \mathbf{X}'$  defined in (6) satisfies the following estimates*

$$\begin{aligned} \|G(\varepsilon; \vec{w})\|_{\mathbf{X}'} &\leq \left\| \vec{F} \right\|_{\mathbf{X}'} + 2C_{\text{NL}} M(\varepsilon)^2 \|\vec{u}_{\text{in}}\|_{H_0^{1/2}(\Gamma_{\text{in}})^d}^2 + \|a\| M(\varepsilon) \|\vec{u}_{\text{in}}\|_{H_0^{1/2}(\Gamma_{\text{in}})^d} \\ &\quad + 2C_{\text{NL}} \|\vec{w}\|_{\mathbf{X}}^2, \end{aligned}$$

$$\|G(\varepsilon; \vec{w}_1) - G(\varepsilon; \vec{w}_2)\|_{\mathbf{X}'} \leq C_L \left( \|\vec{w}_1\|_{\mathbf{X}} + \|\vec{w}_2\|_{\mathbf{X}} + M(\varepsilon) \|\vec{u}_{\text{in}}\|_{H_0^{1/2}(\Gamma_{\text{in}})^d} \right) \|\vec{w}_1 - \vec{w}_2\|_{\mathbf{X}},$$

where  $M(\varepsilon) = C \left\{ \varepsilon_0^{-1} + \varepsilon_0^{-1} \|\nabla \varepsilon\|_{L^3(\Omega)} \right\}$  and  $C_L = C(\Omega) \max \left\{ 1, \|\beta(\varepsilon)\|_{L^\infty(\Omega)} \right\}$  with some generic constant  $C(\Omega) > 0$ .

PROOF. Using (10) with Lemma 16, we get

$$\begin{aligned} |\langle G(\varepsilon; \vec{w}), \vec{v} \rangle_{\mathbf{X}' \times \mathbf{X}}| &\leq \left\| \vec{F} \right\|_{\mathbf{X}'} \|\vec{v}\|_{\mathbf{X}} + \|a\| \left\| \vec{V} \right\|_{\mathbf{X}} \|\vec{v}\|_{\mathbf{X}} + C_{\text{NL}} \|\vec{v}\|_{\mathbf{X}} \left\| \vec{w} + \vec{V} \right\|_{\mathbf{X}}^2 \\ &\leq \|\vec{v}\|_{\mathbf{X}} \left( \left\| \vec{F} \right\|_{\mathbf{X}'} + 2C_{\text{NL}} M(\varepsilon)^2 \|\vec{u}_{\text{in}}\|_{H_0^{1/2}(\Gamma_{\text{in}})^d}^2 \right) \\ &\quad + \|\vec{v}\|_{\mathbf{X}} \left( \|a\| M(\varepsilon) \|\vec{u}_{\text{in}}\|_{H_0^{1/2}(\Gamma_{\text{in}})^d} + 2C_{\text{NL}} \|\vec{w}\|_{\mathbf{X}}^2 \right). \end{aligned}$$

Taking the supremum over  $\vec{v} \in \mathbf{X}$  with  $\|\vec{v}\|_{\mathbf{X}} \leq 1$  then gives the first estimate. For the second estimate, we start by noting that

$$\begin{aligned} \langle G(\varepsilon; \vec{w}_1) - G(\varepsilon; \vec{w}_2), \vec{v} \rangle_{\mathbf{X}' \times \mathbf{X}} &= c(\varepsilon; \vec{w}_2 + \vec{V}, \vec{w}_2 + \vec{V}, \vec{v}) - c(\varepsilon; \vec{w}_1 + \vec{V}, \vec{w}_1 + \vec{V}, \vec{v}) \\ &= \int_{\Omega} \varepsilon \left( ((\vec{w}_2 + \vec{V}) \cdot \nabla)(\vec{w}_2 + \vec{V}) - ((\vec{w}_1 + \vec{V}) \cdot \nabla)(\vec{w}_1 + \vec{V}) \right) \cdot \vec{v} \, dx \\ &\quad + \int_{\Omega} \beta(\varepsilon) \left( \left| \vec{w}_2 + \vec{V} \right| (\vec{w}_2 + \vec{V}) - \left| \vec{w}_1 + \vec{V} \right| (\vec{w}_1 + \vec{V}) \right) \cdot \vec{v} \, dx. \end{aligned}$$

For two vector fields  $\vec{a}, \vec{b}$  we have the following bounds

$$\begin{aligned} \left| \vec{a} |\vec{a}| - \vec{b} |\vec{b}| \right| &= \left| (\vec{a} - \vec{b}) |\vec{a}| - \vec{b} (|\vec{b}| - |\vec{a}|) \right| \leq |\vec{a} - \vec{b}| (|\vec{a}| + |\vec{b}|), \\ \left| \vec{a} \cdot \nabla \vec{a} - \vec{b} \cdot \nabla \vec{b} \right| &\leq |\vec{a} - \vec{b}| |\nabla \vec{a}| + |\vec{b}| |\nabla (\vec{a} - \vec{b})|. \end{aligned}$$

Gathering the previous estimates, there is a constant  $C(\Omega) > 0$  such that

$$\begin{aligned} |\langle G(\varepsilon; \vec{w}_1) - G(\varepsilon; \vec{w}_2), \vec{v} \rangle_{\mathbf{X}' \times \mathbf{X}}| &\leq C(\Omega) \max \left\{ 1, \|\beta(\varepsilon)\|_{L^\infty(\Omega)} \right\} \\ &\times \left( \|\vec{w}_1\|_{\mathbf{X}} + \|\vec{w}_2\|_{\mathbf{X}} + M(\varepsilon) \|\vec{u}_{\text{in}}\|_{H_{00}^{1/2}(\Gamma_{\text{in}})^d} \right) \\ &\times \|\vec{w}_1 - \vec{w}_2\|_{\mathbf{X}}, \end{aligned}$$

and the proof is thus finished.

We are now in position to prove the existence and uniqueness of solution to Problem (5).

**Theorem 5.** *Assume that (3) holds. Then there exists  $\eta > 0$  and  $R > 0$  (see (12) and (13)) such that if*

$$\left\| \vec{F} \right\|_{\mathbf{X}'} + \|\vec{u}_{\text{in}}\|_{H_{00}^{1/2}(\Gamma_{\text{in}})^d} \leq \eta,$$

there exists a unique  $(\vec{w}, p) \in \mathbf{X} \times L^2(\Omega)$  satisfying the weak formulation (5) and the estimate

$$\|\vec{w}\|_{\mathbf{X}} + \|p\|_{L^2(\Omega)} \leq R.$$

PROOF. Let  $\mathcal{S} : \vec{F} \in \mathbf{X}' \mapsto (\vec{w}, p) \in \mathbf{X} \times L^2(\Omega)$  where  $(\vec{w}, p) \in \mathbf{X} \times L^2(\Omega)$  is the unique solution to the linear Problem (7). The non-linear variational problem (5) is then equivalent to the following fixed point equation

$$(\vec{w}, p) = \mathcal{T}(\vec{w}, p), \tag{11}$$

where

$$\mathcal{T} = \mathcal{S}\mathcal{G} \text{ with } \mathcal{G}(\vec{w}, p) = G(\varepsilon; \vec{w}).$$

Let  $\mathbf{B}_R = \left\{ (\vec{v}, p) \in \mathbf{X} \times L^2(\Omega) \mid \|\vec{v}\|_{\mathbf{X}} + \|p\|_{L^2(\Omega)} \leq R \right\}$ . Note that  $\mathcal{T} : \mathbf{X} \times L^2(\Omega) \mapsto \mathbf{X} \times L^2(\Omega)$  and we thus only need to show that  $\mathcal{T} : \mathbf{B}_R \rightarrow \mathbf{B}_R$  for some  $R > 0$  and that  $\mathcal{T}$  is a contraction mapping.

Thanks to Theorem 3, the operator  $\mathcal{S}$  satisfy the next bound

$$\left\| \mathcal{S}\vec{F} \right\|_{\mathbf{X} \times L^2(\Omega)} \leq C_S \left\| \vec{F} \right\|_{\mathbf{X}'}, \quad C_S = \frac{1}{\alpha_0} + \frac{1}{\gamma(\varepsilon)} \left( 1 + \frac{\|a\|}{\alpha_0} \right).$$

Using now Lemma 4, we obtain

$$\|\mathcal{T}\|_{\mathbf{X} \times L^2(\Omega)} \leq C_S \|G(\varepsilon; \vec{w})\|_{\mathbf{X}'} \leq C_S \left( \mathcal{H} + 2C_{\text{NL}} \|\vec{w}\|_{\mathbf{X}}^2 \right)$$

where

$$\mathcal{H} := \left\| \vec{F} \right\|_{\mathbf{X}'} + 2C_{\text{NL}} M(\varepsilon)^2 \|\vec{u}_{\text{in}}\|_{H_{00}^{1/2}(\Gamma_{\text{in}})^d}^2 + \|a\| M(\varepsilon) \|\vec{u}_{\text{in}}\|_{H_{00}^{1/2}(\Gamma_{\text{in}})^d}.$$

If we now assume that  $\|\bar{F}\|_{\mathbf{X}'}$ ,  $\|\bar{u}_{\text{in}}\|_{H_{00}^{1/2}(\Gamma_{\text{in}})^d}$  and  $R_0$  are such that

$$\mathcal{H} < \frac{1}{8C_S^2 C_{\text{NL}}}, \quad R_0 \leq \frac{1 - \sqrt{1 - 8C_S^2 C_{\text{NL}} \mathcal{H}}}{4C_{\text{NL}} C_S} \quad (12)$$

then we obtain that

$$\|\mathcal{T}\|_{\mathbf{X} \times L^2(\Omega)} \leq R_0$$

and we have thus proved that  $\mathcal{T} : \mathbf{B}_{R_0} \rightarrow \mathbf{B}_{R_0}$ .

To prove that  $\mathcal{T}$  is a contraction mapping, we use Lemma 4 with  $(\bar{w}_1, p_1)$ ,  $(\bar{w}_2, p_2) \in \mathbf{B}_{R_1}$  which yields

$$\begin{aligned} & \|\mathcal{T}(\bar{w}_1, p_1) - \mathcal{T}(\bar{w}_2, p_2)\|_{\mathbf{X} \times L^2(\Omega)} \leq C_S \|G(\varepsilon; \bar{w}_1) - G(\varepsilon; \bar{w}_2)\|_{\mathbf{X}'} \\ & \leq C_S C_L \left( \|\bar{w}_1\|_{\mathbf{X}} + \|\bar{w}_2\|_{\mathbf{X}} + M(\varepsilon) \|\bar{u}_{\text{in}}\|_{H_{00}^{1/2}(\Gamma_{\text{in}})^d} \right) \|\bar{w}_1 - \bar{w}_2\|_{\mathbf{X}} \\ & \leq C_S C_L \left( 2R_1 + M(\varepsilon) \|\bar{u}_{\text{in}}\|_{H_{00}^{1/2}(\Gamma_{\text{in}})^d} \right) \|\bar{w}_1 - \bar{w}_2\|_{\mathbf{X}}. \end{aligned}$$

If we now chose  $R_1 > 0$  and  $\bar{u}_{\text{in}} \in H_{00}^{1/2}(\Gamma_{\text{in}})^d$  that comply with

$$\|\bar{u}_{\text{in}}\|_{H_{00}^{1/2}(\Gamma_{\text{in}})^d} \leq \frac{1}{C_L C_S M(\varepsilon)}, \quad R_1 < \frac{1}{2} \left( \frac{1}{C_L C_S} - M(\varepsilon) \|\bar{u}_{\text{in}}\|_{H_{00}^{1/2}(\Gamma_{\text{in}})^d} \right), \quad (13)$$

then the application  $\mathcal{T} : \mathbf{B}_{R_1} \rightarrow \mathbf{B}_{R_1}$  is a contraction mapping.

From (12) and (13), we get that  $\mathcal{T} : \mathbf{B}_R \rightarrow \mathbf{B}_R$  and that  $\mathcal{T}$  is a contraction mapping with  $R \leq \min\{R_0, R_1\}$ . The Banach fixed point theorem then gives the existence and uniqueness of  $(\bar{w}, p) \in \mathbf{B}_R$  which satisfy (11).

We end this section by noting that Theorem 5 gives the existence and uniqueness of a solution to (4) since  $\bar{u} = \bar{w} + \bar{V}$  where  $\bar{V} \in \mathbf{X}_1$  is the divergence-free lifting defined in Lemma 16.

### 3. Finite element approximation of the Darcy-Brinkman-Forchheimer model

We consider a quasi-uniform family of triangulations (see [17, p. 76, Definition 1.140])  $\{\mathcal{T}_h\}_{h>0}$  of  $\Omega$  whose elements are triangles ( $d = 2$ ) or tetrahedrons ( $d = 3$ ) denoted by  $K$ . We emphasize that  $\bar{\Omega} = \cup_{K \in \mathcal{T}_h} K$ . The parameter  $h_K$  is the diameter of the circle or sphere inscribed in  $K$  and we set

$$h = \max_{K \in \mathcal{T}_h} h_K.$$

We consider the Taylor-Hood finite element [44] (see also [19, p. 176, Chapter II, Section 4.2]) which consists in looking for piecewise-polynomial approximations  $(\bar{w}_h, p_h) \in \mathbf{X}_h \times \mathbf{M}_h$  of  $(\bar{w}, p) \in \mathbf{X} \times L^2(\Omega)$  with

$$\begin{aligned} \mathbf{X}_h &= \{\bar{v}_h \in \mathbf{X} \mid \forall K \in \mathcal{T}_h, \bar{v}_h|_K \in \mathbb{P}_2(K)\}, \\ \mathbf{M}_h &= \{q_h \in \mathcal{C}^0(\bar{\Omega}) \mid \forall K \in \mathcal{T}_h, q_h|_K \in \mathbb{P}_1(K)\}. \end{aligned}$$

It is worth noting that the convergence results proved in this section will also hold for other finite element spaces.

We now consider some  $\varepsilon_h \in \mathbf{M}_h$  that approximates  $\varepsilon$  in the following sense

$$\begin{aligned} \forall x \in \Omega, \quad \varepsilon_0 &\leq \varepsilon_h(x) \leq 1, \\ \|\varepsilon_h - \varepsilon\|_{L^\infty(\Omega)} &\leq Ch \|\varepsilon\|_{W^{1,\infty}(\Omega)}, \\ \|\nabla \varepsilon_h - \nabla \varepsilon\|_{L^r(\Omega)} &\leq Ch^l \|\varepsilon\|_{W^{l+1,r}(\Omega)}, \quad r > d. \end{aligned} \quad (14)$$

Note that these assumptions are satisfied if  $\varepsilon \in W^{1,\infty}(\Omega) \cap W^{l+1,r}(\Omega)$  for  $l > 0$  and if one takes  $\varepsilon_h = \mathcal{I}_h \varepsilon$  where  $\mathcal{I}_h$  is the global interpolation operator (see [17, Corollary 1.109 and Corollary 1.110]).

This section is now devoted to finite element discretization of (4) which actually amounts to consider the finite element discretization of (5) since the solution of these two problems are related thanks to  $\vec{u} = \vec{w} + \vec{V}$ . As a result, if  $\vec{w}_h$  denotes the velocity associated to (5), the finite element approximation of the solution to Problem (4) is going to be  $\vec{u}_h = \vec{w}_h + \mathcal{I}_{\mathbf{X}_h} \vec{V}$  where  $\mathcal{I}_{\mathbf{X}_h} : \mathbf{X} \mapsto \mathbf{X}_h$  is the finite element interpolate operator. The discrete problem associated to (5) reads

$$\begin{aligned} &\text{Find } (\vec{w}_h, p_h) \in \mathbf{X}_h \times \mathbf{M}_h \text{ such that} \\ &\begin{cases} a(\varepsilon_h; \vec{w}_h, \vec{v}_h) + b(\varepsilon_h; \vec{v}_h, p_h) = \langle G(\varepsilon_h; \vec{w}_h), \vec{v}_h \rangle_{\mathbf{X}' \times \mathbf{X}}, & \forall \vec{v}_h \in \mathbf{X}_h, \\ b(\varepsilon_h; \vec{w}_h, q_h) = 0, & \forall q_h \in \mathbf{M}_h, \end{cases} \end{aligned} \quad (15)$$

where the non-linear term is defined as in (6). We are going to study the existence and uniqueness of discrete solution to (15) as we did for the continuous problem, using a fixed-point approach. As a result, we study first the discretization of (7) which is

$$\begin{aligned} &\text{Find } (\vec{w}_h, p_h) \in \mathbf{X}_h \times \mathbf{M}_h \text{ such that} \\ &\begin{cases} a(\varepsilon_h; \vec{w}_h, \vec{v}_h) + b(\varepsilon_h; \vec{v}_h, p_h) = \langle \vec{F}, \vec{v}_h \rangle_{\mathbf{X}' \times \mathbf{X}}, & \forall \vec{v}_h \in \mathbf{X}_h, \\ b(\varepsilon_h; \vec{w}_h, q_h) = 0, & \forall q_h \in \mathbf{M}_h. \end{cases} \end{aligned} \quad (16)$$

Note that (16) is again a saddle-point problem with a coercive bilinear form  $a$  (see (8)) and we thus need a (discrete) inf-sup condition in order to get its well-posedness. This is done in the next subsection.

### 3.1. Discrete inf-sup conditions

This section is devoted to prove that the bilinear forms  $b(\varepsilon; \cdot, \cdot)$  and  $b(\varepsilon_h; \cdot, \cdot)$  both satisfy inf-sup conditions. The latter together with the coercivity of the bilinear forms  $a(\varepsilon_h; \cdot, \cdot)$  and  $a(\varepsilon; \cdot, \cdot)$  are necessary to prove that the linear discrete problems (16), either with  $\varepsilon$  or  $\varepsilon_h$ , are well-posed.

**Lemma 6 (Discrete inf-sup with  $\varepsilon$ ).** *Assume that  $\varepsilon$  is regular enough so that (14) holds and that at least one edge ( $d = 2$ ) or a face ( $d = 3$ ) of an*

element of  $\mathcal{T}_h$  is contained in  $\Gamma_{\text{out}}$  (see [9, Assumption 3.1]). Then the following inf-sup condition holds:

$$\inf_{q_h \in \mathbf{M}_h \setminus \{0\}} \sup_{\vec{u}_h \in \mathbf{X}_h} \frac{b(\varepsilon; \vec{u}_h, q_h)}{\|\vec{u}_h\|_{\mathbf{X}} \|q_h\|_{L^2(\Omega)}} \geq \beta(h, \varepsilon),$$

$$\beta(h, \varepsilon) = \frac{\min \left\{ \frac{1}{2}, \frac{\mu c_0 \varepsilon_0}{4|\Omega|} \right\} - C_1 C(\Omega) h M(\varepsilon) \|\varepsilon\|_{W^{1,\infty}(\Omega)}}{\max \left\{ \left( \varepsilon_0^{-1} + \varepsilon_0^{-2} \|\nabla \varepsilon\|_{L^3(\Omega)} \right), \mu C_2 \right\}},$$

where  $c_0, C_1, C_2, C(\Omega)$  are generic positive constants and only  $\mu$  defined in (20) depends on  $\varepsilon$ .

PROOF. The proof is adapted from [7, p. 18, Proposition 3.7] and [9, Lemma 3.2]. Let  $q_h \in \mathbf{M}_h$  be written as

$$q_h = \tilde{q}_h + \bar{q}, \quad \bar{q} = \frac{1}{|\Omega|} \int_{\Omega} q_h(x) dx.$$

Since  $\tilde{q}_h \in L_0^2(\Omega)$ , the continuous inf-sup condition from [19, p. 24, Corollary 2.4] gives the existence of  $\vec{v} \in H_0^1(\Omega)^d$  such that

$$\operatorname{div}(\vec{v}) = -\tilde{q}_h \quad \text{and} \quad \|\vec{v}\|_{\mathbf{X}} \leq C_1 \|\tilde{q}_h\|_{L^2(\Omega)}. \quad (17)$$

Since  $\bar{q} \in \mathbb{R}$ , we have

$$\forall \vec{v} \in \mathbf{X}, \quad b(\varepsilon; \vec{v}, \bar{q}) = -\bar{q} \int_{\Gamma_{\text{out}}} \varepsilon \vec{v} \cdot \vec{n} d\sigma.$$

Let  $\varphi \in \mathcal{C}_c^\infty(\bar{\Omega})$  be a smooth function with  $\varphi(x) > 0$  and let  $c_0 > 0$  be given by  $\int_{\Gamma_{\text{out}}} \varphi d\sigma = c_0 > 0$ . The Cauchy-Schwarz inequality together with the continuity of the trace operator then give

$$\begin{aligned} \int_{\Gamma_{\text{out}}} \varepsilon \mathcal{I}_{\mathbf{X}_h}(\varphi) d\sigma &\geq \varepsilon_0 \int_{\Gamma_{\text{out}}} \mathcal{I}_{\mathbf{X}_h}(\varphi) d\sigma \geq c_0 \varepsilon_0 - C(\Omega) \varepsilon_0 \|\mathcal{I}_{\mathbf{X}_h}(\varphi) - \varphi\|_{\mathbf{X}} \\ &\geq c_0 \varepsilon_0 - C(\Omega) h^l \|\varphi\|_{W^{l+1,2}(\Omega)} \geq \frac{c_0 \varepsilon_0}{2}, \end{aligned}$$

where we used [17, p. 61, Corollary (1.110)] and we assumed that  $h$  is small enough to get the two last lower bounds. From assumptions of Lemma 6, we now consider a regular extension  $\vec{n}_*$  of the unit normal vector  $\vec{n}|_{\Gamma_{\text{out}}}$  inside  $\Omega$  and set

$$\vec{v} = -\bar{q} \mathcal{I}_{\mathbf{X}_h}(\varphi \vec{n}_*).$$

For  $h$  small enough, the previous estimate then gives that

$$b(\varepsilon; \vec{v}, \bar{q}) \geq \frac{c_0 \varepsilon_0}{2|\Omega|} \|\bar{q}\|_{L^2(\Omega)}^2 \quad \text{and} \quad \|\vec{v}\|_{\mathbf{X}} \leq C_2 \|\bar{q}\|_{L^2(\Omega)}. \quad (18)$$

Now setting

$$\vec{u}_h = \mathcal{I}_{\mathbf{X}_h} \left( \varepsilon^{-1} \vec{v} \right) + \mu \vec{v}. \quad (19)$$

and using (17) and (18), we infer

$$\begin{aligned} b(\varepsilon; \vec{u}_h, q_h) &= b\left(\varepsilon; \varepsilon^{-1} \vec{v}, q_h\right) + b\left(\varepsilon; \mathcal{I}_{\mathbf{X}_h} \left( \varepsilon^{-1} \vec{v} \right) - \varepsilon^{-1} \vec{v}, q_h\right) + \mu b(\varepsilon; \vec{v}, q_h) \\ &\geq b\left(\varepsilon; \mathcal{I}_{\mathbf{X}_h} \left( \varepsilon^{-1} \vec{v} \right) - \varepsilon^{-1} \vec{v}, q_h\right) + \|\tilde{q}_h\|_{L^2(\Omega)}^2 + \mu \frac{c_0 \varepsilon_0}{2|\Omega|} \|\bar{q}\|_{L^2(\Omega)}^2 \\ &\quad - \mu C(\Omega) \left( C_2(1 + \|\nabla \varepsilon\|_{L^3(\Omega)}) \right) \|\bar{q}\|_{L^2(\Omega)} \|\tilde{q}_h\|_{L^2(\Omega)}. \\ &\geq b\left(\varepsilon; \mathcal{I}_{\mathbf{X}_h} \left( \varepsilon^{-1} \vec{v} \right) - \varepsilon^{-1} \vec{v}, q_h\right) + \frac{1}{2} \|\tilde{q}_h\|_{L^2(\Omega)}^2 + \frac{1}{2} \left( \mu \frac{c_0 \varepsilon_0}{|\Omega|} - \delta^2 \right) \|\bar{q}\|_{L^2(\Omega)}^2, \end{aligned}$$

where the last lower bound has been obtain thanks to Young inequality  $ab \leq a^2/(2\delta) + b^2\delta/2$  applied with

$$a = \|\tilde{q}_h\|_{L^2(\Omega)}, \quad b = \|\bar{q}\|_{L^2(\Omega)}, \quad \delta = \mu C(\Omega) \left( C_2(1 + \|\nabla \varepsilon\|_{L^3(\Omega)}) \right).$$

If we now chose  $\mu$  as

$$\mu = \frac{c_0 \varepsilon_0}{2|\Omega| \left( C(\Omega) \left( C_2(1 + \|\nabla \varepsilon\|_{L^3(\Omega)}) \right) \right)^2}, \quad (20)$$

we end up with

$$\begin{aligned} b(\varepsilon; \vec{u}_h, q_h) &\geq b\left(\varepsilon; \mathcal{I}_{\mathbf{X}_h} \left( \varepsilon^{-1} \vec{v} \right) - \varepsilon^{-1} \vec{v}, q_h\right) + \frac{1}{2} \|\tilde{q}_h\|_{L^2(\Omega)}^2 + \frac{\mu c_0 \varepsilon_0}{4|\Omega|} \|\bar{q}\|_{L^2(\Omega)}^2 \\ &\geq b\left(\varepsilon; \mathcal{I}_{\mathbf{X}_h} \left( \varepsilon^{-1} \vec{v} \right) - \varepsilon^{-1} \vec{v}, q_h\right) + \min \left\{ \frac{1}{2}, \frac{\mu c_0 \varepsilon_0}{4|\Omega|} \right\} \|q_h\|_{L^2(\Omega)}^2 \end{aligned} \quad (21)$$

Using now [7, Proof of Proposition 3.7, (ii)] and assumptions (14), we have

$$\begin{aligned} \left| b\left(\varepsilon; \mathcal{I}_{\mathbf{X}_h} \left( \varepsilon^{-1} \vec{v} \right) - \varepsilon^{-1} \vec{v}, q_h\right) \right| &\leq Ch \|\varepsilon\|_{W^{1,\infty}(\Omega)} \left\| \varepsilon^{-1} \vec{v} \right\|_{\mathbf{X}} \|q_h\|_{L^2(\Omega)} \\ &\leq Ch \|\varepsilon\|_{W^{1,\infty}(\Omega)} M(\varepsilon) \left\| \vec{v} \right\|_{\mathbf{X}} \|q_h\|_{L^2(\Omega)} \\ &\leq C_1 Ch \|\varepsilon\|_{W^{1,\infty}(\Omega)} M(\varepsilon) \|\tilde{q}_h\|_{L^2(\Omega)} \|q_h\|_{L^2(\Omega)} \\ &\leq C_1 Ch \|\varepsilon\|_{W^{1,\infty}(\Omega)} M(\varepsilon) \|q_h\|_{L^2(\Omega)}^2, \end{aligned}$$

where we used (17) for the last upper bound. From (19), (17) and (18), we get

$$\begin{aligned} \|\vec{u}_h\|_{\mathbf{X}} &\leq \left\| \mathcal{I}_{\mathbf{X}_h} \left( \varepsilon^{-1} \vec{v} \right) \right\|_{\mathbf{X}} + \mu \|\vec{v}\|_{\mathbf{X}} \leq C(\Omega) \left\| \varepsilon^{-1} \vec{v} \right\|_{\mathbf{X}} + \mu C_2 \|\bar{q}\|_{L^2(\Omega)} \\ &\leq \max \left\{ \left( \varepsilon_0^{-1} + \varepsilon_0^{-2} \|\nabla \varepsilon\|_{L^3(\Omega)} \right), \mu C_2 \right\} \|q_h\|_{L^2(\Omega)}. \end{aligned} \quad (22)$$

The desired inf-sup condition is finally proved by gathering (22), (21) and (22).

We emphasize that we are interested in solving the discrete problem (15) which involve the discretization of  $\varepsilon$ . Therefore, we extend Lemma 6 for the bilinear form  $b(\varepsilon_h; \cdot, \cdot)$ .

**Lemma 7 (Discrete inf-sup with  $\varepsilon_h$ ).** *Let the assumptions of Lemma 6 be satisfied. Then one has the following inf-sup condition*

$$\inf_{q_h \in \mathcal{M}_h \setminus \{0\}} \sup_{\vec{u}_h \in \mathbf{X}_h} \frac{b(\varepsilon_h; \vec{u}_h, q_h)}{\|\vec{u}_h\|_{\mathbf{X}} \|q_h\|_{L^2(\Omega)}} \geq \beta_2(h, \varepsilon),$$

$$\beta_2(h, \varepsilon) = \beta(h, \varepsilon) - h^l \frac{C(\Omega) \|\varepsilon\|_{W^{l+1, r}(\Omega)}}{\max \left\{ \left( \varepsilon_0^{-1} + \varepsilon_0^{-2} \|\nabla \varepsilon\|_{L^3(\Omega)} \right), \mu C_2 \right\}},$$

where  $\beta(h, \varepsilon)$  is defined in Lemma 6,  $\mu$  in (20) and  $C_2 > 0$  is a generic constant.

PROOF. We have that

$$b(\varepsilon_h; \vec{u}_h, q_h) = b(\varepsilon; \vec{u}_h, q_h) + \int_{\Omega} \operatorname{div}((\varepsilon - \varepsilon_h) \vec{u}_h) q_h \, dx.$$

From [7, p. 13; Lemma 3.1], we have the estimate

$$\forall \vec{w} \in H^1(\Omega), \quad \|(\varepsilon - \varepsilon_h) \vec{w}\|_{H^1(\Omega)} \leq Ch^l \|\varepsilon\|_{W^{l+1, r}(\Omega)} \|\vec{w}\|_{H^1(\Omega)},$$

which gives

$$b(\varepsilon_h; \vec{u}_h, q_h) \geq b(\varepsilon; \vec{u}_h, q_h) - Ch^l \|\varepsilon\|_{W^{l+1, r}(\Omega)} \|\vec{u}_h\|_{\mathbf{X}} \|q_h\|_{L^2(\Omega)}.$$

Using now  $q_h$  as in the proof of Lemma 6,  $\vec{u}_h$  given by (19) and the estimate (22), we obtain

$$b(\varepsilon_h; \vec{u}_h, q_h) \geq \left( \min \left\{ \frac{1}{2}, \frac{\mu c_0 \varepsilon_0}{4|\Omega|} \right\} - C_1 C(\Omega) h - Ch^l \|\varepsilon\|_{W^{l+1, r}(\Omega)} \right) \|q_h\|_{L^2(\Omega)}^2,$$

and the desired inf-sup condition then follows from the estimate (22) on  $\vec{u}_h$ .

We end this section by noting that both inf-sup constants satisfy  $\beta(h, \varepsilon) \geq \beta_*(\varepsilon)$  and  $\beta_2(h, \varepsilon) \geq \gamma_*(\varepsilon)$  with

$$\gamma_*(\varepsilon) = \frac{1}{8|\Omega|} \frac{\min \{2|\Omega|, \mu c_0 \varepsilon_0\}}{\max \left\{ \left( \varepsilon_0^{-1} + \varepsilon_0^{-2} \|\nabla \varepsilon\|_{L^3(\Omega)} \right), \mu C_2 \right\}},$$

if the mesh-size  $h$  is assumed to be small enough.

### 3.2. Convergence of the finite element method

We now have all the necessary tools to show that  $(\vec{w}_h, p_h)$  satisfying (15) converge toward  $(\vec{w}, p)$  which is the solution to the variational formulation (5). First we are going to prove that (15) has a unique solution under assumptions similar to those giving the well-posedness of the continuous problem (5) (see Theorem 5), namely for source terms that are small enough.

**Theorem 8.** *Assume that (3) holds. Then one has a  $\eta > 0$ , a  $R > 0$  and a  $h_{\min} > 0$  such that if*

$$\left\| \vec{F} \right\|_{\mathbf{X}'} + \|\vec{u}_{\text{in}}\|_{H_{00}^{1/2}(\Gamma_{\text{in}})^d} \leq \eta, \quad h \leq h_{\min},$$

*there exists a unique  $(\vec{w}_h, p_h) \in \mathbf{X}_h \times \mathbf{M}_h$  satisfying the weak formulation (15) and the estimate*

$$\|\vec{w}_h\|_{\mathbf{X}} + \|p_h\|_{L^2(\Omega)} \leq R.$$

PROOF. We proceed as in the proof of Theorem 5 and we thus write (15) as a fixed point equation on  $\mathbf{X}_h \times \mathbf{M}_h$ . Owing to [19, p. 59, Theorem 4.1], the coercivity of the bilinear form  $a$  (8) and Lemma 7, Problem (16) has a unique solution. We introduce the operator  $\mathcal{S}_h : \vec{F} \in \mathbf{X}' \mapsto (\vec{w}_h, p_h) \in \mathbf{X}_h \times \mathbf{M}_h$  where  $(\vec{w}_h, p_h)$  is the unique solution to (16). Assuming  $h$  is small enough so that  $\beta_2(h, \varepsilon) \geq \gamma_*(\varepsilon)$ , we also have the bound

$$\left\| \mathcal{S}_h \vec{F} \right\|_{\mathbf{X} \times L^2(\Omega)} \leq C_{\mathcal{S}_h} \left\| \vec{F} \right\|_{\mathbf{X}'}, \quad C_{\mathcal{S}_h} = \frac{1}{\alpha_0} + \frac{1}{\gamma_*(\varepsilon)} \left( 1 + \frac{\|a\|}{\alpha_0} \right).$$

The non-linear discrete problem (15) can then be written as the next fixed-point equation

$$(\vec{w}_h, p_h) = \mathcal{S}_h \mathcal{G}_h(\vec{w}_h, p_h) \quad \text{with} \quad \mathcal{G}_h(\vec{w}_h, p_h) = G(\varepsilon_h; \vec{w}_h),$$

where  $\mathcal{S}_h \mathcal{G}_h : \mathbf{X}_h \times \mathbf{M}_h \mapsto \mathbf{X}_h \times \mathbf{M}_h$ . Since the properties of  $G(\varepsilon; \cdot) : \mathbf{X} \rightarrow \mathbf{X}'$  proved in Lemma 4 are also valid for  $G(\varepsilon_h; \cdot) : \mathbf{X}_h \rightarrow \mathbf{X}'$ , the proof of the present theorem can be done exactly as the proof of Theorem 5. In addition, the smallness assumptions on the data and the definitions of  $R_0, R_1$  from (12) and (13) are the same where the only change is that  $C_{\mathcal{S}}$  has to be replaced by  $C_{\mathcal{S}_h}$ .

In the sequel, we prove optimal error estimate for the finite element approximation of the linear and non-linear problems (5) and (7), respectively.

#### *Convergence estimate for the linear problem*

We consider here the linear problem (7) whose finite element discretization is (16) when the given porosity is also discretized. We emphasize that Problem (16) falls into the class of discrete saddle-point problem such as those studied in [12, p. 65, II.2.6]. The existence and uniqueness of  $(\vec{w}_h, p_h)$  satisfying (16) is ensured by the inf-sup condition from Lemma 7 and the coercivity and continuity of the bilinear form  $a_h$  (see (8)). We also have the following convergence result.

**Theorem 9.** *Assume that  $\alpha : ]0, 1] \rightarrow \mathbb{R}^+$  is Lipschitz continuous. Let  $(\vec{w}, p)$  be the unique solution to (7). Assume that  $h$  is small enough so that the inf-sup condition from Lemma 7 holds uniformly in  $h$  and let  $(\vec{w}_h, p_h)$  be the unique solution to (16). We then have*

$$\begin{aligned} \|\vec{w}_h - \vec{w}\|_{\mathbf{X}} + \|p_h - p\|_{L^2(\Omega)} &\leq C \left( \inf_{\vec{v}_h \in \mathbf{X}_h} \|\vec{w} - \vec{v}_h\|_{\mathbf{X}} + \inf_{q_h \in \mathbf{M}_h} \|p - q_h\|_{L^2(\Omega)} \right) \\ &\quad + C \max \left\{ \|\varepsilon_h - \varepsilon\|_{L^\infty(\Omega)}, \|\varepsilon - \varepsilon_h\|_{W^{1,r}(\Omega)} \right\}. \end{aligned}$$



PROOF. We apply [12, p. 67, Proposition 2.16] to get

$$\begin{aligned} \|\vec{w}_h - \vec{w}\|_{\mathbf{X}} + \|p_h - p\|_{L^2(\Omega)} &\leq C \left( \inf_{\vec{v}_h \in \mathbf{X}_h} \|\vec{w} - \vec{v}_h\|_{\mathbf{X}} + \inf_{q_h \in \mathbf{M}_h} \|p - q_h\|_{L^2(\Omega)} \right) \\ &+ \sup_{\vec{v}_h \in \mathbf{X}_h} \frac{|a(\varepsilon; \vec{w}, \vec{v}_h) - a(\varepsilon_h; \vec{w}, \vec{v}_h)|}{\|\vec{v}_h\|_{\mathbf{X}}} + \sup_{\vec{v}_h \in \mathbf{X}_h} \frac{|b(\varepsilon; \vec{v}_h, p) - b(\varepsilon_h; \vec{v}_h, p)|}{\|\vec{v}_h\|_{\mathbf{X}}} \\ &+ \sup_{q_h \in \mathbf{M}_h} \frac{|b(\varepsilon; \vec{w}, q_h) - b(\varepsilon_h; \vec{w}, q_h)|}{\|q_h\|_{L^2(\Omega)}}. \end{aligned}$$

Note that

$$\begin{aligned} a(\varepsilon; \vec{w}, \vec{v}) - a(\varepsilon_h; \vec{w}, \vec{v}) &= \int_{\Omega} 2\text{Re}^{-1}(\varepsilon - \varepsilon_h) S(\vec{w}) : S(\vec{v}) + (\alpha(\varepsilon) - \alpha(\varepsilon_h)) \vec{w} \cdot \vec{v} \, dx, \\ b(\varepsilon; \vec{w}, q) - b(\varepsilon_h; \vec{w}, q) &= \int_{\Omega} q \text{div}((\varepsilon_h - \varepsilon)\vec{w}) \, dx. \end{aligned}$$

The error estimate then follows easily thanks to the Lipschitz continuity of  $s \in ]0, 1[ \mapsto \alpha(s) \in \mathbb{R}^+$  and the Hölder inequality.

If  $\varepsilon \in W^{1,\infty}(\Omega) \cap W^{l+1,r}(\Omega)$  for some  $l > 0$  then (14) hold. Assuming also that the solution  $(\vec{w}, p)$  to (7) are in  $H^{s+1}(\Omega)^d \times H^s(\Omega)$  then the error estimate from Theorem 9 reads

$$\begin{aligned} \|\vec{w}_h - \vec{w}\|_{\mathbf{X}} + \|p_h - p\|_{L^2(\Omega)} &\leq Ch^s \left( \|\vec{w}\|_{H^{s+1}(\Omega)} + \|p\|_{H^s(\Omega)} \right) \\ &+ C \max \left\{ h \|\varepsilon\|_{W^{1,\infty}(\Omega)}, h^l \|\varepsilon\|_{W^{l+1,r}(\Omega)} \right\}, \end{aligned}$$

where we used [17, p. 61, Corollary 1.110] to get the dependence of the inf with respect to the meshsize.

Let us now consider  $\vec{u} = \vec{w} + \vec{V}$  where  $\vec{V}$  is the divergence-free lifting of the inhomogeneous Dirichlet boundary condition introduced in Lemma 16. It is worth noting that  $(\vec{u}, p)$  satisfy the linear DBF problem with inhomogeneous Dirichlet boundary condition on  $\Gamma_{\text{in}}$ . The finite element discretization of  $\vec{u}$  is then  $\vec{u}_h = \vec{w}_h + \mathcal{I}_{\mathbf{X}_h} \vec{V}$  and one has the error estimate

$$\begin{aligned} \|\vec{u}_h - \vec{u}\|_{\mathbf{X}} + \|p_h - p\|_{L^2(\Omega)} &\leq Ch^s \left( \|\vec{w}\|_{H^{s+1}(\Omega)} + \|p\|_{H^s(\Omega)} \right) + \left\| \vec{V} - \mathcal{I}_{\mathbf{X}_h} \vec{V} \right\|_{\mathbf{X}} \\ &+ C \max \left\{ h \|\varepsilon\|_{W^{1,\infty}(\Omega)}, h^l \|\varepsilon\|_{W^{l+1,r}(\Omega)} \right\}. \end{aligned}$$

*The discrete non-linear problem without using a discrete porosity*

We consider now the discrete problem associated to (5) where the porosity is not discretized. The latter is very similar to (15) and reads

$$\begin{aligned} &\text{Find } (\vec{w}_h, p_h) \in \mathbf{X}_h \times \mathbf{M}_h \text{ such that} \\ &\begin{cases} a(\varepsilon; \vec{w}_h, \vec{v}_h) + b(\varepsilon; \vec{v}_h, p_h) = \langle G(\varepsilon; \vec{w}_h), \vec{v}_h \rangle_{\mathbf{X}' \times \mathbf{X}}, & \forall \vec{v}_h \in \mathbf{X}_h, \\ b(\varepsilon; \vec{w}_h, q_h) = 0, & \forall q_h \in \mathbf{M}_h. \end{cases} \end{aligned} \quad (23)$$

We emphasize that the existence and uniqueness of solution to (23) can be proved with arguments similar to those used to get Theorem 8. We are now going to compute the effective order of convergence of  $(\vec{w}_h, p_h)$  toward  $(\vec{w}, p)$  which satisfy (5). This can be done using the results from [10] (see also [22, p. 14, Section 4.2], [23] and [9, Theorem 4.3]) and relies on several properties that we check below.

We recall that (5) is equivalent to

$$0 = \mathcal{F}(\vec{w}, p) := (\vec{w}, p) - \mathcal{S}\mathcal{G}(\vec{w}, p) \text{ with } \mathcal{G}(\vec{w}, p) = G(\varepsilon; \vec{w}),$$

where  $\mathcal{S} : \vec{F} \in \mathbf{X}' \mapsto (\vec{w}, p) \in \mathbf{X} \times L^2(\Omega)$  is the unique solution of (7). Now, let  $\mathcal{S}_h : \vec{F} \in \mathbf{X}' \mapsto (\vec{w}_h, p_h) \in \mathbf{X}_h \times \mathbf{M}_h$  be the operator associated to any right-hand side  $\vec{F}$  the solution to the following linear discrete problem

$$\begin{aligned} & \text{Find } (\vec{w}_h, p_h) \in \mathbf{X}_h \times \mathbf{M}_h \text{ such that} \\ & \begin{cases} a(\varepsilon; \vec{w}_h, \vec{v}_h) + b(\varepsilon; \vec{v}_h, p_h) = \langle \vec{F}, \vec{v}_h \rangle_{\mathbf{X}' \times \mathbf{X}}, & \forall \vec{v}_h \in \mathbf{X}_h, \\ b(\varepsilon; \vec{w}_h, q_h) = 0, & \forall q_h \in \mathbf{M}_h. \end{cases} \end{aligned} \quad (24)$$

Then (23) is equivalent to the non-linear equation

$$0 = \mathcal{F}_h(\vec{w}_h, p_h) := (\vec{w}_h, p_h) - \mathcal{S}_h \mathcal{G}(\vec{w}_h, p_h) \text{ with } \mathcal{G}(\vec{w}_h, p_h) = G(\varepsilon; \vec{w}_h). \quad (25)$$

Since (24) is a linear saddle-point problem where the bilinear form  $a$  is coercive and continuous (see (8)) and the bilinear form  $b$  satisfy an inf-sup condition (see Lemma 6), we can apply [12, p. 54, Proposition 2.4] and get that the operator  $\mathcal{S}_h$  verifies

$$\begin{aligned} \|\mathcal{S}_h \vec{F}\|_{\mathbf{X} \times L^2(\Omega)} &\leq C \|\vec{F}\|_{\mathbf{X}'}, \\ \|(\mathcal{S}_h - \mathcal{S}) \vec{F}\|_{\mathbf{X} \times L^2(\Omega)} &\leq C \inf_{(\vec{v}_h, q_h) \in \mathbf{X}_h \times \mathbf{M}_h} \|\mathcal{S} \vec{F} - (\vec{v}_h, q_h)\|_{\mathbf{X} \times L^2(\Omega)}. \end{aligned} \quad (26)$$

We emphasize that, for  $h$  small enough to ensure that  $\beta(h, \varepsilon) \geq \gamma_*(\varepsilon)$ , the constant  $C$  in (26) does not depend on  $h$  but it may depends on  $\varepsilon$ . From (26) and the density of smooth function in  $\mathbf{X} \times L^2(\Omega)$ , we obtain

$$\lim_{h \rightarrow 0} \|(\mathcal{S}_h - \mathcal{S}) \vec{F}\|_{\mathbf{X} \times L^2(\Omega)} = 0. \quad (27)$$

We prove below that the differential  $D_{\mathcal{F}}(\vec{w}, p)$  of  $\mathcal{F}$  at  $(\vec{w}, p)$  is an isomorphism of  $\mathbf{X} \times L^2(\Omega)$ .

**Lemma 10.** *Let  $(\vec{w}, p) \in \mathbf{X} \times L^2(\Omega)$  be the solution to  $\mathcal{F}(\vec{w}, p) = 0$  satisfying  $\|\vec{w}\|_{\mathbf{X}} + \|p\|_{L^2(\Omega)} \leq R$  where  $R \leq \min\{R_0, R_1\}$  (see (12) and (13)) can be as small as we wish. Then there exists  $\eta > 0$  such that if*

$$\|\vec{w}\|_{\mathbf{X}} + \|\vec{u}_{\text{in}}\|_{H_0^{1/2}(\Gamma_{\text{in}})^d} \leq \eta,$$

*then  $D_{\mathcal{F}}(\vec{w}, p) : \mathbf{X} \times L^2(\Omega) \rightarrow \mathbf{X} \times L^2(\Omega)$  is an isomorphism with bounded inverse.*

PROOF. A computation gives

$$D_{\mathcal{F}}(\vec{w}, p)[\delta\vec{w}, \delta p] = (\delta\vec{w}, \delta p) - \mathcal{S}D_{\mathcal{G}}(\vec{w}, p)[\delta\vec{w}, \delta p].$$

We recall that  $\mathcal{G}(\vec{w}, p) = G(\varepsilon; \vec{w})$  where the non-linear term  $G(\varepsilon; \cdot)$  is defined in (6). Then

$$\begin{aligned} \langle D_{\mathcal{G}}(\vec{w}, p)[\delta\vec{w}, \delta p], \vec{v} \rangle_{\mathbf{X}' \times \mathbf{X}} &= - \int_{\Omega} \varepsilon \left( (\vec{w} + \vec{V}) \cdot \nabla \right) \delta\vec{w} \cdot \vec{v} \, dx \\ &\quad - \int_{\Omega} \varepsilon \left( (\delta\vec{w}) \cdot \nabla \right) (\vec{w} + \vec{V}) \cdot \vec{v} \, dx \\ &\quad - \int_{\Omega} \beta(\varepsilon) \left\{ |\vec{w} + \vec{V}| \delta\vec{w} \cdot \vec{v} \right\} \, dx \\ &\quad - \int_{\Omega} \beta(\varepsilon) \left\{ \frac{\vec{w} + \vec{V}}{|\vec{w} + \vec{V}|} \cdot \delta\vec{w} \left( (\vec{w} + \vec{V}) \cdot \vec{v} \right) \right\} \, dx. \end{aligned} \quad (28)$$

To study the inverse of the operator  $[\delta\vec{w}, \delta p] \rightarrow D_{\mathcal{F}}(\vec{w}, p)[\delta\vec{w}, \delta p]$ , we consider the equation  $D_{\mathcal{F}}(\vec{w}, p)[\delta\vec{w}, \delta p] = \vec{F}$  which is equivalent to the linear saddle-point problem

$$\begin{cases} \text{Find } (\delta\vec{w}, \delta p) \in \mathbf{X} \times L^2(\Omega) \text{ such that } \forall \vec{v} \in \mathbf{X}, q \in L^2(\Omega) : \\ a(\varepsilon; \delta\vec{w}, \vec{v}) - \langle D_{\mathcal{G}}(\vec{w}, p)[\delta\vec{w}, \delta p], \vec{v} \rangle_{\mathbf{X}' \times \mathbf{X}} + b(\varepsilon; \vec{v}, \delta p) = \langle \vec{F}, \vec{v} \rangle_{\mathbf{X}' \times \mathbf{X}}, \\ b(\varepsilon; \delta\vec{w}, q) = 0. \end{cases} \quad (29)$$

Theorem 5 and Lemma 16 give that  $(\vec{w}, p) \in \mathbf{X} \times L^2(\Omega)$  and  $\vec{V}$  satisfy the next estimate

$$\|\vec{w}\|_{\mathbf{X}} + \|p\|_{L^2(\Omega)} \leq R, \quad \|\vec{V}\|_{\mathbf{X}} \leq M(\varepsilon) \|u_{\text{in}}^{\vec{w}}\|_{H_{00}^{1/2}(\Gamma_{\text{in}})^d}.$$

with  $R \leq \min\{R_0, R_1\}$  where  $R_0$  and  $R_1$  are respectively defined in (12) and (13). As a result, there exists a constant  $C > 0$  such that

$$|\langle D_{\mathcal{G}}(\vec{w}, p)[\delta\vec{w}, \delta p], \delta\vec{w} \rangle_{\mathbf{X}' \times \mathbf{X}}| \leq C \left( \|\vec{w}\|_{\mathbf{X}} + \|u_{\text{in}}^{\vec{w}}\|_{H_{00}^{1/2}(\Gamma_{\text{in}})^d} \right) \|\delta\vec{w}\|_{\mathbf{X}}^2.$$

Since  $R$  can be as small as we want, there exists  $\eta > 0$  such that if  $\|\vec{w}\|_{\mathbf{X}} + \|u_{\text{in}}^{\vec{w}}\|_{H_{00}^{1/2}(\Gamma_{\text{in}})^d} \leq \eta$ , then

$$|\langle D_{\mathcal{G}}(\vec{w}, p)[\delta\vec{w}, \delta p], \delta\vec{w} \rangle_{\mathbf{X}' \times \mathbf{X}}| \leq \text{Re}^{-1} \varepsilon_0 \|\delta\vec{w}\|_{\mathbf{X}}^2.$$

Using now (8), we obtain that the bilinear form

$$A(\delta\vec{w}, \vec{v}) := a(\varepsilon; \delta\vec{w}, \vec{v}) - \langle D_{\mathcal{G}}(\vec{w}, p)[\delta\vec{w}, \delta p], \vec{v} \rangle_{\mathbf{X}' \times \mathbf{X}},$$

is coercive and continuous on  $\mathbf{X} \times \mathbf{X}$ . Lemma 2 gives that  $b(\varepsilon; \cdot, \cdot)$  satisfies an inf-sup condition and [12, II.1, Proposition 1.3] then show that (29) is well-posed and the solutions satisfy a bound similar to those of Theorem 3. This proves that  $D_{\mathcal{F}}(\vec{w}, p) : \mathbf{X} \times L^2(\Omega) \rightarrow \mathbf{X} \times L^2(\Omega)$  is an isomorphism with bounded inverse.

We now show the properties needed to apply the results from [10].

**Theorem 11.** *Assume that (3) holds and the solution  $(\vec{w}, p)$  to (5) is in  $H^{s+1}(\Omega)^d \times H^s(\Omega)$ . Assume also that  $h$  is small enough so that  $\mathcal{S}_h$  is well-defined. Then we have the following properties*

(i) *The next error estimate is valid*

$$\left\| (\mathcal{S}_h - \mathcal{S}) \vec{F} \right\|_{\mathbf{X} \times L^2(\Omega)} \leq Ch^s \left\| \mathcal{S} \vec{F} \right\|_{H^{s+1}(\Omega)^d \times H^s(\Omega)}.$$

(ii) *There exists a constant  $C(\vec{w}, p) > 0$  that does not depend on  $h$  so that*

$$\|\mathcal{F}_h(\vec{w}, p)\|_{\mathbf{X} \times L^2(\Omega)} \leq C(\vec{w}, p)h^s.$$

(iii) *There exists  $\eta > 0$  such that if  $\|\vec{w}\|_{\mathbf{X}} + \|\vec{u}_{\text{in}}\|_{H_{00}^{1/2}(\Gamma_{\text{in}})^d} \leq \eta$ , then  $D_{\mathcal{F}_h}(\vec{w}, p)$  is an isomorphism of  $\mathbf{X} \times L^2(\Omega)$  and the norm of its inverse is bounded independently of  $h$ .*

(iv) *There exists a neighborhood  $\mathcal{U}$  of  $(\vec{w}, p) \in \mathbf{X} \times L^2(\Omega)$  and a constant  $L > 0$  such that*

$$\forall (\vec{v}, q) \in \mathcal{U}, \|D_{\mathcal{F}_h}(\vec{w}, p) - D_{\mathcal{F}_h}(\vec{v}, q)\|_{\mathcal{L}(\mathbf{X} \times L^2(\Omega))} \leq L \|(\vec{w} - \vec{v}, p - q)\|_{\mathbf{X} \times L^2(\Omega)}.$$

PROOF. The proof of (i) follows from (26), the regularity of  $(\vec{w}, p)$  and [17, p. 61, Corollary 1.110]. To get (ii), we note that  $\mathcal{F}(\vec{w}, p) = 0$ . Using then (25), we obtain that

$$\mathcal{F}_h(\vec{w}, p) = \mathcal{F}_h(\vec{w}, p) - \mathcal{F}(\vec{w}, p) = (\mathcal{S}_h - \mathcal{S}) \mathcal{G}(\vec{w}, p).$$

Lemma 4 and (i) then prove (ii). Regarding (iii), the invertibility of  $D_{\mathcal{F}_h}(\vec{w}, p)$  can be obtained as in the proof of Lemma 10. The fact that the inverse of  $D_{\mathcal{F}_h}(\vec{w}, p)$  has a norm that does not depend on  $h$  follows from the fact that the coercivity constant of  $a(\varepsilon; \cdot, \cdot)$  and the inf-sup constant of  $b(\varepsilon; \cdot, \cdot)$  does not depend on the mesh-size if  $h$  is small enough.

We now prove (iv). The differential  $D_{\mathcal{F}_h}(\vec{w}, p)$  of  $\mathcal{F}_h$  at  $(\vec{w}, p)$  is given by

$$D_{\mathcal{F}_h}(\vec{w}, p)[\delta\vec{w}, \delta p] = (\delta\vec{w}, \delta p) - \mathcal{S}_h D_{\mathcal{G}}(\vec{w}, p)[\delta\vec{w}, \delta p],$$

where  $D_{\mathcal{G}}(\vec{w}, p)$  is defined in (28). From (26), we only have to study the local Lipschitz property of the application  $(\vec{v}, q) \rightarrow D_{\mathcal{G}}(\vec{v}, q)$ . Using (28), one can see that the three first terms appearing in  $D_{\mathcal{G}}(\vec{v}, q)$  are locally Lipschitz. It only remains to prove that the next application  $\Psi : (\vec{v}, q) \in \mathbf{X} \times L^2(\Omega) \mapsto \Psi(\vec{v}, q) \in \mathbf{X}'$  defined for all  $\vec{u} \in \mathbf{X}$  by

$$\langle \Psi(\vec{v}, q), \vec{u} \rangle_{\mathbf{X}' \times \mathbf{X}} = - \int_{\Omega} \beta(\varepsilon) \left\{ \frac{\vec{v} + \vec{V}}{|\vec{v} + \vec{V}|} \cdot \delta\vec{w} \right\} ((\vec{v} + \vec{V}) \cdot \vec{u}) dx,$$

is Lipschitz in a neighborhood of  $(\vec{w}, p)$  satisfying  $\mathcal{F}(\vec{w}, p) = 0$ .

We start by the case where  $(\vec{w}, p) = (-\vec{V}, p)$ . It is worth noting that

$$\|\Psi(\vec{v}, q)\|_{\mathbf{X}'} \leq \|\beta\|_{L^\infty(\Omega)} \left\| \vec{v} + \vec{V} \right\|_{\mathbf{X}} \|\delta\vec{w}\|_{\mathbf{X}},$$

and thus  $\Psi(-\vec{V}, p) = 0$ . This shows that

$$\left\| \Psi(\vec{v}, q) - \Psi(-\vec{V}, p) \right\|_{\mathbf{X}'} \leq \|\beta\|_{L^\infty(\Omega)} \left\| \vec{v} - (-\vec{V}) \right\|_{\mathbf{X}} \|\delta\vec{w}\|_{\mathbf{X}},$$

and thus the application  $(\vec{v}, q) \rightarrow \Psi(\vec{v}, q)$  is Lipschitz in a neighborhood of  $(\vec{w}, p) = (-\vec{V}, p)$ .

The application  $(\vec{v}, q) \rightarrow \Psi(\vec{v}, q)$  is smooth on  $(\mathbf{X} \setminus \{-\vec{V}\}) \times L^2(\Omega)$ . The differential of  $\Psi$  for all  $\vec{w} \neq -\vec{V}$  is:

$$\begin{aligned} \langle D_\Psi(\vec{w}, p)[\delta\vec{u}], \vec{v} \rangle_{\mathbf{X}' \times \mathbf{X}} &= - \int_{\Omega} \beta(\varepsilon) \left\{ \frac{\delta\vec{u}}{|\vec{w} + \vec{V}|} \cdot \delta\vec{w} \right\} ((\vec{w} + \vec{V}) \cdot \vec{v}) \, dx \\ &\quad - \int_{\Omega} \beta(\varepsilon) \left\{ \frac{\vec{w} + \vec{V}}{|\vec{w} + \vec{V}|} \cdot \delta\vec{w} \right\} (\delta\vec{u} \cdot \vec{v}) \, dx \\ &\quad + \int_{\Omega} \beta(\varepsilon) \left\{ (\vec{w} + \vec{V}) \cdot \delta\vec{w} \right\} ((\vec{w} + \vec{V}) \cdot \vec{v}) \left( \frac{(\vec{w} + \vec{V}) \cdot \delta\vec{u}}{|\vec{w} + \vec{V}|^3} \right) \, dx. \end{aligned}$$

This yields

$$\|D_\Psi(\vec{w}, p)[\delta\vec{u}]\|_{\mathbf{X}'} \leq 3 \|\beta\|_{L^\infty(\Omega)} \|\delta\vec{w}\|_{\mathbf{X}} \|\delta\vec{u}\|_{\mathbf{X}},$$

and a Taylor expansion finally shows that  $(\vec{v}, q) \rightarrow \Psi(\vec{v}, q)$  is also locally Lipschitz in a neighborhood of  $(\vec{w}, p)$  satisfying  $\mathcal{F}(\vec{w}, p) = 0$  if  $\vec{w} \neq -\vec{V}$ .

Thanks to Lemma 10 and Theorem 11, we can use [19, p. 302, Theorem 3.1] (see also [10], [22, p. 14, Section 4.2], [23]) to get the following error estimate.

**Theorem 12.** *Let the assumptions of Lemmas 6 and 10 and of Theorem 11 be valid. Then there exists a constant  $C(\vec{w}, p) > 0$  such that*

$$\|\vec{w}_h - \vec{w}\|_{\mathbf{X}} + \|p_h - p\|_{L^2(\Omega)} \leq C(\vec{w}, p) h^s.$$

*In addition, if  $(\vec{u}, p)$  denotes the solution to (4) then its finite element approximation  $(\vec{u}_h, p_h)$  satisfies the error estimate*

$$\|\vec{u}_h - \vec{u}\|_{\mathbf{X}} + \|p_h - p\|_{L^2(\Omega)} \leq Ch^s C(\vec{w}, p) + \left\| \vec{V} - \mathcal{I}_{\mathbf{X}_h} \vec{V} \right\|_{\mathbf{X}}.$$

Theorem 12 gives optimal error estimate. Note nevertheless that the  $O(h^s)$  can be deteriorated if the divergence-free lifting  $\vec{V}$  is not regular enough.

*The non-linear problem using a discrete porosity*

We prove here some error estimates for the finite element approximation of  $(\vec{w}, p)$  which satisfies (5). To ease the presentation, we introduce some notations. The solution to (5) is denoted by  $\Phi(\varepsilon) = (\vec{w}(\varepsilon), p(\varepsilon))$ ,  $\Phi_h(\varepsilon_h) = (\vec{w}_h(\varepsilon_h), p_h(\varepsilon_h))$  satisfy the discrete problem (15) so that  $\Phi_h(\varepsilon) = (\vec{w}_h(\varepsilon), p_h(\varepsilon))$  is the solution to the non-linear discrete problem (23). We now write

$$\Phi(\varepsilon) - \Phi_h(\varepsilon_h) = (\Phi(\varepsilon) - \Phi_h(\varepsilon)) + (\Phi_h(\varepsilon) - \Phi_h(\varepsilon_h)) = E_1 + E_2.$$

It is worth noting that Theorem 12 can be used to bound  $E_1$  and yields

$$\|E_1\|_{\mathbf{X} \times L^2(\Omega)} = \|\vec{w}_h - \vec{w}\|_{\mathbf{X}} + \|p_h - p\|_{L^2(\Omega)} \leq C(\vec{w}, p)h^s. \quad (30)$$

Regarding the second error term  $E_2$ , we recall that  $\Phi_h(\varepsilon)$  is a solution to  $\mathcal{H}(\varepsilon, \Phi_h) = 0$  where

$$\mathcal{H}(\varepsilon, \Phi_h) = \mathcal{L}(\varepsilon; \vec{w}_h, p_h) - \mathcal{G}(\varepsilon; \vec{w}_h, p_h) \text{ with } \mathcal{G}(\varepsilon; \vec{w}, p) = (G(\varepsilon; \vec{w}_h), 0).$$

Above,  $\mathcal{L}(\varepsilon; \cdot, \cdot) : (\vec{w}_h, p_h) \in \mathbf{X}_h \times \mathbf{M}_h \mapsto \mathbf{X}' \times L^2(\Omega)$  is the operator associated to (24) defined by

$$\langle \mathcal{L}(\varepsilon; \vec{w}, p), (\vec{v}, q) \rangle = (a(\varepsilon; \vec{w}, \vec{v}) + b(\varepsilon; \vec{v}, p), b(\varepsilon; \vec{w}, q)).$$

We are now in position to study the regularity of the mapping  $\varepsilon \mapsto \Phi_h(\varepsilon)$  and get a bound on  $E_2$ .

**Theorem 13.** *Let (3) and the assumptions under which Problem (23) has a unique solution be valid (see Theorem 8). Then there exists  $h_0$  such that if  $h < h_0$ , we have the next estimate*

$$\|E_2\|_{\mathbf{X} \times L^2(\Omega)} \leq C \|\varepsilon - \varepsilon_h\|_{L^\infty(\Omega) \cap W^{1,r}(\Omega)},$$

where  $C > 0$  does not depend on  $h$ .

PROOF. Let  $\mathbf{Y} = L^\infty(\Omega) \cap W^{1,r}(\Omega)$  and

$$U = \{\varepsilon \in \mathbf{Y} \mid \varepsilon_0 \leq \varepsilon(x) \leq 1, \forall x \in \Omega\}.$$

For any  $\varepsilon \in U$  and if  $\vec{F}$  and  $\vec{u}_{\text{in}}$  have small enough norms, we have the existence and uniqueness of  $\Phi_h \in \mathbf{X}_h \times \mathbf{M}_h$  satisfying  $\mathcal{H}(\varepsilon, \Phi_h) = 0$  together with the estimate

$$\|\Phi_h\|_{\mathbf{X}_h \times \mathbf{M}_h} \leq R,$$

where  $R > 0$  does not depend on  $h$  since the coercivity and inf-sup constants does not depend on the mesh-size for  $h$  small enough. We also note that  $R$  can be as small as wanted. Using similar techniques as those from the demonstration of Lemma 10, one can show that  $\delta\Phi \mapsto D_{\mathcal{H}}(\varepsilon, \Phi_h)[0, \delta\Phi]$  is an isomorphism (this actually amount to solve a discrete version of (29)). We emphasize that this application is also an isomorphism even if  $\Phi_h$  is not a solution to  $\mathcal{H}(\varepsilon, \Phi_h) = 0$ , as

soon as  $\|\Phi_h\|_{\mathbf{X} \times L^2(\Omega)}$ ,  $\|\vec{F}\|_{\mathbf{X}}$  and  $\|\vec{u}_{\text{in}}\|_{H_{00}^{1/2}(\Gamma_{\text{in}})^d}$  are small enough. In addition, since the coercivity and inf-sup constants does not depend on the mesh-size the norm of the inverse of the application  $\delta\Phi \mapsto D_{\mathcal{H}}(\varepsilon, \Phi_h)[0, \delta\Phi]$  is bounded with a constant independent of  $h$ .

We now apply the implicit function theorem around some fixed  $\varepsilon \in U$  to the application  $\mathcal{H} : \mathbf{Y} \times (\mathbf{X}_h \times \mathbf{M}_h) \rightarrow \mathbf{X}' \times L^2(\Omega)$  which is continuous Fréchet differentiable. This yields two neighborhoods  $\mathcal{O} \subset U \subset \mathbf{Y}$  of  $\varepsilon$  and  $\mathcal{V}_h \subset \mathbf{X}_h \times \mathbf{M}_h$  of  $\Phi_h(\varepsilon)$  such that the application  $\varepsilon \in \mathcal{O} \mapsto \Phi_h(\varepsilon) \in \mathcal{V}_h$  is Fréchet differentiable and that

$$\forall (\varepsilon, \Phi) \in \mathcal{O} \times \mathcal{V}_h, \mathcal{H}(\varepsilon, \Phi_h(\varepsilon)) = 0. \quad (31)$$

We also have some  $\delta > 0$  such that the ball centered at  $\Phi_h(\varepsilon)$  of radius  $\delta$  is included into  $\mathcal{V}_h$ . This yields

$$\forall \varepsilon \in \mathcal{O}, \|\Phi_h(\varepsilon)\| \leq \delta + R. \quad (32)$$

Differentiating (31) with respect to  $\varepsilon$  gives

$$D_{\mathcal{H}}(\varepsilon, \Phi_h(\varepsilon)) [0, D_{\Phi_h}(\varepsilon)[\delta\varepsilon]] = -D_{\mathcal{H}}(\varepsilon, \Phi_h(\varepsilon))[\delta\varepsilon, 0].$$

A direct calculation gives that

$$\begin{aligned} \langle D_{\mathcal{H}}(\varepsilon, \Phi)[\delta\varepsilon, 0], (\vec{v}, q) \rangle &= (2\text{Re}^{-1} \int_{\Omega} \delta\varepsilon S(\vec{u}) : S(\vec{v}) + \alpha'(\varepsilon)(\delta\varepsilon)\vec{u} \cdot \vec{v} \, dx \\ &\quad - \int_{\Omega} p \text{div}(\delta\varepsilon\vec{v}) \, dx \\ &\quad + \int_{\Omega} \delta\varepsilon(\vec{u} \cdot \nabla)\vec{v} \cdot \vec{w} + \beta'(\varepsilon)(\delta\varepsilon)|\vec{u}|\vec{v} \cdot \vec{w} \, dx \\ &\quad , \quad - \int_{\Omega} q \text{div}(\delta\varepsilon\vec{u}) \, dx), \end{aligned}$$

where  $\Phi = (\vec{u}, p)$ . From the Hölder inequality, we obtain

$$\|D_{\mathcal{H}}(\varepsilon, \Phi)[\cdot, 0]\|_{\mathcal{L}(\mathbf{Y}, \mathbf{X}' \times L^2(\Omega))} \leq C \max \left\{ \|\alpha'(\varepsilon)\|_{L^\infty(\Omega)}, \|\beta'(\varepsilon)\|_{L^\infty(\Omega)} \right\} \|\Phi\|_{\mathbf{X} \times L^2(\Omega)},$$

where  $C > 0$  only depend on  $\Omega$ . From (32), we can take  $\delta, R > 0$  small enough so that  $\delta\Phi \mapsto D_{\mathcal{H}}(\varepsilon, \Phi_h)[0, \delta\Phi]$  is an isomorphism. It is worth noting that its inverse is bounded independently of  $h$ . This yields

$$\begin{aligned} \sup_{\varepsilon \in \mathcal{O}} \|D_{\Phi_h}(\varepsilon)\|_{\mathcal{L}(\mathbf{Y}, \mathbf{X}_h \times \mathbf{M}_h)} &\leq C \sup_{\varepsilon \in \mathcal{O}} \|D_{\mathcal{H}}(\varepsilon, \Phi)[\cdot, 0]\|_{\mathcal{L}(\mathbf{Y}, \mathbf{X}' \times L^2(\Omega))} \\ &\leq C \sup_{\varepsilon \in \mathcal{O}} \left( \max \left\{ \|\alpha'(\varepsilon)\|_{L^\infty(\Omega)}, \|\beta'(\varepsilon)\|_{L^\infty(\Omega)} \right\} \right). \end{aligned}$$

Owning to (14) we can chose  $h_0$  such that  $\|\varepsilon - \varepsilon_h\|_{L^\infty(\Omega) \cap W^{1,r}(\Omega)}$  is small enough so that  $\varepsilon_h \in \mathcal{O}$  for any  $h < h_0$ . Since both functions  $s \in [\varepsilon_0, 1] \mapsto \alpha'(s) \in \mathbb{R}^+$

and  $s \in [\varepsilon_0, 1] \mapsto \beta'(s) \in \mathbb{R}^+$  are bounded, one gets

$$\begin{aligned} \|\Phi_h(\varepsilon) - \Phi_h(\varepsilon_h)\|_{\mathbf{X} \times L^2(\Omega)} &\leq \sup_{\gamma \in \mathcal{O}} \|D_{\Phi_h}(\gamma)\|_{\mathcal{L}(\mathbf{Y}, \mathbf{X}_h \times \mathbf{M}_h)} \|\varepsilon - \varepsilon_h\|_{L^\infty(\Omega) \cap W^{1,r}(\Omega)} \\ &\leq C \|\varepsilon - \varepsilon_h\|_{L^\infty(\Omega) \cap W^{1,r}(\Omega)}, \end{aligned}$$

and we have finally proved the desired result.

Assuming that  $(\vec{w}, p) \in H^{s+1}(\Omega)^d \times H^s(\Omega)$  and using Theorem (13) and (30), we have proved that there exist  $\eta > 0$ ,  $R > 0$  and  $h_0$  such that if  $h < h_0$  and

$$\|\vec{F}\|_{\mathbf{X}} + \|\vec{u}_{\text{in}}\|_{H_{00}^{1/2}(\Gamma_{\text{in}})^d} \leq \eta \text{ and } \|\vec{w}\|_{\mathbf{X}} + \|p\|_{L^2(\Omega)} \leq R,$$

then

$$\begin{aligned} \|\vec{u}_h - \vec{u}\|_{\mathbf{X}} + \|p_h - p\|_{L^2(\Omega)} &\leq C(\vec{w}, p)h^s + \left\| \vec{V} - \mathcal{I}_{\mathbf{X}_h} \vec{V} \right\| \\ &\quad + C \|\varepsilon - \varepsilon_h\|_{L^\infty(\Omega) \cap W^{1,r}(\Omega)}, \end{aligned} \quad (33)$$

where  $(\vec{u}, p)$  satisfies (4) and  $(\vec{u}_h, p_h) = (\vec{w}_h + \mathcal{I}_{\mathbf{X}_h} \vec{V}, p_h)$  satisfies (15). We have then proved optimal error estimates for the finite element approximation using a discretization of the porosity of the solution to the DBF model with mixed boundary conditions.

#### 4. Numerical analysis of the DBF model

In this section, we present some numerical results related to the DBF model. First, we present the method used to solve the non-linear discrete problem. The latter relies on a fixed-point method also known as Picard iteration and we are going to prove its convergence. We consider next a smoothly varying porosity, such as those appearing in packed beds (see e.g. [46, 1, 41]), to illustrate the convergence properties of the finite element method.

##### 4.1. Picard-like iteration for solving the non-linear discrete problem

We introduce the following finite element space

$$\mathbf{X}_{1,h} = \{\vec{v}_h \in \mathbf{X}_1 \mid \forall K \in \mathcal{T}_h, \vec{v}_h|_K \in \mathbb{P}_2(K)\}.$$

We recall that the non-linear discrete problem can be written as

$$\begin{aligned} &\text{Find } (\vec{u}_h, p_h) \in \mathbf{X}_{1,h} \times \mathbf{M}_h \text{ such that for all } (\vec{v}_h, p_h) \in \mathbf{X}_h \times \mathbf{M}_h \\ &\vec{u}_h|_{\Gamma_{\text{in}}} = \vec{u}_{\text{in}} \text{ and} \\ &\begin{cases} a(\varepsilon; \vec{u}_h, \vec{v}_h) + \delta c(\varepsilon; \vec{u}_h, \vec{u}_h, \vec{v}_h) + b(\varepsilon; \vec{v}_h, p_h) &= \left\langle \vec{F}, \vec{v}_h \right\rangle_{\mathbf{X}' \times \mathbf{X}}, \\ b(\varepsilon; \vec{u}_h, p_h) &= 0, \end{cases} \end{aligned} \quad (34)$$

where one could use either  $\varepsilon$  or the finite element interpolant of the porosity, namely taking  $\varepsilon = \varepsilon_h$  in (34) and  $\delta \in \{0, 1\}$  allows to go from the linear ( $\delta = 0$ )



to the non-linear problem ( $\delta = 1$ ). The Picard-like iteration used to solve (34) with  $\delta = 1$  is obtained by computing, for some  $n$ ,  $(\vec{u}_{h,n}, p_{h,n}) \in \mathbf{X}_{1,h} \times \mathbf{M}_h$  such that  $\vec{u}_{h,n}|_{\Gamma_{\text{in}}} = \vec{u}_{\text{in}}$  and

$$\begin{cases} \forall (\vec{v}_h, p_h) \in \mathbf{X}_h \times \mathbf{M}_h \\ a(\varepsilon; \vec{u}_{h,n}, \vec{v}_h) + c(\varepsilon; \vec{u}_{h,n-1}, \vec{u}_{h,n}, \vec{v}_h) + b(\varepsilon; \vec{v}_h, p_{h,n}) = \langle \vec{F}, \vec{v}_h \rangle_{\mathbf{X}' \times \mathbf{X}}, \\ b(\varepsilon; \vec{u}_{h,n}, q_h) = 0. \end{cases} \quad (35)$$

We now assume there is no volumic right hand side to lighten the overall expressions, that is  $\vec{F} = \vec{0}$ . We study below the convergence of the iterative method (35) in this setting.

**Theorem 14.** *We consider  $\varepsilon = \varepsilon_h$  in (35). Assume that*

$$\|\alpha(\varepsilon)\|_{L^\infty(\Omega)} = O(\text{Re}^{-1}), \quad \|\beta(\varepsilon)\|_{L^\infty(\Omega)} = O(1),$$

where the  $O(\cdot)$  are used to highlight the dependance with respect to  $\text{Re}^{-1}$ . Then there exists a generic constant  $C_{\text{CV}} > 0$  that may depend on  $\varepsilon$  such that if

$$\|\vec{u}_{\text{in}}\|_{H_0^{1/2}(\Gamma_{\text{in}})^d} \leq C_{\text{CV}} \text{Re}^{-1},$$

then the sequence  $(\vec{u}_{h,n}, p_{h,n})$  generated by (35) converges toward the solution to (34) in the strong topology of  $\mathbf{X} \times L^2(\Omega)$ .

PROOF. Note first that  $(\vec{u}_{h,1}, p_{h,1}) = (\vec{w}_{h,1} + \mathcal{I}_{\mathbf{X}_{1,h}} \vec{V}, p_{h,1})$  where  $(\vec{w}_{h,1}, p_{h,1}) \in \mathbf{X}_h \times \mathbf{M}_h$  satisfies (16) with the next right hand side

$$\langle \vec{F}, \vec{v}_h \rangle_{\mathbf{X}' \times \mathbf{X}} = -a(\varepsilon; \mathcal{I}_{\mathbf{X}_{1,h}} \vec{V}, \vec{v}_h).$$

Using [12, II.1, Proposition 1.3], we get that  $(\vec{w}_{h,1}, p_{h,1})$  exists uniquely and satisfies

$$\|\vec{w}_{h,1}\|_{\mathbf{X}} \leq \frac{\|a\|}{\alpha_0} \|\vec{V}\|_{\mathbf{X}}, \quad \|p_{h,1}\|_{L^2(\Omega)} \leq \frac{1}{\beta_2(h, \varepsilon)} \left(1 + \frac{\|a\|}{\alpha_0}\right) \|a\| \|\vec{V}\|_{\mathbf{X}},$$

from which we infer

$$\|\vec{u}_{h,1}\|_{\mathbf{X}} \leq \left(\frac{\|a\|}{\alpha_0} + 1\right) \|\vec{V}\|_{\mathbf{X}} \leq \left(\frac{\|a\|}{\alpha_0} + 1\right) M(\varepsilon) \|\vec{u}_{\text{in}}\|_{H_0^{1/2}(\Gamma_{\text{in}})^d}.$$

If we now assume that  $\|\vec{u}_{\text{in}}\|_{H_0^{1/2}(\Gamma_{\text{in}})^d}$  is small enough so that

$$\|\vec{u}_{h,n-1}\|_{\mathbf{X}} \leq \frac{\text{Re}^{-1} \varepsilon_0}{C_{\text{NL}}}, \quad (36)$$

then the bilinear form  $a(\varepsilon; \cdot, \cdot) + c(\varepsilon; \vec{u}_{h,n-1}, \cdot, \cdot)$  is coercive with coercivity constant  $\text{Re}^{-1} \varepsilon_0$ . As a result,  $(\vec{w}_{h,n}, p_{h,n}) \in \mathbf{X}_h \times \mathbf{M}_h$  is well-defined and satisfies

$$\begin{aligned} \|\vec{w}_{h,n}\|_{\mathbf{X}} &\leq \frac{\|a\| + C_{\text{NL}} \|\vec{u}_{h,n-1}\|_{\mathbf{X}}}{\text{Re}^{-1} \varepsilon_0} \|\vec{V}\|_{\mathbf{X}}, \\ \|p_{h,n}\|_{L^2(\Omega)} &\leq \frac{1}{\beta_2(h, \varepsilon)} \left(1 + \frac{\|a\| + C_{\text{NL}} \|\vec{u}_{h,n-1}\|_{\mathbf{X}}}{\text{Re}^{-1} \varepsilon_0}\right) (\|a\| + C_{\text{NL}} \|\vec{u}_{h,n-1}\|_{\mathbf{X}}) \|\vec{V}\|_{\mathbf{X}}. \end{aligned}$$

Since  $(\vec{u}_{h,n}, p_{h,n}) = (\vec{w}_{h,n} + \mathcal{I}_{\mathbf{X}_{1,h}} \vec{V}, p_{h,n})$  and  $c(\varepsilon; \vec{u}_{h,n-1}, \cdot, \cdot)$  is bilinear, we infer

$$\|\vec{u}_{h,n}\|_{\mathbf{X}} \leq \left( \frac{\|a\| + C_{\text{NL}} \|\vec{u}_{h,n-1}\|_{\mathbf{X}}}{\text{Re}^{-1} \varepsilon_0} + 1 \right) \|\vec{V}\|_{\mathbf{X}}. \quad (37)$$

Now setting  $(\vec{\Phi}_{h,n}, \pi_{h,n}) = (\vec{u}_{h,n} - \vec{u}_{h,n-1}, p_{h,n} - p_{h,n-1})$ , we get that  $(\vec{\Phi}_{h,n}, \pi_{h,n}) \in \mathbf{X}_h \times \mathbf{M}_h$  satisfies the following discrete linear saddle-point problem

$$\begin{cases} a(\varepsilon; \vec{\Phi}_{h,n}, \vec{v}_h) + c(\varepsilon; \vec{u}_{h,n-1}, \vec{\Phi}_{h,n}, \vec{v}_h) + b(\varepsilon; \vec{v}_h, \pi_{h,n}) = \langle \vec{G}_n, \vec{v}_h \rangle_{\mathbf{X}' \times \mathbf{X}}, \\ b(\varepsilon; \vec{\Phi}_{h,n}, q_h) = 0, \end{cases} \quad (38)$$

with  $\vec{G}_n \in \mathbf{X}'$  defined for all  $\vec{v} \in \mathbf{X}$  as follow

$$\langle \vec{G}_n, \vec{v} \rangle_{\mathbf{X}' \times \mathbf{X}} = c(\varepsilon; \vec{u}_{h,n-2}, \vec{u}_{h,n-1}, \vec{v}) - c(\varepsilon; \vec{u}_{h,n-1}, \vec{u}_{h,n-1}, \vec{v}).$$

From (36), we know that (38) is well-posed and that  $(\vec{\Phi}_{h,n}, \pi_{h,n})$  satisfies

$$\begin{aligned} \|\Phi_{h,n}\|_{\mathbf{X}} &\leq \frac{1}{\text{Re}^{-1} \varepsilon_0} \|\vec{G}_n\|_{\mathbf{X}'}, \\ \|\pi_{h,n}\|_{L^2(\Omega)} &\leq \frac{1}{\beta_2(h, \varepsilon)} \left( 1 + \frac{\|a\| + C_{\text{NL}} \|\vec{u}_{h,n-1}\|_{\mathbf{X}}}{\text{Re}^{-1} \varepsilon_0} \right) \|\vec{G}_n\|_{\mathbf{X}'}. \end{aligned}$$

A computation gives

$$\sup_{\|v\|_{\mathbf{X}}=1} \left| \langle \vec{G}_n, \vec{v} \rangle_{\mathbf{X}' \times \mathbf{X}} \right| \leq C(\Omega) \max \left\{ 1, \|\beta(\varepsilon)\|_{L^\infty(\Omega)} \right\} \|\Phi_{h,n-1}\|_{\mathbf{X}} \|\vec{u}_{h,n-1}\|_{\mathbf{X}},$$

where  $C(\Omega)$  is a generic constant. We can thus finally infer that

$$\begin{aligned} \|\Phi_{h,n}\|_{\mathbf{X}} &\leq C(\Omega) \frac{1}{\text{Re}^{-1} \varepsilon_0} \max \left\{ 1, \|\beta(\varepsilon)\|_{L^\infty(\Omega)} \right\} \|\vec{u}_{h,n-1}\|_{\mathbf{X}} \|\Phi_{h,n-1}\|_{\mathbf{X}}, \\ \|\pi_{h,n}\|_{L^2(\Omega)} &\leq C(\Omega) \frac{1}{\beta_2(h, \varepsilon)} \left( 1 + \frac{\|a\| + C_{\text{NL}} \|\vec{u}_{h,n-1}\|_{\mathbf{X}}}{\text{Re}^{-1} \varepsilon_0} \right) \\ &\quad \times \max \left\{ 1, \|\beta(\varepsilon)\|_{L^\infty(\Omega)} \right\} \|\Phi_{h,n-1}\|_{\mathbf{X}} \|\vec{u}_{h,n-1}\|_{\mathbf{X}}. \end{aligned} \quad (39)$$

The  $O(\cdot)$  below are used to highlight the dependance of some parameters with respect to  $\text{Re}^{-1}$ . Note that  $\|a\| = C_K^2 \|\alpha(\varepsilon)\|_{L^\infty(\Omega)} + 2\text{Re}^{-1} = O(\text{Re}^{-1})$  and  $\alpha_0 = 2\text{Re}^{-1} \varepsilon_0$  (see (8)) and that  $C_{\text{NL}} = C(\Omega) \max(1, \|\beta(\varepsilon)\|_{L^\infty(\Omega)}) = O(1)$ . As a result, one has that  $\|a\| / \alpha_0 = O(1)$  and if  $\|\vec{u}_{\text{in}}\|_{H_0^{1/2}(\Gamma_{\text{in}})} \leq C_{\text{CV}} \text{Re}^{-1}$  with

$$C_{\text{CV}} \leq \frac{\alpha_0 \varepsilon_0}{C_{\text{NL}} M(\varepsilon) (\alpha_0 + \|a\|)},$$

then (36) is satisfied by  $\vec{u}_{h,1}$ . Now, if we assume that

$$C_{\text{CV}} \leq \frac{1}{2} \frac{\alpha_0 \varepsilon_0}{C_{\text{NL}} M(\varepsilon) (\alpha_0 + \|a\|)}, \quad (40)$$

we get from (37) that  $\|u_{h,2}\|_{\mathbf{X}} \leq \text{Re}^{-1}\varepsilon_0/C_{\text{NL}}$  hence it satisfies (36) and by induction the whole sequence is determined and satisfies (36) for any  $n \in \mathbb{N}^*$ . Now combining (37) with (39) give

$$\begin{aligned} \|\Phi_{h,n}\|_{\mathbf{X}} &\leq C(\Omega) \frac{1}{\text{Re}^{-1}\varepsilon_0} \max\left\{1, \|\beta(\varepsilon)\|_{L^\infty(\Omega)}\right\} \left(\frac{\|a\| + C_{\text{NL}}\|\vec{u}_{h,n-2}\|_{\mathbf{X}}}{\text{Re}^{-1}\varepsilon_0} + 1\right) \\ &\quad \times M(\varepsilon) \|\vec{u}_{\text{in}}\|_{H_{00}^{1/2}(\Gamma_{\text{in}})^d} \|\Phi_{h,n-1}\|_{\mathbf{X}} \\ &\leq 2C_{\text{CV}}C(\Omega) \frac{M(\varepsilon)\text{Re}^{-1}}{\text{Re}^{-1}\varepsilon_0} \max\left\{1, \|\beta(\varepsilon)\|_{L^\infty(\Omega)}\right\} \left(\frac{\|a\|}{\alpha_0} + 1\right) \|\Phi_{h,n-1}\|_{\mathbf{X}}. \end{aligned}$$

As a result, if in addition to (40), we have

$$2C_{\text{CV}}C(\Omega) \frac{M(\varepsilon)}{\varepsilon_0} \max\left\{1, \|\beta(\varepsilon)\|_{L^\infty(\Omega)}\right\} \left(\frac{\|a\|}{\alpha_0} + 1\right) < 1, \quad (41)$$

then the sequence  $(\vec{u}_{h,n})_{n \in \mathbb{N}^*} \subset \mathbf{X}_1$  is a Cauchy sequence and therefore converges toward some  $\vec{u}_h \in \mathbf{X}_1$ . We actually have  $\vec{u}_h \in \mathbf{X}_{1,h}$  since for all  $n \in \mathbb{N}^*$   $(\vec{u}_{h,n})_{n \in \mathbb{N}^*} \subset \mathbf{X}_{1,h}$ .

To get the convergence for the pressures  $(p_{h,n})_{n \in \mathbb{N}^*}$ , we note that (39) and (36) give the next bound

$$\|\pi_{h,n}\|_{L^2(\Omega)} \leq C(\Omega) \frac{\text{Re}^{-1}\varepsilon_0}{C_{\text{NL}}\beta_2(h,\varepsilon)} \left(2 + \frac{\|a\|}{\text{Re}^{-1}\varepsilon_0}\right) \max\left\{1, \|\beta(\varepsilon)\|_{L^\infty(\Omega)}\right\} \|\Phi_{h,n-1}\|_{\mathbf{X}}. \quad (42)$$

Since  $(\Phi_{h,n})_{n \in \mathbb{N}^*}$  is a sequence that converges to 0 in  $\mathbf{X}$ , we infer that there exists  $n_0 \in \mathbb{N}$  such that for all  $n \geq n_0$ , one has  $\|\pi_{h,n}\|_{L^2(\Omega)} \leq 1/2$  and thus the sequence  $(p_{h,n})_{n \in \mathbb{N}^*} \subset \mathbf{M}_h$  is a Cauchy sequence. This gives the existence of some  $p_h \in \mathbf{M}_h$  such that  $(p_{h,n})_{n \in \mathbb{N}^*}$  converges toward  $p_h$ . It now only remains to pass to the limit as  $n \rightarrow +\infty$  to get that  $(\vec{u}_h, p_h) \in \mathbf{X}_{1,h} \times \mathbf{M}_h$  converges toward the solution to (34).

**Remark 15.** *If  $\varepsilon$  is used then  $\beta(h, \varepsilon)$  appears in the bounds where the pressure is involved (see (39) and (42)).*

*We emphasize that the constant  $C_{\text{CV}}$  satisfy (40,41) which reduce to*

$$C_{\text{CV}} < \frac{\alpha_0\varepsilon_0}{2(\|a\| + \alpha_0)} \min\left\{\frac{1}{C_{\text{NL}}M(\varepsilon)}, \frac{1}{C(\Omega)M(\varepsilon) \max\left\{1, \|\beta(\varepsilon)\|_{L^\infty(\Omega)}\right\}}\right\},$$

*where  $C(\Omega)$  only depends on the geometry of the domain. It is also worth noting that the proof of Theorem 14 can be used to prove the existence of solution to the continuous and discrete problems (4), (34) for  $\vec{F} = \vec{0}$  and  $\alpha(\varepsilon) = O(\text{Re}^{-1})$  as soon as  $h$  is small enough to ensure that  $\beta_2(h, \varepsilon) > 0$ . Nevertheless, we found that it was easier to rely on (25) sine it fits in the framework of [19, p. 302, Theorem 3.1] (see also [10], [22, p. 14, Section 4.2], [23]) and thus allow to prove optimal error estimates for the finite element method.*

One can also compute the speed of convergence of the fixed-point iteration (35). Let us assume that  $C_{CV}$  is given as above and that  $C_{CV} \leq q < 1$ . Since  $\|\Phi_{h,n}\|_{\mathbf{X}} \leq q \|\Phi_{h,n-1}\|_{\mathbf{X}}$ , we have

$$\|\vec{u}_{h,n} - \vec{u}\|_{\mathbf{X}} \leq \frac{q^{n-1}}{1-q} \|\vec{u}_{h,1} - \vec{u}_{h,2}\|_{\mathbf{X}},$$

where  $\vec{u}_h$  satisfy (34). In addition, we can prove by induction that  $\|\Phi_{h,n-1}\|_{\mathbf{X}} \leq q^{n-2} \|\Phi_{h,2}\|_{\mathbf{X}}$  and (42) thus gives

$$\|\pi_{h,n}\|_{L^2(\Omega)} \leq C(\Omega) \frac{\text{Re}^{-1}\varepsilon_0}{C_{\text{NL}}\beta_2(h,\varepsilon)} \left(2 + \frac{\|a\|}{\text{Re}^{-1}\varepsilon_0}\right) \max\left\{1, \|\beta(\varepsilon)\|_{L^\infty(\Omega)}\right\} q^{n-2} \|\Phi_{h,2}\|_{\mathbf{X}}.$$

Note that the fixed-point iteration defined in (35) is well-defined if  $\vec{u}_{\text{in}}$  have a small enough norm. We also emphasize that, under these assumptions, this method is globally convergent. It is worth noting that the method may not converge otherwise and that the upper bound above which the method diverges also depends on the Reynolds number. As a result, divergence may occurs if  $\text{Re}$  or  $\|\vec{u}_{\text{in}}\|_{H_{00}^{1/2}(\Gamma_{\text{in}})^d}$  are too large. Nevertheless note that, for any Reynolds number, one can find some  $\vec{u}_{\text{in}}$  for which (35) actually converges and conversely, for any  $\vec{u}_{\text{in}}$ , we have a  $\text{Re}_0$  such that for any  $\text{Re} < \text{Re}_0$  the method converges. We illustrate this behavior in our numerical experiments.

#### 4.2. Numerical experiments

For this test case, we choose the following smooth porosity

$$\varepsilon(x, y) = 0.45 \left(1 + \frac{1 - 0.45}{0.45} \exp(-(1 - y))\right),$$

and recall that this can be obtained when considering packed beds such as those studied in [46, 1]. The Darcy and Forchheimer terms are defined in [30, p. 3, Eq (8,9)] (see also [46]) and read

$$\alpha(\varepsilon) = \frac{150}{\text{Re}} \left(\frac{1 - \varepsilon}{\varepsilon}\right)^2, \quad \beta(\varepsilon) = 1.75 \left(\frac{1 - \varepsilon}{\varepsilon}\right). \quad (43)$$

It is easy to see that they satisfy all the assumptions (3). We set  $\Omega = (0, 2) \times (0, 1)$ ,  $\Gamma_{\text{in}} = \{0\} \times [0, 1]$ ,  $\Gamma_{\text{out}} = \{2\} \times [0, 1]$  and  $\Gamma_w = [0, 2] \times (\{0\} \cup \{1\})$ . For the inlet velocity, we take the parabolic profile

$$\vec{u}_{\text{in}}(y) = c_{\text{in}} y(1 - y) \vec{e}_x,$$

where  $c_{\text{in}}$  is a constant which is going to be tuned in order for  $\|\vec{u}_{\text{in}}\|_{H_{00}^{1/2}(\Gamma_{\text{in}})^2}$  to be small enough to ensure that the discrete problem has a unique solution and also that (35) converges.

All the following numerical computations are done with FreeFem ++ [24]. We use the LU solver to solve the linear problems (35) which needs every submatrices to be invertible. We thus add the term  $\gamma p$  in the incompressibility




Figure 1: Value of  $c_{in,min}$  with  $it_{max} = 10$ . Top:  $N = 40$ , Bottom:  $N = 120$ , Left row:  $tol = 1e - 5$ , Right row:  $tol = 1e - 10$ .

condition with  $\gamma = 1e - 07$ . We also used a  $\mathbb{P}_1$  finite element approximation of  $\varepsilon$ , that is  $\varepsilon_h$  defined as the finite element interpolate of the porosity. We also emphasize that all the convergence theorems proved in the previous section apply to the considered test case. Finally, the mesh is obtained thanks to the Freefem command *buildmesh* ( $a(N) + b(N) + c(N) + d(N)$ ) with  $N$  being the number of vertices on each part of the boundary denoted by  $a, b, c, d$ . As a result, the mesh-size  $h$  is

$$h = \frac{\sqrt{2}}{N},$$

and we can consider only  $N$  in our numerical simulations.

To set the constant  $c_{in}$ , we are going to compute the error between the last two iterates of (35) after  $it_{max}$  iterations have been performed. This amount to compute the following quantity

$$\text{err}(\text{Re}, it_{max}) = \max \left\{ \|p_{h,it_{max}} - p_{h,it_{max}-1}\|_{L^2(\Omega)}, \|\vec{u}_{h,it_{max}} - \vec{u}_{h,it_{max}-1}\|_{L^2(\Omega)^2} \right\}.$$

We now want to find numerically  $c_{in,min}$  such that

$$\forall c > c_{in,min}, \text{err}(\text{Re}, it_{max}) > tol,$$

for a given  $it_{max}$  and tolerance. Since the fixed point iteration (35) converges if  $\vec{u}_{in}$  has small enough norm, finding such  $c_{in,min}$  is useful to setup the parameters of our numerical experiments, namely the Reynolds number and the inlet velocity.

In Figure 1 are shown values of  $c_{in,min}$  for  $it_{max} = 10$ ,  $N = 40, 120$  and  $tol = 1e - 5, 1e - 10$  for several values of the Reynolds number. We used a

discrete porosity defined as the  $\mathbb{P}_1$  finite element interpolation of  $\varepsilon$ . For all cases considered, the value of  $c_{\text{in},\text{min}}$  behaves like  $C \times \text{Re}^{-1}$ . We emphasize that this is in agreement with Theorem 14 which shows that one needs to have  $\|\vec{u}_{\text{in}}\|_{H_{00}^{1/2}(\Gamma_{\text{in}})^2} \leq C_{\text{CV}}/\text{Re}$  for the fixed point iteration to converge. It is also worth noting that, if one keeps the same number of iterations while diminishing the tolerance, then one gets a value of  $c_{\text{in},\text{min}}$  that is greatly reduced. This is actually expected from the theoretical speed of convergence of the iteration (35) computed in Remark 15 since a smaller value for  $c_{\text{in}}$  means a smaller value for  $C_{\text{CV}}$  which is directly linked to the speed of convergence. Note also that the value of  $c_{\text{in},\text{min}}$  slightly depends on the mesh-size. This is again expected from Remark 15 since the speed of convergence of the pressure depend on the discrete inf-sup constant  $\beta_2(h, \varepsilon)$  which depends on  $h$ .

We now give some illustrations of the convergence order of the finite element approximation toward the continuous solution. We first discuss the regularity of the weak solution to the Darcy-Brinkman-Forchheimer problem. Since  $\varepsilon$  is smooth and bounded over  $\bar{\Omega}$ , any solution to (1) also satisfies a Stokes problem

$$\begin{cases} -2\text{Re}^{-1}\text{div}(S(\vec{u})) + \nabla p &= \vec{F}, & \text{in } \Omega \\ \text{div}(\vec{u}) &= -\varepsilon^{-1}\nabla\varepsilon \cdot \vec{u}, & \text{in } \Omega, \end{cases} \quad (44)$$

with

$$\vec{F} = -\varepsilon^{-1}\alpha(\varepsilon)\vec{u} - \varepsilon^{-1}\beta(\varepsilon)\vec{u}|\vec{u}| - (\vec{u} \cdot \nabla)\vec{u} - 2\text{Re}^{-1}\varepsilon^{-1}S(\vec{u})\nabla\varepsilon.$$

Regarding the boundary condition on  $\Gamma_{\text{out}}$ , one can prove as in the demonstration of Lemma 16 that  $\mu \in H_{00}^{1/2}(\Gamma_{\text{out}}) \mapsto \varepsilon\mu \in H_{00}^{1/2}(\Gamma_{\text{out}})$  is a continuous linear mapping with inverse mapping given by  $\mu \in H_{00}^{1/2}(\Gamma_{\text{out}}) \mapsto \varepsilon^{-1}\mu \in H_{00}^{1/2}(\Gamma_{\text{out}})$ . The boundary condition on  $\Gamma_{\text{out}}$  thus reduces to  $(2\text{Re}^{-1}S(\vec{u}) - p)\vec{n} = 0$  in  $H_{00}^{1/2}(\Gamma_{\text{out}})'$  which corresponds to a traction boundary condition on  $\Gamma_{\text{out}}$ . As a result, any weak solution to (1) satisfies a Stokes problem with mixed boundary conditions, a right hand side  $\vec{F}$  and inhomogeneous divergence  $\text{div}(\vec{u}) \in L^2(\Omega)$ . Since  $\vec{u}_{\text{in}} = c_{\text{in}}y(1-y)$  is smooth with  $\vec{u}_{\text{in}}|_{\partial\Gamma_{\text{in}}} = 0$ , it is actually at least in  $H_{00}^{3-1/2}(\Gamma_{\text{in}})$ . The results from [28] (see also [29]) then ensure that any weak solution  $(\vec{u}, p)$  to (1) is at least in  $H^3(\Omega)^2 \times H^2(\Omega)$ . The convergence Theorem 12 (see also (33)) then gives

$$\begin{aligned} Err_{\text{tot}} &= \|\vec{u}_h - \vec{u}\|_{\mathbf{X}} + \|p_h - p\|_{L^2(\Omega)} \\ &\leq Ch^2C(\vec{w}, p) + \left\| \vec{V} - \mathcal{I}_{\mathbf{X}_h}\vec{V} \right\|_{\mathbf{X}} + C\|\varepsilon - \varepsilon_h\|_{L^\infty(\Omega) \cap W^{1,r}(\Omega)}. \end{aligned}$$

This error estimate is optimal when no finite element approximation of the porosity is used. Since we do not have explicit solution, we note  $(\vec{u}_{\text{ex}}, p_{\text{ex}})$  the solution obtained with  $N = 200$  and we compute the error between the discrete solution for  $N \leq 100$  and  $(\vec{u}_{\text{ex}}, p_{\text{ex}})$ .

The errors are shown in Figures 2 for  $(\text{Re}, c_{\text{in}}) = (500, 0.5)$  and in Figure 3 for  $(\text{Re}, c_{\text{in}}) = (1000, 1)$ . The optimal order of convergence, namely  $Err_{\text{tot}} = O(h^2)$ ,




Figure 2:  $(Re, c_{in}) = (500, 0.5)$ : (Left)  $Err_{tot}$ , (Right)  $L^2$  error for the velocity.




Figure 3:  $(Re, c_{in}) = (1000, 1)$ : (Left)  $Err_{tot}$ , (Right)  $L^2$  error for the velocity.

is obtained. Since we used an approximate porosity  $\varepsilon_h \in \mathbf{M}_h$  the convergence order is actually expected to be smaller. Nevertheless, the smoothness of  $\varepsilon$  ensures that  $\|\varepsilon - \varepsilon_h\|_{L^\infty(\Omega)} = O(h^2)$  and that  $\|\varepsilon - \varepsilon_h\|_{W^{1,r}(\Omega)} = O(h)$ . We could therefore conclude that the theoretical error estimates involving the gradient of  $(\varepsilon_h - \varepsilon)$  is not optimal and that only the  $L^\infty$  norm of  $(\varepsilon - \varepsilon_h)$  should appear in  $Err_{tot}$ . It is worth noting that this could be achieved by considering the bilinear form  $\tilde{b}(\varepsilon; \vec{u}, q) = \int_{\Omega} \varepsilon \nabla p \cdot \vec{u} \, dx$  instead of  $b(\varepsilon; \cdot, \cdot)$  which is well-defined as  $p \in H^1(\Omega)$ . Note also that the Reynolds number does not have a significant effect on the total error. Regarding the  $L^2$  error of the velocity, one could have expected one extra order of convergence as in the case of Stokes flow (see e.g. [17, p. 185, Proposition 4.18]) or elliptic problems. Note nevertheless that the convergence order in the  $L^2$ -norm is the same as the one of the total error. Once again, we have  $\|\varepsilon - \varepsilon_h\|_{L^\infty(\Omega)} = O(h^2)$  which may cause the  $L^2$  error to be second order accurate even if  $\mathbb{P}_2$  element to approximate the velocity. This claim is confirmed by Figure 4 which shows the  $L^2$ -error for the velocity where  $\varepsilon_h \in \mathbb{P}_2$  instead of  $\varepsilon_h \in \mathbb{P}_1$  as in Figures 2, 3. Indeed, in that case, we have  $\|\varepsilon - \varepsilon_h\|_{L^\infty(\Omega)} = O(h^3)$  and one can see that the  $L^2$ -norm of the velocity now behaves like  $O(h^3)$  hence recovering the optimal convergence rate.

## 5. Conclusions and outlook

We proved the well-posedness of the DBF model with mixed boundary conditions as well as the convergence of the Taylor-Hood finite element method when using a discrete porosity. We also provided a fixed point iteration, and




Figure 4:  $L^2$  error for the velocity using  $\varepsilon_h \in \mathbb{P}_2$ . (Left):  $(Re, c_{in}) = (500, 0.5)$ , (Right):  $(Re, c_{in}) = (1000, 1)$ .

proved its convergence, to solve the discrete non-linear problem and gave some numerical experiments to illustrate the optimal error estimates obtained theoretically. We emphasize that all our results hold for small enough source terms and that we showed numerically how small these terms have to be. It is also worth noting that all our results extend without any change to the incompressible Navier-Stokes equation since the latter is obtained from the DBF model when  $\varepsilon(x) = 1$  over the domain.

Regarding the perspectives, it could be interesting to try first to extend the results of this paper to the DBF problem involving the generalized Forchheimer term, namely  $\vec{u}|\vec{u}|^s$  for  $1 \leq s \leq 2$  (see e.g. [13, 47]). Secondly, it may be very interesting to study the problem of finding some optimal porous media minimizing a given cost function. Such problem falls into the class of PDE-constrained optimization problems such as those studied in [4, 5] and involve the DBF model as constraint equation. Some theoretical questions like for instance the existence of optimal porosity or the convergence of discrete optimal porosity toward the optimal continuous ones are central when working on such problems and the results proved in this paper will be of great help to provide some answers.

## Appendix A. Divergence-free lifting

We provide here the existence of a divergence-free lifting of the inhomogeneous Dirichlet boundary condition.

**Lemma 16.** *For any  $\vec{u}_{in} \in H_{00}^{1/2}(\Gamma_{in})^d$ , there exists a vector field  $\vec{V}$  solution of*

$$\begin{cases} \operatorname{div}(\varepsilon \vec{V}) = 0 & \text{in } \Omega, \\ \vec{V} = \vec{u}_{in} & \text{on } \Gamma_{in}, \\ \vec{V} = 0 & \text{on } \Gamma_w, \end{cases}$$

which satisfies

$$\|\vec{V}\|_{\mathbf{X}} \leq M(\varepsilon) \|\vec{u}_{in}\|_{H_{00}^{1/2}(\Gamma_{in})^d}.$$

where  $M(\varepsilon) = C \left\{ \varepsilon_0^{-1} + \varepsilon_0^{-1} \|\nabla \varepsilon\|_{L^3(\Omega)} \right\}$ .



PROOF. Let  $\vec{a}$  be defined as follows

$$\vec{a} = \begin{cases} \varepsilon^{-1}\vec{u}_{\text{in}} & \text{on } \Gamma_{\text{in}}, \\ 0 & \text{on } \Gamma_{\text{w}}, \\ \vec{a}_{\text{out}} & \text{on } \Gamma_{\text{out}}, \end{cases}$$

where

$$\vec{a}_{\text{out}} := \left( - \int_{\Gamma_{\text{in}}} \varepsilon^{-1}\vec{u}_{\text{in}} \cdot \vec{n} \, d\sigma \right) \vec{u}_{\text{out}}$$

with  $\vec{u}_{\text{out}} \in H_{00}^{1/2}(\Gamma_{\text{out}})^d$  being a given function such that

$$\int_{\Gamma_{\text{out}}} \vec{u}_{\text{out}} \cdot \vec{n} \, d\sigma = 1.$$

Note that, since  $\vec{u}_{\text{in}} \in H_{00}^{1/2}(\Gamma_{\text{in}})^d$ , one has  $E_0(\vec{u}_{\text{in}}) \in H^{1/2}(\partial\Omega)^d$  and thus there exists  $\vec{\Phi}_{\text{in}} \in H^1(\Omega)^d$  such that  $\vec{\Phi}_{\text{in}}|_{\partial\Omega} = E_0(\vec{u}_{\text{in}})$ . Since  $\varepsilon^{-1}\vec{\Phi}_{\text{in}} \in H^1(\Omega)^d$ ,  $\varepsilon^{-1}\vec{\Phi}_{\text{in}} \in H^{1/2}(\partial\Omega)^d$  and then  $\varepsilon^{-1}\vec{\Phi}_{\text{in}}|_{\Gamma_{\text{in}}} = \varepsilon^{-1}\vec{u}_{\text{in}} \in H^{1/2}(\Gamma_{\text{in}})^d$ . Note that

$$\int_{\Gamma_{\text{in}}} \frac{\varepsilon(s)^{-2}|\vec{u}_{\text{in}}|^2}{\text{dist}(s, \partial\Gamma_{\text{in}})} < +\infty,$$

since  $\vec{u}_{\text{in}} \in H_{00}^{1/2}(\Gamma_{\text{in}})^d$  and  $\varepsilon \in L^\infty(\bar{\Omega})$  and we obtain  $\varepsilon^{-1}\vec{u}_{\text{in}} \in H_{00}^{1/2}(\Gamma_{\text{in}})^d$ .

Since  $\vec{a}_{\text{out}} \in H_{00}^{1/2}(\Gamma_{\text{out}})^d$ , we have some  $\vec{\Phi}_{\text{out}} \in H^1(\Omega)^d$  such that  $\vec{\Phi}_{\text{out}}|_{\partial\Omega} = E_0(\vec{u}_{\text{out}})$ . Now setting  $\vec{\Phi} = \vec{\Phi}_{\text{out}} + \vec{\Phi}_{\text{in}} \in H^1(\Omega)^d$ , we have that  $\vec{\Phi}|_{\partial\Omega} = \vec{a}$  and thus  $\vec{a} \in H^{1/2}(\partial\Omega)^d$ . Since

$$\int_{\partial\Omega} \vec{a} \cdot \vec{n} \, d\sigma = 0,$$

we can use [18, p. 176, Exercise III 3.5] ( $\Omega$  needs to be bounded and locally Lipschitz) to get the existence of  $\vec{U} \in H^1(\Omega)$  that satisfies

$$\begin{cases} \text{div}(\vec{U}) & = 0 \text{ in } \Omega, \\ \vec{U} & = \vec{a} \text{ on } \partial\Omega, \end{cases} \quad (\text{A.1})$$

together with the bound

$$\|\vec{U}\|_{H^1(\Omega)^d} \leq C \|\vec{a}\|_{H^{1/2}(\partial\Omega)^d} \leq C \|\vec{u}_{\text{in}}\|_{H_{00}^{1/2}(\Gamma_{\text{in}})^d}$$

From [7, p. 3, Lemma 2.1], the application  $\vec{u} \in H^1(\Omega)^d \mapsto \varepsilon\vec{u} \in H^1(\Omega)^d$  is an isomorphism. Therefore, there exists  $\vec{V} \in \mathbf{X}$  such that  $\vec{U} = \varepsilon\vec{V}$  and

$$\begin{cases} \text{div}(\varepsilon\vec{V}) & = 0 & \text{in } \Omega, \\ \vec{V} & = \vec{u}_{\text{in}} & \text{on } \Gamma_{\text{in}}, \\ \vec{V} & = 0 & \text{on } \Gamma_{\text{w}}, \\ \vec{V} & = \varepsilon\vec{a}_{\text{out}} & \text{on } \Gamma_{\text{out}}. \end{cases} \quad (\text{A.2})$$

A computation also gives the bound (see also [41, p. 3, Theorem 2]):

$$\begin{aligned} \|\vec{V}\|_{H^1(\Omega)^d} &= \|\varepsilon^{-1}\vec{U}\|_{H^1(\Omega)^d} \leq C \left\{ \varepsilon_0^{-1} + \varepsilon_0^{-2} \|\nabla\varepsilon\|_{L^3(\Omega)} \right\} \|\vec{U}\|_{H^1(\Omega)^d} \\ &\leq C \left\{ \varepsilon_0^{-1} + \varepsilon_0^{-1} \|\nabla\varepsilon\|_{L^3(\Omega)} \right\} \|u_{\text{in}}\|_{H_{00}^{1/2}(\Gamma_{\text{in}})^d}. \end{aligned}$$

The proof is then finished thanks to Korn inequality.

## Appendix B. Regularity of the boundary stress tensor

We give here a result about the regularity of the boundary stress tensor that justify the equivalence between (1,2) and its weak formulation (4). Such result can also be used when considering inhomogeneous traction boundary conditions such as

$$\varepsilon (2\text{Re}^{-1}S(\vec{u}) - p) \vec{n} = \vec{\varphi},$$

on some part of the boundary. It is worth noting that all the results proved in the paper apply if such boundary conditions are considered since this only amount to change the right-hand-side. The regularity of the boundary stress tensor  $\varepsilon (2\text{Re}^{-1}S(\vec{u}) - p) \vec{n}$  is given in the next result.

**Lemma 17.** *Assume that  $\varepsilon \in L^\infty(\Omega) \cap W^{1,r(d)}(\Omega)$  with  $r(2) > 2$  and  $r(3) \geq 3$  and that  $\alpha(\varepsilon), \beta(\varepsilon) \in L^\infty(\Omega)$ . Let  $\vec{f} \in L^2(\Omega)^d$  and  $(\vec{u}, p) \in H^1(\Omega)^d \times L^2(\Omega)$  satisfying (1). Then, for any  $\Gamma_c \subset \partial\Omega$ , we have*

$$\mathbb{G}\vec{n} := \varepsilon (2\text{Re}^{-1}S(\vec{u}) - p) \vec{n} \in \left( H_{00}^{1/2}(\Gamma_c)^d \right)'.$$

PROOF. We begin with the proof for  $d = 3$ . From Remark 1 and the incompressibility condition, the non-linear term can be written as

$$\text{div}(\varepsilon\vec{u} \otimes \vec{u}) = \varepsilon(\vec{u} \cdot \nabla)\vec{u}.$$

Using  $\varepsilon\nabla p = \nabla(\varepsilon p) - p\nabla\varepsilon$ , one has

$$-\text{div}(2\text{Re}^{-1}\varepsilon S(\vec{u}) - \varepsilon p\mathbb{I}) = p\nabla\varepsilon + \varepsilon(\vec{u} \cdot \nabla)\vec{u} - \alpha(\varepsilon)\vec{u} - \beta(\varepsilon)\vec{u}|\vec{u}| + \varepsilon\vec{f}.$$

For  $p = 6/5$ , the Hölder inequality gives

$$\begin{aligned} \|\text{div}(2\text{Re}^{-1}\varepsilon S(\vec{u}) - \varepsilon p\mathbb{I})\|_{L^p(\Omega)} &\leq \|\varepsilon\|_{L^\infty(\Omega)} \left( \|\vec{f}\|_{L^p(\Omega)} + \|\vec{u}\|_{L^3(\Omega)} \|\nabla\vec{u}\|_{L^2(\Omega)} \right) \\ &\quad + \|\alpha(\varepsilon)\|_{L^\infty(\Omega)} \|\vec{u}\|_{L^p(\Omega)} + \|\beta(\varepsilon)\|_{L^\infty(\Omega)} \|\vec{u}\|_{L^{2p}(\Omega)}^2 \\ &\quad + \|p\|_{L^2(\Omega)} \|\nabla\varepsilon\|_{L^3(\Omega)^d}. \end{aligned}$$

From the continuous embedding  $H^1(\Omega) \subset L^6(\Omega)$  that holds in  $\mathbb{R}^3$ , one gets

$$\text{div}(2\text{Re}^{-1}\varepsilon S(\vec{u}) - \varepsilon p\mathbb{I}) \in L^{6/5}(\Omega)^d.$$

For any  $\vec{v} \in W^{1,q}(\Omega)^d$  with  $q = 6$ , we have the Green's identity

$$\int_{\Omega} \vec{v} \cdot \operatorname{div} \mathbb{G} dx + \int_{\Omega} \mathbb{G} : \nabla \vec{v} dx = \langle \mathbb{G} \cdot \vec{n}, \vec{v} \rangle, \quad (\text{B.1})$$

where  $\langle \cdot, \cdot \rangle$  is the duality product between  $(W^{1-1/q,q}(\partial\Omega))'$  and  $W^{1-1/q,q}(\partial\Omega)$ . Note that (B.1) also holds for  $\vec{v} \in H^1(\Omega)$  since  $\mathbb{G} \in L^2(\Omega)^{d \times d}$ ,  $\vec{v} \in L^6(\Omega)$  and  $\operatorname{div} \mathbb{G} \in L^{6/5}(\Omega)^d$ .

Now let  $\mu \in H_{00}^{1/2}(\Gamma_c)^d$  then  $E_0\mu \in H^{1/2}(\partial\Omega)^d$  and, since the trace operator  $\tau : H^1(\Omega) \rightarrow H^{1/2}(\partial\Omega)$  has a continuous bounded right inverse  $\tau^{-1} : H^{1/2}(\partial\Omega) \rightarrow H^1(\Omega)$ , there exists a  $\vec{v} = \tau^{-1}(E_0\mu) \in H^1(\Omega)^d$  such that  $\vec{v}|_{\partial\Omega} = E_0\mu$  and

$$\|\vec{v}\|_{H^1(\Omega)^d} \leq C(\Omega) \|E_0\mu\|_{H^{1/2}(\partial\Omega)^d} \leq C \|\mu\|_{H_{00}^{1/2}(\Gamma_c)^d},$$

for a generic constant  $C(\Omega) > 0$ . Observe now that

$$\langle \mathbb{G}\vec{n}, \mu \rangle_{(H_{00}^{1/2}(\Gamma_c)^d)' \times H_{00}^{1/2}(\Gamma_c)^d} = \langle \mathbb{G}\vec{n}, E_0\mu \rangle = \langle \mathbb{G}\vec{n}, \vec{v} \rangle.$$

The Green's formula (B.1) together with Hölder inequality and the continuous embedding  $H^1(\Omega) \subset L^6(\Omega)$  then gives that

$$\begin{aligned} |\langle \mathbb{G}\vec{n}, \mu \rangle_{\Gamma_c}| &\leq \|\vec{v}\|_{H^1(\Omega)^d} \|\mathbb{G}\|_{L^2(\Omega)^{d \times d}} + \|\operatorname{div}(\mathbb{G})\|_{L^{6/5}(\Omega)^d} \|\vec{v}\|_{L^6(\Omega)^d} \\ &\leq C(\Omega) \|\vec{v}\|_{H^1(\Omega)^d} \left( \|\mathbb{G}\|_{L^2(\Omega)^{d \times d}} + \|\operatorname{div}(\mathbb{G})\|_{L^{6/5}(\Omega)^d} \right) \\ &\leq C(\Omega) \|\mu\|_{H_{00}^{1/2}(\Gamma_c)^d} \left( \|\mathbb{G}\|_{L^2(\Omega)^{d \times d}} + \|\operatorname{div}(\mathbb{G})\|_{L^{6/5}(\Omega)^d} \right), \end{aligned}$$

which finally shows that  $\mathbb{G} \cdot \vec{n} \in (H_{00}^{1/2}(\Gamma_c)^d)'$  by taking the supremum over all  $\mu \in H_{00}^{1/2}(\Gamma_c)^d$ .

For the case  $d = 2$ , this is very similar and only amount to take  $p = 2r/(r+2)$  and use next the Sobolev's embedding  $H^1(\Omega) \subset L^s(\Omega)$  that holds with  $1 \leq s < +\infty$  when  $d = 2$ .

## References

- [1] Abelman, S., Parsa, A. B., & Sayehvand, H. O. (2018). Nanofluid flow and heat transfer in a Brinkman porous channel with variable porosity. *Quaestiones Mathematicae*, 41(4), 449-467.
- [2] Adams, R. A., & Fournier, J. J. (2003). *Sobolev spaces* (Vol. 140). Elsevier.
- [3] Akolkar, A., & Petrasch, J. (2012). Tomography-based characterization and optimization of fluid flow through porous media. *Transport in porous media*, 1-16.
- [4] Amigo, R. C. R., Prado, D. S., Paiva, J. L., Hewson, R. W., & Silva, E. C. N. (2018). Topology optimisation of biphasic adsorbent beds for gas storage. *Structural and Multidisciplinary Optimization*, 58(6), 2431-2454.

- [5] Bastide, A., Cocquet, P. H., & Ramalingom, D. (2018). Penalization model for Navier-Stokes-Darcy equations with application to porosity-oriented topology optimization. *Mathematical Models and Methods in Applied Sciences*, 28(08), 1481-1512.
- [6] Banks, H. T., & Kunisch, K. (2012). *Estimation techniques for distributed parameter systems*. Springer Science Business Media.
- [7] Bernardi, C., Laval, F., Métivet, B., & Pernaud-Thomas, B. (1992). Finite element approximation of viscous flows with varying density. *SIAM journal on numerical analysis*, 29(5), 1203-1243.
- [8] Bernardi, C., Canuto, C., & Maday, Y. (1991). Spectral approximations of the Stokes equations with boundary conditions on the pressure. *SIAM journal on numerical analysis*, 28(2), 333-362.
- [9] Bernardi, C., Chacoón Rebollo, T., & Yakoubi, D. (2015). Finite Element Discretization of the Stokes and Navier–Stokes Equations with Boundary Conditions on the Pressure. *SIAM Journal on Numerical Analysis*, 53(3), 1256-1279.
- [10] Brezzi, F., Rappaz, J., & Raviart, P. A. (1980). Finite dimensional approximation of nonlinear problems. *Numerische Mathematik*, 36(1), 1-25.
- [11] Brezis, H. (2010). *Functional analysis, Sobolev spaces and partial differential equations*. Springer Science & Business Media.
- [12] Brezzi, F., & Fortin, M. (1991). *Mixed and hybrid finite element methods* (Vol. 15). Springer Science & Business Media.
- [13] Coulaud, O., Morel, P., & Caltagirone, J. P. (1988). Numerical modelling of nonlinear effects in laminar flow through a porous medium. *Journal of Fluid Mechanics*, 190, 393-407.
- [14] Discacciati, M., & Quarteroni, A. (2009). *Navier-Stokes/Darcy coupling: modeling, analysis, and numerical approximation*.
- [15] Ebmeyer, C., & Frehse, J. (2001). Steady Navier-Stokes equations with mixed boundary value conditions in three-dimensional Lipschitzian domains. *Mathematische Annalen*, 319(2), 349-381.
- [16] Diersch, H. J. G. (2013). *FEFLOW: finite element modeling of flow, mass and heat transport in porous and fractured media*. Springer Science & Business Media.
- [17] Ern, A., & Guermond, J. L. (2013). *Theory and practice of finite elements* (Vol. 159). Springer Science & Business Media.
- [18] Galdi, G. (2011). *An introduction to the mathematical theory of the Navier-Stokes equations: Steady-state problems*. Springer Science & Business Media.

- [19] Girault, V., & Raviart, P. A. (1979). Finite element approximation of the Navier-Stokes equations. Lecture Notes in Mathematics, Berlin Springer Verlag, 749.
- [20] Girault, V., & Rivi re, B. (2009). DG approximation of coupled Navier-Stokes and Darcy equations by Beaver-Joseph-Saffman interface condition. SIAM Journal on Numerical Analysis, 47(3), 2052-2089.
- [21] Guo, Z., & Zhao, T. S. (2005). A lattice Boltzmann model for convection heat transfer in porous media. Numerical Heat Transfer, Part B, 47(2), 157-177.
- [22] Gunzburger, M. D., Hou, L., & Svobodny, T. P. (1991). Analysis and finite element approximation of optimal control problems for the stationary Navier-Stokes equations with distributed and Neumann controls. Mathematics of Computation, 57(195), 123-151.
- [23] Gunzburger, M. D., & Hou, L. S. (1996). Finite-dimensional approximation of a class of constrained nonlinear optimal control problems. SIAM journal on control and optimization, 34(3), 1001-1043.
- [24] Hecht, F. New development in FreeFem++. J. Numer. Math. 20 (2012), no. 3-4, 251-265. 65Y15
- [25] Huang, J., Chen, L., & Rui, H. (2018). Multigrid methods for a mixed finite element method of the Darcy-Forchheimer model. Journal of scientific computing, 74(1), 396-411.
- [26] Salas, J. J., L pez, H., & Molina, B. (2013). An analysis of a mixed finite element method for a Darcy-Forchheimer model. Mathematical and Computer Modelling, 57(9-10), 2325-2338.
- [27] Kaloni, P. N., & Guo, J. (1996). Steady nonlinear double-diffusive convection in a porous medium based upon the Brinkman-Forchheimer model. Journal of Mathematical Analysis and Applications, 204(1), 138-155.
- [28] Maz'ya, V., & Rossmann, J. (2007).  $L^p$  estimates of solutions to mixed boundary value problems for the Stokes system in polyhedral domains. Mathematische Nachrichten, 280(7), 751-793.
- [29] Maz'ya, V., & Rossmann, J. (2009). Mixed boundary value problems for the stationary Navier-Stokes system in polyhedral domains. Archive for rational mechanics and analysis, 194(2), 669-712.
- [30] Nithiarasu, P., Seetharamu, K. N., & Sundararajan, T. (1997). Natural convective heat transfer in a fluid saturated variable porosity medium. International Journal of Heat and Mass Transfer, 40(16), 3955-3967.
- [31] Layton, W. J., Schieweck, F., & Yotov, I. (2002). Coupling fluid flow with porous media flow. SIAM Journal on Numerical Analysis, 40(6), 2195-2218.

- [32] Lions, J. L., & Magenes, E. (1968). *Problèmes aux limites non-homogènes et applications*, Dunod.
- [33] Cao, Y., Gunzburger, M., Hua, F., & Wang, X. (2010). Coupled Stokes-Darcy model with Beavers-Joseph interface boundary condition. *Communications in Mathematical Sciences*, 8(1), 1-25.
- [34] Kieu, T. T. (2016). Analysis of expanded mixed finite element methods for the generalized Forchheimer flows of slightly compressible fluids. *Numerical Methods for Partial Differential Equations*, 32(1), 60-85.
- [35] Li, X., & Rui, H. (2019). Superconvergence of a fully conservative finite difference method on non-uniform staggered grids for simulating wormhole propagation with the Darcy-Brinkman-Forchheimer framework. *Journal of Fluid Mechanics*, 872, 438-471.
- [36] Manzoni, A., Quarteroni, A., & Salsa, S. (2019). A saddle point approach to an optimal boundary control problem for steady Navier-Stokes equations. *Mathematics in Engineering*, 1(2), 252-280.
- [37] Nield, D. A. (1991). The limitations of the Brinkman-Forchheimer equation in modeling flow in a saturated porous medium and at an interface. *International Journal of Heat and Fluid Flow*, 12(3), 269-272.
- [38] Park, E. J. (2005). Mixed finite element methods for generalized Forchheimer flow in porous media. *Numerical Methods for Partial Differential Equations: An International Journal*, 21(2), 213-228.
- [39] Ramalingom, D., Cocquet, P. H., & Bastide, A. (2018). A new interpolation technique to deal with fluid-porous media interfaces for topology optimization of heat transfer. *Computers & Fluids*, 168, 144-158.
- [40] Rui, H., & Pan, H. (2012). A Block-Centered Finite Difference Method for the Darcy-Forchheimer Model. *SIAM Journal on Numerical Analysis*, 50(5), 2612-2631.
- [41] Skrzypacz, P., & Wei, D. (2017). Solvability of the Brinkman-Forchheimer-Darcy Equation. *Journal of Applied Mathematics*.
- [42] Skrzypacz, P. (2010). Finite element analysis for flows in chemical reactors. In: Magdeburg, Univ., Fak. für Mathematik, Diss. .
- [43] Sigmund, O., & Maute, K. (2013). Topology optimization approaches. *Structural and Multidisciplinary Optimization*, 48(6), 1031-1055.
- [44] Taylor, C., & Hood, P. (1973). A numerical solution of the Navier-Stokes equations using the finite element technique. *Computers & Fluids*, 1(1), 73-100.
- [45] Temam, R. (2001). *Navier-Stokes equations: theory and numerical analysis* (Vol. 343). American Mathematical Soc..

- [46] Vafai, K. (1984). Convective flow and heat transfer in variable-porosity media. *Journal of Fluid Mechanics*, 147, 233-259.
- [47] Varsakelis, C., & Papalexandris, M. V. (2017). On the well-posedness of the Darcy-Brinkman-Forchheimer equations for coupled porous media-clear fluid flow. *Nonlinearity*, 30(4), 1449.
- [48] Whitaker, S. (2013). *The method of volume averaging* (Vol. 13). Springer Science & Business Media.
- [49] Whitaker, S. (1996). The Forchheimer equation: a theoretical development. *Transport in Porous media*, 25(1), 27-61.