



**HAL**  
open science

## Fungal Pigments and Their Prospects in Different Industries

Ajay Lagashetti, Laurent Dufossé, Sanjay Singh, Paras Singh

► **To cite this version:**

Ajay Lagashetti, Laurent Dufossé, Sanjay Singh, Paras Singh. Fungal Pigments and Their Prospects in Different Industries. *Microorganisms*, 2019, 7 (12), 10.3390/microorganisms7120604 . hal-02562464

**HAL Id: hal-02562464**

**<https://hal.univ-reunion.fr/hal-02562464v1>**

Submitted on 13 May 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Review

# Fungal Pigments and Their Prospects in Different Industries

Ajay C. Lagashetti <sup>1</sup>, Laurent Dufossé <sup>2,\*</sup> , Sanjay K. Singh <sup>1,\*</sup> and Paras N. Singh <sup>1</sup>

<sup>1</sup> Biodiversity and Palaeobiology Group, National Fungal Culture Collection of India (NFCCI), MACS' Agharkar Research Institute, G.G. Agarkar Road, Pune 411004, India; lagashettiajay@gmail.com (A.C.L.); pnsingh@aripune.org (P.N.S.)

<sup>2</sup> Chimie et Biotechnologie des Produits Naturels & ESIROI Agroalimentaire, Université de la Réunion, 15 Avenue René Cassin, CS 92003, F-97744 Saint-Denis CEDEX, France

\* Correspondence: laurent.dufosse@univ-reunion.fr (L.D.); sksingh@aripune.org (S.K.S.); Tel.: +91-020-2532-5103 (S.K.S.)

Received: 17 October 2019; Accepted: 18 November 2019; Published: 22 November 2019



**Abstract:** The public's demand for natural, eco-friendly, and safe pigments is significantly increasing in the current era. Natural pigments, especially fungal pigments, are receiving more attention and seem to be in high demand worldwide. The immense advantages of fungal pigments over other natural or synthetic pigments have opened new avenues in the market for a wide range of applications in different industries. In addition to coloring properties, other beneficial attributes of fungal pigments, such as antimicrobial, anticancer, antioxidant, and cytotoxic activity, have expanded their use in different sectors. This review deals with the study of fungal pigments and their applications and sheds light on future prospects and challenges in the field of fungal pigments. Furthermore, the possible application of fungal pigments in the textile industry is also addressed.

**Keywords:** color; natural pigments; fungal pigments; dyeing; textile fabrics

## 1. Introduction

Color has always played an important role in the life of all organisms on Earth. Human life has become truly "colorful" due to the use of colors in all its aspects, including clothes, food, and furniture. Much archaeological evidence has shown that the use of pigments as coloring agents has been practiced since ancient times [1]. Pigments, especially synthetic ones, have occupied the entire market due to their wide range of applications in different industries since their discovery in the 19th century. Different attributes such as low production costs, ease of production, and superior coloring properties have largely contributed to the establishment of synthetic pigments in the market. However, the use of synthetic colors has been found to be detrimental to human health and the environment because of their many adverse impacts [2–7]. Many disadvantages of synthetic pigments, such as poor degradation, longer persistence, potential to cause cancers/allergies, etc., have increased the demand for natural, organic, and eco-friendly pigments in the current era.

The global response, as well as the demand for eco-friendly natural pigments, has significantly increased in recent decades due to their advantages over hazardous synthetic pigments. They are used as colorants, color intensifiers, additives, antioxidants, etc., in many industries including the textile, pharmaceutical, cosmetic, painting, food, and beverage industries [1,8]. In recent years, fungi have emerged among the prominent, eco-friendly sources of natural pigments. Easy processing, fast growth in cheap media, and weather-independent growth make them an excellent alternative to natural pigments. The present review highlights the role of fungi as small factories in pigment production and their potential application in different industries, including the textile industry.

## 2. Natural Pigments

Natural pigments are naturally derived pigments synthesized mainly by plants, animals, and microbes [5,9]. Most of the natural pigments used for different purposes since ancient times are produced from plants, such as annatto, grapes, indigo, beetroot, turmeric, madder, saffron, etc. [10,11]. However, the process of pigment production from plants may not be a good option because of various problems, such as season dependency, loss of vulnerable plant species due to their extensive use, variations in color shades and intensity, expensive production, and issues related to stability and solubility [2].

Nowadays, microorganisms, including bacteria, fungi, and algae, have been shown to be an excellent alternative source of natural pigments. For the large-scale production of pigments, microorganisms are more suitable, due to a clear understanding of their cultural techniques, processing, and ease of handling. Natural pigments from microbes, especially from bacteria and fungi, have been reported worldwide by many researchers [1,10,12–20]. Many bacterial species have been reported to possess potential for pigment production [10,21–23], but their pathogenic nature as well as associated toxicity have blocked production and commercialization. This eventually opened a new avenue for producing pigments from fungi and for their various applications.

## 3. Fungal Pigments

Fungi have been shown to be a good and readily available alternative source of natural pigments [1,20,24–26]. Fungi have immense advantages over plants such as season-independent pigment production, easy and fast growth in a cheap culture medium, production of pigments with different color shades and of more stable, soluble pigments, and easy processing [10,27]. Fungi belonging to the *Monascaceae*, *Trichocomaceae*, *Nectriaceae*, *Hypocreaceae*, *Pleosporaceae*, *Cordycipitaceae*, *Xylariaceae*, *Chaetomiaceae*, *Sordariaceae*, *Chlorociboriaceae*, *Hyaloscyphaceae*, *Hymenochaetaceae*, *Polyporaceae*, *Ophiostomataceae*, *Tremellaceae*, *Herpotrichiellaceae*, and *Tuberaceae* families have been described as potent pigment producers [8,12,20,25,26,28–45] (Table 1). These fungi are known to synthesize a variety of pigments as secondary metabolites. They are prolific producers of pigments belonging to several chemical classes, such as carotenoids, melanins, azaphilones, flavins, phenazines, quinones, monascin, violacein, indigo, etc. [16,25,26,46–49] (Table 1).

The use of *Monascus* pigments for the production of red mold rice (ang-kak) is the oldest recorded use of fungal pigments by humans. Certain species of *Monascus*, viz., *Monascus ruber* and *Monascus purpureus*, have been reported to be good potential producers of pigments worldwide. Studies have shown the potential of the red pigment produced by *M. ruber* as an important food colorant as well as food additive [50,51]. Many new pigments produced by *M. ruber*, such as *N*-glucosylrubropunctamine, *N*-glucosylmonascorubramine, monarubrin, rubropunctin, etc., have been discovered (Figure 1) [52–54]. Recently, researchers revealed the first detailed biosynthetic pathway of *Monascus* azophilone pigments (MonAzPs) in *M. ruber* M7, based on targeted gene knockouts, heterologous gene expression, as well as in vitro enzymatic and chemical reactions [55]. Along with *M. ruber*, *M. purpureus* was also reported to produce a variety of novel pigments, such as monapurone A–C, monasphilone A–B, monapilol A–D, and 9-(1-hydroxyhexyl)-3-(2-hydroxypropyl)-6a-methyl-9,9a-dihydrofuro[2,3-h] isoquinoline-6,8 (2H,6aH)-dione (Figure 1) [56–59]. Another study reports on the physicochemical (pH, light, and heat stability) properties of the red pigment of *M. purpureus* [60].

**Table 1.** Updated list of pigment-producing fungi and their respective pigments [25,61].

Fungal Species	Pigments	References
<i>Monascus species</i>		
<i>Monascus pilosus</i>	Citrinin (yellow)	[61]
<i>Monascus purpureus</i>	Monascin (yellow), monascorubrin (orange), monascorubramine (red), monapurone A–C (yellow), monasphilone A and B (yellow), ankaflavin (yellow), rubropunctamine (purple-red), rubropunctatin (orange), monopilol A–D (yellow), citrinin (yellow), 9-(1-hydroxyhexyl)-3-(2-hydroxypropyl)-6a-methyl-9,9a-dihydrofuro[2,3-h]isoquinoline-6,8(2H,6aH)-dione (red), uncharacterized (red)	[56–61]
<i>Monascus ruber</i>	Monascin (yellow), monascorubramine (red), monascorubrin (orange), ankaflavin (yellow), citrinin (yellow), rubropunctamine (purple-red), rubropunctatin (orange), <i>N</i> -glucosylrubropunctamine (red), <i>N</i> -glucosylmonascorubramine (red), monarubrin (pale yellow), rubropunctin (pale yellow)	[52,54,61]
<i>Monascus species</i>	Ankaflavin (yellow) *, monascorubramine (red) *, rubropunctatin (orange) *	[25]
<i>Fusarium species</i>		
<i>Fusarium acuminatum</i> , <i>F. avenaceum</i> , <i>F. tricinctum</i>	Antibiotic Y (yellow), aurofusarin (red)	[61]
<i>Fusarium chlamyosporum</i>	Uncharacterized (red)	[62]
<i>Fusarium culmorum</i>	Aurofusarin (red), fuscofusarin (yellow), rubrofusarin (red)	[61]
<i>Fusarium fujikuroi</i> (formerly known as <i>Fusarium moniliforme</i> / <i>Fusarium verticillioides</i> )	Bikaverin (red), norbikaverin (red), <i>O</i> -demethylanhydrofusarubin (red), 8- <i>O</i> -methoxybostrycoidin, 2-(4-((3E,5E)-14-aminotetradeca-3,5-dienyloxy)butyl)-1,2,3,4-tetrahydroisoquinolin-4-ol (ATDBTHIQN) (pink), neurosporaxanthin (orange), $\beta$ -carotene (red-orange), fusarubin (red), <i>O</i> -demethylfusarubin, <i>O</i> -methyljavanicin, <i>O</i> -methylsolaniol (orange-red)	[43,61,63–65]
<i>Fusarium graminearum</i>	Aurofusarin (red), rubrofusarin (red), 5-deoxybostrycoidin anthrone (green), 6- <i>O</i> -demethyl-5-deoxybostrycoidin anthrone (blue), purpurfusarin (purple), 6- <i>O</i> -demethyl-5-deoxybostrycoidin (yellow), 5-deoxybostrycoidin (red)	[64,66]
<i>Fusarium oxysporum</i>	2,7-dimethoxy-6-(acetoxylethyl)juglone (yellow), bikaverin (red), bostrycoidin (red), nectriafurone (yellow), norjavanicin (red), <i>O</i> -methyl-6-hydroxynorjavanicin (yellow), <i>O</i> -methylanhydrofusarubin (orange-red), <i>O</i> -methylfusarubin (red), <i>O</i> -methyljavanicin, 2-acetyl-3,8-dihydroxy-6-methoxy anthraquinone (yellow), 2-(1-hydroxyethyl)-3,8-dihydroxy-6-methoxy anthraquinone (orange), neurosporaxanthin (orange), $\beta$ -carotene (red-orange), uncharacterized naphthaquinones (purple)	[43,47,61,64,67]
<i>Fusarium poae</i> , <i>F. sambucinum</i>	Aurofusarin (red)	[61]
<i>Fusarium solani</i>	Fusarubin (red), <i>O</i> -methyl-dihydrofusarubin (red), <i>O</i> -ethylfusarubin (red), isomartincins (red)	
<i>Fusarium sporotrichioides</i>	Aurofusarin (red), $\beta$ -carotene (yellow-orange) **, lycopene (red) **	[25,61]
<i>Fusarium stilboides</i>	Antibiotic Y (yellow), aurofusarin (red), nectriafurone (yellow)	[61]
<i>Fusarium venenatum</i>	Aurofusarin (red), rubrofusarin (red)	
<i>Fusarium sp.</i>	Benzoquinone (yellow)	[68]
<i>Fusarium sp.</i> PSU-F14 and PSU-F135	Fusarnaphthoquinones B (red), fusarnaphthoquinones C (red)	[69]
<i>Fusicolla aquaeductuum</i> (Formerly Known as <i>Fusarium aquaeductuum</i> )		
<i>Fusicolla aquaeductuum</i>	Neurosporaxanthin (orange), $\beta$ -carotene (red-orange)	[43]
<i>Albonectria rigidiuscula</i> (Formerly Known as <i>Fusarium decemcellulare</i> )		
<i>Albonectria rigidiuscula</i>	Javanicin (red-orange), fusarubin (red), anhydrojavanicin, anhydrofusarubin, bostrycoidin (red), novarubin	[64]

Table 1. Cont.

Fungal Species	Pigments	References
<b>Trichoderma species</b>		
<i>Trichoderma harzianum</i>	Pachybasin (yellow), chrysophanol (orange-red), emodin (yellow), 1-hydroxy-3-methyl-anthraquinone, 1,8-dihydroxy-3-methyl-anthraquinone, T22 azaphilone	[25]
<i>Trichoderma polysporum</i>	Pachybasin (yellow), chrysophanol (orange-red), emodin (yellow)	
<i>Trichoderma viride</i>	Pachybasin (yellow), chrysophanol (orange-red), emodin (yellow), 1,3,6,8-tetrahydroxyanthraquinone, 2,4,5,7-tetrahydroxyanthraquinone	
<i>Trichoderma aureoviride</i>	Pachybasin (yellow), chrysophanol (orange-red)	
<i>Trichoderma afzharzianum</i> , <i>Trichoderma pyramidale</i> , <i>Trichoderma parareesei</i> (formerly known as <i>Trichoderma atroviride</i> ), <i>Trichoderma</i> sp. 1	Uncharacterized (yellow)	[70,71]
<i>Trichoderma parceramosum</i>	Uncharacterized (red)	[72]
<b>Cordyceps farinosa (Formerly Known as Isaria farinosa)</b>		
<i>Cordyceps farinosa</i>	Anthraquinone derivative	[73]
<b>Ophiocordyceps unilateralis (Formerly Known as Cordyceps unilateralis)</b>		
<i>Ophiocordyceps unilateralis</i>	Erythrostrominone (red), 3,5,8-TMON * (red), deoxyerythrostrominone (red), deoxyerythrostrominol (red), 4-O-methyl erythrostrominone (red), epierythrostrominol (red), naphthoquinones (deep blood red) **	[25]
<b>Beauveria species</b>		
<i>Beauveria basiana</i>	Tenellin (yellow), bassianin (yellow), pyridovericin (pale yellow), pyridomacrolidin (pale yellow), oosporein (red)	
<i>Beauveria brongniartii</i> (formerly known as <i>Beauveria tenella</i> )	Tenellin (yellow), bassianin (yellow)	[25,74]
<b>Torrubiella species</b>		
<i>Torrubiella</i> sp.	Torrubiellones A–D (yellow)	[75]
<b>Lecanicillium species</b>		
<i>Lecanicillium aphanocladii</i>	Oosporein (red)	[41]
<b>Hyperdermium species</b>		
<i>Hyperdermium bertonii</i>	Skyrin (orange-red)	[25]
<b>Daldinia species</b>		
<i>Daldinia bambusicol</i> , <i>Daldinia caldariorum</i> , <i>Daldinia childiae</i> , <i>Daldinia clavata</i> , <i>Daldinia fissa</i> , <i>Daldinia grandis</i> , <i>Daldinia lloydi</i> , <i>Daldinia loculata</i> , <i>Daldinia petriniae</i> , <i>Daldinia singularis</i>	BNT (1,1'-Binaphthalene-4,4'-5,5'-tetrol) (yellow), daldinol (dark brown), 8-methoxy-1-naphthol, 2-hydroxy-5-methylchromone	[25]
<i>Daldinia concentrica</i>	BNT (1,1'-Binaphthalene-4,4'-5,5'-tetrol) (yellow), daldinol, 8-methoxy-1-naphthol, 2-hydroxy-5-methylchromone, daldinal A–C (yellow), daldinin A–C (green-olivaceous-isabelline)	
<i>Daldinia eschscholzii</i>	BNT (1,1'-Binaphthalene-4,4'-5,5'-tetrol) (yellow), daldiol (dark brown), 8-methoxy-1-naphthol, 2-hydroxy-5-methylchromone, daldinal A–C (yellow)	

Table 1. Cont.

Fungal Species	Pigments	References
<i>Jackrogersella cohaerens</i> (Formerly Known as <i>Annulohypoxylon cohaerens</i> )		
<i>Jackrogersella cohaerens</i>	Cohaerin A	[25]
<i>Hypoxylon</i> species		
<i>Hypoxylon fragiforme</i>	Hypoxyxylone (green), fragiformins A–B, cytochalasin H (white), mitorubrin azaphilones (red)	
<i>Hypoxylon howeanum</i>	Mitorubrin azaphilones (red)	
<i>Hypoxylon lechatii</i>	Vermelhotin (orange-red), hypoxyvermelhotins A–C (orange-red)	
<i>Hypoxylon fuscum</i>	Daldinin A–C (green-olivaceous-isabelline)	
<i>Hypoxylon fulvo-sulphureum</i>	Mitorubrinol derivatives	[25]
<i>Hypoxylon sclerophaeum</i>	Hypoxytone (orange)	
<i>Hypoxylon rickii</i>	Rickenyl B (red), rickenyl D (brown)	
<i>Hypoxylon lenormandii</i> , <i>Hypoxylon jaklitschii</i>	Lenormandins A–G (yellow)	
<i>Hypoxylon rubiginosum</i>	Mitorubrin (orange), rubiginosin (orange-brown), hypomiltin (yellowish-green)	
<i>Alternaria</i> species		
<i>Alternaria alternata</i>	Alternariol (red), altenuene (red-violet), alternarienoic acid (red), alternariol-5-methyl ether (red-brown), tenuazoic acid (orange-red), alterperyleneol (red), stemphyperyleneol (yellow–orange-red)	[76]
<i>Alternaria dauci</i>	Uncharacterized (red)	[25,61]
<i>Alternaria porri</i>	Altersolanol A (yellow-orange), dactylariol	[25,61,77]
<i>Alternaria solani</i> , <i>Alternaria tomatophila</i>	Altersolanol A (yellow-orange)	[25,61]
<i>Alternaria</i> species	Alterperyleneol (red), dihydroalterperyleneol (dark purple)	[78]
<i>Alternaria</i> sp. ZJ9–6B	Alterporriol K–M (red)	[79]
<i>Curvularia</i> species		
<i>Curvularia lunata</i>	Chrysophanol (red), cynodontin (bronze), helminthosporin (maroon), erythroglaucin (red), catenarin (red)	[25,61]
<i>Sanghuangporus</i> species		
<i>Sanghuangporus baumii</i>	Uncharacterized (yellow)	[71]
<i>Clonostachys</i> species		
<i>Clonostachys intermedia</i>	Uncharacterized (yellow)	[71]
<i>Pyrenophora</i> species (Previously Known as species of <i>Drechslera</i> )		
<i>Pyrenophora teres</i> , <i>Pyrenophora graminea</i> , <i>Pyrenophora tritici-repentis</i> , <i>Pyrenophora grahamii</i> , <i>Pyrenophora dictyoides</i> , <i>Pyrenophora chaetomioides</i>	Catenarin (red), cynodontin (bronze), helminthosporin (maroon), tritispurin (reddish-brown), erythroglaucin (red)	[25,61]
<i>Exophiala</i> species		
<i>Exophiala dermatitidis</i> (formerly known as <i>Wangiella dermatitidis</i> )	Melanin (black-brown)	[44]
<i>Sporothrix</i> species		
<i>Sporothrix schenckii</i>	Melanin (black-brown)	[44]

Table 1. Cont.

Fungal Species	Pigments	References
<i>Cryptococcus</i> species		
<i>Cryptococcus neoformans</i>	Dihydroxy phenyl alanine-melanin	[29,80]
<i>Tuber</i> species		
<i>Tuber melanosporum</i>	Melanin (black)	[29,81]
<i>Polyporus</i> species		
<i>Lentinus brumalis</i> (formerly known as <i>Polyporus brumalis</i> )	Melanin (black)	[34,35]
<i>Cerioporus squamosus</i> (formerly known as <i>Polyporus squamosus</i> )	Melanin (black)	
<i>Xylaria</i> species		
<i>Xylaria polymorpha</i>	Melanin (black)	[34,35]
<i>Fomes</i> species		
<i>Fomes fomentarius</i>	Melanin (black)	[34,35]
<i>Oxyporus</i> species		
<i>Oxyporus populinus</i>	Melanin (black)	[34]
<i>Trametes</i> species		
<i>Trametes versicolor</i>	Melanin (black)	[34,35]
<i>Inonotus</i> species		
<i>Inonotus hispidus</i>	Melanin (black), uncharacterized (yellow)	[34–36]
<i>Chlorociboria</i> species		
<i>Chlorociboria aeruginascens</i>	Xylindein (green), xylindein quinol (yellow)	[33]
<i>Chlorociboria aeruginosa</i>	Xylindein (green)	[37,39]
<i>Scytalidium</i> species		
<i>Scytalidium cuboideum</i>	Draconin red (red)	[37,39]
<i>Scytalidium ganodermophthorum</i>	Uncharacterized (yellow)	[36,39]
<i>Scytalidium lignicola</i>	Uncharacterized (yellow)	[36,39]
<i>Epicoccum</i> species		
<i>Epicoccum nigrum</i>	Carotenoids, chromanone (yellow), epicoccarines A–B, epicocconone (fluorescent yellow), epipyridone (red), flavipin (brown), isobenzofuran derivatives (yellow to brown), orevactaene (yellow)	[41,61]
<i>Chaetomium</i> species		
<i>Chaetomium cupreum</i>	Oosporein (red), rotiorinols A–C (red), rubrorotiorin (red)	[25]
<i>Chaetomium globosum</i>	Chaetoviridins A–D (yellow), chaetoglobins A–B, chaetomugilins A–F, cochliodinol (purple)	
<i>Chaetomium</i> sp. NA-S01-R1	Chaephilone–C (yellow), chaetoviridides A–C (red)	[82]
<i>Achaetomium</i> species		
<i>Achaetomium</i> sp.	Parietin (orange)	[25]
<i>Phyllosticta</i> species		
<i>Phyllosticta capitalensis</i>	Melanin (black)	[83]
<i>Cladosporium</i> species		
<i>Cladosporium cladosporioides</i>	Calphostins A–D and I (red)	[61]

Table 1. Cont.

Fungal Species	Pigments	References
<b><i>Nodulisporium</i> species</b>		
<i>Nodulisporium hinnuleum</i>	Hinnuliquinone (red)	[84]
<b><i>Astrosphaeriella</i> species</b>		
<i>Astrosphaeriella papuana</i>	Astropaquinones A–C (orange)	[85]
<b><i>Arthrotrys</i> species</b>		
<i>Arthrotrys ferox</i>	Carotenoid	[86]
<b><i>Thelebolus</i> species</b>		
<i>Thelebolus microsporus</i>	$\beta$ -carotene (orange)	[86,87]
<b><i>Shiraia</i> species</b>		
<i>Shiraia bambusicola</i>	Shiraiarin (red), hypocrellin D (orange-red)	[88,89]
<b><i>Paecilomyces</i> species</b>		
<i>Paecilomyces sinclairii</i>	Uncharacterized (red) **	[25,61]
<b><i>Neurospora</i> species</b>		
<i>Neurospora crassa</i>	Neurosporaxanthin (yellow-orange), phytoene (yellow-orange), $\beta$ -carotene (red-orange), lycopene (red), neurosporen (yellow-orange), spirilloxanthin (violet), Y-carotene (yellow-orange), $\beta$ -carotene (yellow-orange) **	[25,90]
<i>Neurospora sitophila</i>	Neurosporaxanthin (yellow-orange)	[26]
<i>Neurospora intermedia</i>	Uncharacterized (yellow-orange), a mixture of carotenoids	
<b><i>Blakeslea</i> species</b>		
<i>Blakeslea trispora</i>	$\beta$ -carotene (yellow-orange) *, lycopene (red) *	[25]
<b><i>Ashbya</i> species</b>		
<i>Ashbya gossypi</i>	Riboflavin (yellow) *	[25]
<b><i>Phycomyces</i> species</b>		
<i>Phycomyces blakesleeanus</i>	$\beta$ -carotene (yellow-orange) **	[25]
<b><i>Mucor</i> species</b>		
<i>Mucor circinelloides</i>	$\beta$ -carotene (yellow-orange) ***	[25]
<b><i>Lactarius</i> species</b>		
<i>Lactarius</i> sp.	Azulenes (blue) **	[25]
<b><i>Penicillium</i> species</b>		
<i>Penicillium atramentosum</i>	Uncharacterized (dark brown)	
<i>Penicillium atosanguineum</i>	Phoenicin (red), uncharacterized (yellow and red)	
<i>Penicillium atrovenetum</i>	Atrovenetin (yellow), norherqueinone (red)	[61,91]
<i>Penicillium aurantiogriseum</i>	Uncharacterized	
<i>Penicillium brevicompactum</i> , <i>Penicillium simplicissimum</i>	Xanthoepocin (yellow)	
<i>Penicillium chrysogenum</i>	Sorbicillins (yellow), xanthocillin (yellow), chrysogine (yellow)	[61,92]
<i>Penicillium citrinum</i>	Anthraquinones (yellow), citrinin (yellow)	[61]
<i>Penicillium convolutum</i> (formerly known as <i>Talaromyces convolutus</i> )	Talaroconvolutins A–D, ZG-1494 $\alpha$	[93]
<i>Penicillium cyclopium</i>	Viomellein (reddish-brown), xanthomegnin (orange)	[61]
<i>Penicillium discolor</i>	Uncharacterized	
<i>Penicillium echinulatum</i>	Uncharacterized (yellow)	
<i>Penicillium flavigenum</i>	Xanthocillin (yellow), dihydrotrichodimerol (yellow)	[41,61]




Table 1. Cont.

Fungal Species	Pigments	References
<i>Penicillium</i> species		
<i>Penicillium freii</i> , <i>Penicillium viridicatum</i>	Viomellein (reddish-brown), viioxanthin, xanthomegnin (orange)	[61]
<i>Penicillium herquei</i>	Atrovenetin (yellow), herqueinones (red and yellow)	
<i>Penicillium melinii</i>	Atrovenetin (yellow)	[91]
<i>Penicillium miczynskii</i>	Uncharacterized (red)	[71]
<i>Penicillium mallochii</i>	Sclerotiorin (yellow)	[94]
<i>Penicillium oxalicum</i>	Arpink red™, anthraquinone derivative (red), secalononic acid D (yellow), anthraquinones (red and other hues) *	[25,61]
<i>Penicillium paneum</i>	Uncharacterized (red)	[61]
<i>Penicillium persicinum</i>	Uncharacterized (cherry red)	
<i>Penicillium</i> sp. AZ	PP-V (violet), PP-R (red)	[95]
<i>Penicillium</i> sp. (GBPI_P155)	Uncharacterized (orange)	[96]
<i>Penicillium</i> sp. NIOM-02	Uncharacterized (red)	[97]
<i>Penicillium</i> sp.	Uncharacterized (red)	[98,99]
<i>Talaromyces</i> species		
<i>Talaromyces aculeatus</i> (formerly known as <i>Penicillium aculeatum</i> )	Uncharacterized	[61]
<i>Talaromyces atroseus</i>	Mitorubrin (red), monascorubrin (red), PP-R (red), glauconic acid (red), purpuride (red), ZG-1494 $\alpha$ (red), azaphilones (red) ***	[25,100]
<i>Talaromyces albobiverticillius</i> , <i>Talaromyces amestolkiae</i> , <i>Talaromyces stollii</i>	<i>Monascus</i> -like azaphilones (red)	[25]
<i>Talaromyces cnidii</i> , <i>Talaromyces coalescens</i>	<i>Monascus</i> -like azaphilones (red), uncharacterized (red)	
<i>Talaromyces funiculosus</i> (formerly known as <i>Penicillium funiculosum</i> )	Ankaflavain (yellow), uncharacterized	[61]
<i>Talaromyces islandicus</i> (formerly known as <i>Penicillium islandicum</i> )	Emodin (yellow), skyrin (orange), erythroscopyrin (orange-red), luteoskyrin (yellow)	
<i>Talaromyces marneffei</i> (formerly known as <i>Penicillium marneffei</i> )	Monascorubramine (purple-red), mitorubrinol (orange-red), rubropunctatin (orange), purpactin, herqueinone like (brick red), secalononic acid D (yellow)	[61,101]
<i>Talaromyces pinophilus</i> (formerly known as <i>Penicillium pinophilum</i> )	Azaphilones, uncharacterized	[25,61]
<i>Talaromyces purpureogenus</i> (formerly known as <i>Penicillium purpureogenum</i> )	Mitorubrin (yellow), mitorubrinol (orange-red), PP-R (purple-red), purpurogenone (yellow-orange), rubropunctatin (red), N-glutarylmonascorubramine, N-glutaryl-rubropunctamine, uncharacterized (red), azaphilones (red) ***	[25,61,102–105]
<i>Talaromyces ruber</i> (formerly known as <i>Penicillium crateriforme</i> )	Uncharacterized, <i>Monascus</i> -like azaphilones	[25]
<i>Talaromyces rugulosus</i> (formerly known as <i>Penicillium rugulosum</i> )	Rugulosin (yellow)	[61]
<i>Talaromyces variabilis</i> (formerly known as <i>Penicillium variabile</i> )	Rugulosin (yellow)	[61]

Table 1. Cont.

Fungal Species	Pigments	References
<i>Talaromyces vericulosus</i>	Uncharacterized (red)	[106]
<i>Talaromyces</i> sp. DgCr22.1b	Talaroxanthone (yellow)	[107]
<i>Talaromyces siamensis</i> , <i>Talaromyces</i> sp.	Uncharacterized (red)	[71,108]
<i>Talaromyces</i> sp.	N-threonine rubropunctamine (red)	[72]
<b><i>Hamigera avellanea</i> (Formerly Known as <i>Talaromyces avellaneus</i>)</b>		
<i>Hamigera avellanea</i>	Emodin (yellow), erythroglaucon (red), catenarin (red)	[109]
<b><i>Aspergillus</i> species</b>		
<i>Aspergillus amstelodami</i>	Physcion (yellow), erythroglaucon (red), flavoglaucine (yellow), auroglaucon (orange-red)	[25]
<i>Aspergillus awamori</i>	Asperenone (yellow)	[110]
<i>Aspergillus chevalieri</i>	Physcion (yellow), erythroglaucon (red), flavoglaucine (yellow), auroglaucon (orange-red), catenarin (red), rubrocristin (red)	[25]
<i>Aspergillus cristatus</i>	Emodin (yellow), questin (yellow to orange-brown), erythroglaucon (red), physcion (yellow), catenarin (red), rubrocristin (red)	[25,61]
<i>Aspergillus echinulatum</i> , <i>Aspergillus glaber</i> , <i>Aspergillus spiculosus</i> , <i>Aspergillus umbrosus</i>	Erythroglaucon (red), physcion (yellow), catenarin (red), rubrocristin (red)	[25]
<i>Aspergillus fumigatus</i>	Melanin (dark brown-black)	[25,111]
<i>Aspergillus falconensis</i> , <i>Aspergillus fruticosus</i>	Falconensins A–H (yellow), falconensones A1 and B2 (yellow), zeorin (yellow)	[25]
<i>Aspergillus glaucus</i>	Physcion (yellow), emodin (yellow), questin (yellow to orange-brown), erythroglaucon (red), catenarin (red), rubrocristin (red), flavoglaucine (yellow), auroglaucon (orange-red), aspergin (yellow)	
<i>Aspergillus intermedius</i> , <i>Aspergillus leucocarpus</i> , <i>Aspergillus tonophilus</i>	Physcion (yellow), erythroglaucon (red)	[25,61]
<i>Aspergillus ochraceus</i>	Viomellein (reddish-brown), vioxanthin, xanthomegnin (orange)	
<i>Aspergillus melleus</i> , <i>Aspergillus sulphureus</i> , <i>Aspergillus westerdijkiae</i>	Viomellein (reddish-brown), rubrosulphin (red), viopurpurin (purple), xanthomegnin (orange)	
<i>Aspergillus nidulans</i>	Ascoquinone A (red), norsolorinic acid, sterigmatocystin (yellow), melanin (dark brown-black)	[25,112,113]
<i>Aspergillus niger</i>	Flavioline (orange-red), N-naphtho- $\gamma$ -pyrones (yellow), aspergillin (black), azanigerones A–F, asperenone (yellow), melanin (dark brown-black)	[25,61,110,114,115]
<i>Aspergillus nishimurae</i>	Anishidiol (yellow)	[116]
<i>Aspergillus parvathecia</i> , <i>Aspergillus rugulosus</i> , <i>Aspergillus versicolor</i>	Sterigmatocystin (yellow)	[25]
<i>Aspergillus purpureus</i>	Epurpurins A–C (yellow)	
<i>Aspergillus repens</i>	Emodin (yellow), physcion (yellow), erythroglaucon (red), catenarin (red), rubrocristin (red), questin (yellow to orange-brown)	
<i>Aspergillus ruber</i>	Catenarin (red), rubrocristin (red), emodin (orange), asperflavin (yellow), eurorubrin (Brown), questin (yellow to orange-brown), 3-O-( $\alpha$ -D-ribofuranosyl)-questin (orange), 2-O-methyl-9-dehydroxyeurotinone, 2-O-methyl-4-O-( $\alpha$ -D-ribofuranosyl)-9-dehydroxyeurotinone, 2-O-methyleurotinone	[25,117]
<i>Aspergillus sclerotioniger</i>	Uncharacterized (yellow)	[61]
<i>Aspergillus sclerotiorum</i>	Neoaspergilliac acid (yellow-green)	[91]
<i>Aspergillus terreus</i>	Uncharacterized (yellow)	[118]
<i>Aspergillus</i> sp.	Ferriaspergillin (red), ferrineoaspergillin (red)	[119]
<i>Aspergillus</i> sp.	Uncharacterized (yellow)	[120]


\* Industrial production (IP), \*\* research project (RP), \*\*\* development stage (DS).



**Figure 1.** Pigments reported from *Monascus* species (*M. ruber* and *M. purpureus*), re-drawn from [52,54,56–59].

Along with *Monascus*, many species of *Fusarium* have been reported for their capability to produce pigments. Studies have reported pigments such as bikaverin, nor-bikaverin, fusarubins, some naphthoquinone (8-*O*-methylbostrycoidin, 8-*O*-methylfusarubin, 8-*O*-methylnectriafurone, 8-*O*-methyl-13-hydroxynorjavanicin, 8-*O*-methylanhydrofusarubinlactol, and 13-hydroxynorjavanicin), and a novel isoquinoline-type, pigment 2-(4-((3*E*,5*E*)-14-aminotetradeca-3,5-dienyloxy)butyl)-1,2,3,4-tetrahydroisoquinolin-4-ol (ATDBTHIQN), from *Fusarium fujikuroi* (formerly known as *Fusarium moniliforme*) (Figure 2) [25,63,65]. Similarly, differently colored naphthoquinones [bostrycoidin, 9-*O*-methylfusarubin, 5-*O*-methyljavanicin, 8-*O*-methylbostrycoidin, 1,4-naphthalenedione-3,8-dihydroxy-5,7-dimethoxy-2-(2-oxopropyl), 5-*O*-methylsolaniol, and 9-*O*-methylanhydrofusarubin], two anthraquinone compounds [2-acetyl-3,8-dihydroxy-6-methoxy anthraquinone and 2-(1-hydroxyethyl)-3,8-dihydroxy-6-methoxy anthraquinone], and polyketide pigment (bikaverin) were reported from *Fusarium oxysporum* (Figure 2) [25,47,64,67]. Another species of *Fusarium*, *Fusarium graminearum*, has

been found to produce a variety of pigments such as 5-deoxybostrycoidin anthrone, 6-*O*-dimethyl-5-deoxybostrycoidin anthrone, purpurfusarin, 6-*O*-demethyl-5-deoxybostrycoidin, 5-deoxybostrycoidin, and aurofusarin (Figure 2) [25,64,66,121].



**Figure 2.** Pigments from fungal genera of Nectriaceae (*Fusarium*, *Fusicolla*, and *Albonectria*), re-drawn from [25,47,63,65,66,68].


A red pigment aurofusarin has been found to be produced by many species of *Fusarium* such as *Fusarium culmorum*, *Fusarium sporotrichioides*, *Fusarium acuminatum*, *Fusarium avenaceum*, *Fusarium poae*, *Fusarium crookwellens*, *Fusarium pseudograminearum*, *Fusarium sambucinum*, and *Fusarium tricinctum*. Bikaverin has been reported to be produced by *Fusarium lycopersici*, and *Fusarium vasinfectum*. *Fusarium solani* and *Fusarium verticillioides* (currently known as *F. fujikuroi*) have been described to produce both aurofusarin and bikaverin (Figure 2) [25]. Similarly, benzoquinone has been reported from *Fusarium* sp. JN158 (Figure 2) [68]. A study has shown that the synthesis of major *Fusarium* carotenoids (neurosporaxanthin and  $\beta$ -carotene) is induced by light via transcriptional induction of the structural genes *carRA*, *carB*, *carT*, and *carD* [43]. Similarly, other members of the fungal family Nectriaceae, such as *Albonectria rigidiuscula* and *Fusicolla aquaeductuum* (formerly known as *Fusarium decemcellulare* and *Fusarium aquaeductuum* respectively) were reported for their pigment production potential (Figure 2) [43,64]. Recently, the biosynthetic pathway of chrysogine mediated by two-module non-ribosomal peptide synthetase (NRPS) gene cluster was discovered in *Fusarium graminearum* in which enhanced chrysogine production was observed upon overexpression of NRPS14 [122].

Many investigations report *Penicillium* as potent producers of pigment [25,61,96–98], such as arpink red<sup>TM</sup> (first commercial red colorant), talaroconvolutins A–D, sclerotiorin, xanthoepocin, atrovenetin, and dihydrotrichodimerol discovered from *Penicillium oxalicum* var. *armeniaca*, *Penicillium convolutum* (formerly known as *Talaromyces convolutes*), *Penicillium mallochii*, *Penicillium simplicissimum*, *Penicillium melinii*, and *Penicillium flavigenum*, respectively (Figure 3a) [41,91,93,94,123]. An uncharacterized red pigment has been reported from *Penicillium miczynskii* [71]. Besides, many other *Monascus*-like pigments such as PP-V [(10Z)-12-carboxylmonascorubramine] and PP-R [(10Z)-7-(2-hydroxyethyl)-monascorubramine] have been reported from *Penicillium* (Figure 4) [95]. A biosynthetic pathway for the yellow pigment chrysogine from *Penicillium chrysogenum* has been proposed recently [92].


*Talaromyces* spp. have been reported as a source of pigments by many researchers. The pigment production ability of *Talaromyces purpureogenus* (formerly known as *Penicillium purpureogenum*) was evaluated by many researchers [102,104,105]. Studies report the production of a herqueinone-like pigment from *Talaromyces marneffeii* (formerly known as *Penicillium marneffeii*), *Monascus*-like azaphilone pigments (*N*-glutarylmonascorubramine and *N*-glutarylubropunctamine) from *Talaromyces purpureogenus* (formerly known as *Penicillium purpureogenum*), industrially important red pigments (mitorubrin, monascorubrin, PP-R, glauconic acid, purpuride, and ZG-1494 $\alpha$ ) from *Talaromyces atroseus*, trihydroxyanthraquinones (emodin, erythroglauclin, and catenarin) from *Talaromyces stipitatus*, and a xanthone dimer (talaroxanthone) from *Talaromyces* sp. (Figure 3b) [100,101,103,107,109]. An uncharacterized red pigment was discovered from *Talaromyces siamensis* under submerged fermentation [71]. Moreover, other species of *Talaromyces*, *Talaromyces aculeatus*, *Talaromyces atroseus*, *Talaromyces albobiverticillius*, *Talaromyces cnidii*, *Talaromyces coalescens*, *Talaromyces pinophilus*, *Talaromyces purpureogenus*, *Talaromyces funiculosus*, *Talaromyces amestolkiae*, *Talaromyces ruber*, *Talaromyces stollii*, and *Talaromyces verruculosus* have been reported to have the ability to produce *Monascus*-like azaphilone pigments (Figure 4) [25,106].

Several members of the genus *Aspergillus*, such as *Aspergillus niger*, have been known to synthesize a wide variety of pigments, such as aspergillin, asperenone, azaphilones (azanigerones A–F), and melanin (Figure 5a) [25,110,114,115]. *Aspergillus nidulans* was reported to produce ascoquinone A, norsolorinic acid, and melanin [25,112,113], whereas *Aspergillus fumigatus* was reported to produce melanin and melanin-like pigments [25,111]. In addition, a variety of other pigments such as asperenone, anishidiol, neoaspergillilic acid, sterigmatocystin, and an uncharacterized yellow pigment have been discovered from *Aspergillus nishimurae*, *Aspergillus awamori*, *Aspergillus sclerotiorum*, *Aspergillus versicolor*, and *Aspergillus terreus*, respectively [25,91,110,116,118]. Many other species of *Aspergillus* such as *Aspergillus glaucus*, *Aspergillus cristatus*, and *Aspergillus repens* have been reported to produce a variety of hydroxyanthraquinone pigments, emodin, physcion, questin, erythroglauclin, catenarin, and rubrocristin; while *Aspergillus melleus*, *Aspergillus ochraceus*, *Aspergillus sulphureus*, and *Aspergillus*

*westerdijkiae* have been described to be major producers of polyketide-based pigments (rubrosulfon, viomellein, viopurpurin, and xanthomegnin) (Figure 5a) [25]. In addition to this, other pigments such as ferriaspergillin, ferrineoaspergillin, and an uncharacterized yellow pigment have also been reported from the genus *Aspergillus* (Figure 5a) [119,120].



**Figure 3.** Pigments from the genera *Penicillium* and *Talaromyces*. (a) Different pigments produced by *Penicillium* species, re-drawn from [41,91,93,94,123]. (b) Various pigments produced by *Talaromyces* species, re-drawn from [100,101,107,109].




**Figure 4.** *Monascus*-like azaphilone pigments of *Penicillium* and *Talaromyces* species, re-drawn from [25,95,106].

Certain teleomorphic species of *Aspergillus* have been described as producers of a variety of pigments. Some of the well-known azaphilone pigments such as falconensins A–H, zeorin, falconensones A1 and B2 have been reported from *Emericella falconensis* and *Emericella fruticulosa* (currently known as *Aspergillus falconensis* and *Aspergillus fruticosus*, respectively), epurpurins A–C from *Emericella purpurea* (currently known as *Aspergillus purpureus*), and the pigment sterigmatocystin from *Emericella rugulosus*, *Emericella parvathecia*, and *Emericella nidulans* (currently known as *Aspergillus rugulosus*, *Aspergillus parvathecia*, and *Aspergillus nidulans*) (Figure 5c). Similarly, other *Aspergillus* spp. such as *Aspergillus amstelodami*, *Aspergillus chevalieri*, *Aspergillus glaucus*, *Aspergillus umbrosus*, *Aspergillus spiculosus*, *Aspergillus glaber*, *Aspergillus echinulatum*, *Aspergillus tonophilus*, *Aspergillus intermedius*, *Aspergillus leucocarpus*, *Aspergillus ruber*, and *Aspergillus cristatus* (which were formerly known as *Eurotium amstelodami*, *Eurotium chevalieri*, *Eurotium herbariorum*, *Eurotium umbrosus*, *Eurotium spiculosum*, *Eurotium spiculosum*, *Eurotium echinulatum*, *Eurotium tonophilum*, *Eurotium intermedium*, *Eurotium leucocarpum*, *Eurotium rubrum*, and *Eurotium cristatum*, respectively) have also been reported to produce pigments such as physcion, erythroglaucon, flavoglaucan, auroglaucon, catenarin, rubrocristin, and emodin (Figure 5b) [25].


Members of different genera of the fungal family Pleosporaceae (*Alternaria*, *Curvularia*, *Pyrenophora*, etc.) have immense potential for pigment production. Species of *Alternaria* such as *Alternaria alternata*, *Alternaria solani*, *Alternaria porri*, and *Alternaria tomatophila* have been reported to produce a variety of pigments such as dactylariol, alterperyleneol, dihydroalterperyleneol, alternariol, alternariol-5-methyl ether, altenuene, alternarienoic acid, tenuazoic acid, stemphyperyleneol, and altersolanol A (Figure 6) [25,76–78]. Also, other members of the Pleosporaceae, *Curvularia* and *Pyrenophora*, have been known to produce different types of pigments, e.g., *Curvularia lunata* produces hydroxyanthraquinone pigments such as chrysophanol, cynodontin, helminthosporin, erythroglaucon, and catenarin, whereas different species of *Pyrenophora* such as *Pyrenophora teres*, *Pyrenophora graminea*, *Pyrenophora tritici-repentis*, *Pyrenophora grahamii*, *Pyrenophora dictyoides*, and *Pyrenophora chaetomioides* (which were previously known as *Drechslera teres*, *Drechslera graminea*, *Drechslera tritici-repentis*, *Drechslera phlei*, *Drechslera dictyoides*, *Drechslera avenae*, respectively) have also been reported to produce hydroxyanthraquinone pigments such as cynodontin, erythroglaucon, catenarin, helminthosporin, and tritisorin (Figure 6) [25,61]. *Trichoderma*, a well-known bio-control agent, has been known to produce a variety of pigments [25,124]. Several hydroxyanthraquinones such as pachybasin, chrysophanol, emodin, T22 azaphilone, 1-hydroxy-3-methyl-anthraquinone, 2,4,5,7-tetrahydroxyanthraquinone,

1,3,6,8-tetrahydroxyanthraquinone, and 1,8-dihydroxy-3-methyl-anthraquinone, have been reported from different species of *Trichoderma* (*Trichoderma harzianum*, *Trichoderma polysporum*, *Trichoderma viride*, and *Trichoderma aureoviride*) (Figure 7a) [25], whereas *Trichoderma afrharzianum*, *Trichoderma pyramidale*, and *Trichoderma* sp. 1 are reported to produce uncharacterized yellow pigments in submerged fermentation [71]. Studies have also revealed that certain species of *Neurospora*, such as *Neurospora crassa*, *Neurospora sitophila*, and *Neurospora intermedia* produce a variety of carotenoids such as phytoene,  $\beta$ -carotene,  $\gamma$ -carotene, lycopene, neurosporene, and neurosporaxanthin (Figure 7b) [25,26,90].




**Figure 5.** Pigments from the genus *Aspergillus* and its teleomorphic genera. (a) Structures of pigments produced by *Aspergillus* species. (b) Pigments produced by species of *Eurotium* (teleomorph of *Aspergillus*). (c) Pigments produced by species of *Emericella* (teleomorph of *Aspergillus*), re-drawn from [25].






**Figure 6.** Pigments produced by members of the fungal family Pleosporaceae (species of *Alternaria*, *Curvularia*, *Astrospheariella*, and *Pyrenophora*), re-drawn from [25,76–78].



**Figure 7.** Pigments from other fungi. (a) Pigments from *Trichoderma* species, based on [25]. (b) Pigments from *Neurospora* species, re-drawn from [25,90].


Many genera of the Xylariaceae family, such as *Daldinia*, *Hypoxylon*, *Jackrogersella*, etc., have a great capability to synthesize pigments of very diverse colors and hues [25]. A variety of interesting pigments such as BNT (1,1'-Binaphthalene-4,4'-5,5'-tetrol), daldinol, daldinal A–C, and daldinin A–C have been reported from different species of *Daldinia*, such as *Daldinia bambusicola*, *Daldinia caldariorum*, *Daldinia concentrica*, *Daldinia eschscholzii*, *Daldinia childiae*, *Daldinia clavata*, *Daldinia fissa*, *Daldinia grandis*, *Daldinia*

*lloydii*, *Daldinia loculata*, *Daldinia petriniae*, *Daldinia singularis* (Figure 8a). Similarly, several cohaerin variants (cohaerin A–K), multiformin A, and sassafrins D have been obtained from *Jackrogersella cohaerens* (formerly known as *Annulohypoxylon cohaerens*) (Figure 8a). Besides this, several species of *Hypoxylon* were declared to produce diverse pigments e.g., *Hypoxylon fragiforme* (hypoxyxylone, cytochalasin H, fragiformins A–B, and mitorubrin), *Hypoxylon howeanum* (mitorubrin and azaphilones), *Hypoxylon lechatii* (vermelhotin and hypoxyvermelhotins A–C), *Hypoxylon fuscum* (daldinin A–C), *Hypoxylon fulvo-sulphureum* (mitorubrinol derivatives), *Hypoxylon sclerophaeum* (hypoxylone), *Hypoxylon rickii* (rickenyl B and D), *Hypoxylon lenormandii* and *Hypoxylon jaklitschii* (lenormandins A–G), *Hypoxylon rubiginosum* (mitorubrin, rubiginosin, and hypomiltin) (Figure 8a). Members of the Chaetomiaceae family also exhibit potential of pigment production. *Chaetomium cupreum* has been mentioned to produce red azaphilone pigments, oosporein, rotiorinols A–C, rubrorotiorin, whereas *Chaetomium globosum* produces yellow azaphilone pigments (chaetoviridins A–D), chaetoglobin A–B, chaetomugilins A–F, and cochliodinol (Figure 8b). Production of parietin (hydroxyanthraquinone pigment) has also been revealed from the *Achaetomium* sp. (Figure 8b) [25].




**Figure 8.** Pigments from the fungi of Xylariaceae and Chaetomiaceae families. (a) Pigments from members of the Xylariaceae family (species of *Daldinia*, *Hypoxylon*, and *Jackrogersella*), re-drawn from [25]. (b) Pigments from members of the Chaetomiaceae family (species of *Chaetomium* and *Achaetomium*) and Hypoxyllaceae, re-drawn from [25,84].

Also, the genera belonging to the family Cordycipitaceae such as *Torrubiella*, *Cordyceps*, *Beauveria*, *Hyperdermium*, and *Lecanicillium* have been revealed to be promising producers of bioactive pigments, e.g., tenellin and bassianin are reported from *Beauveria bassiana* and *Beauveria brongniartii* (formerly known as *Beauveria tenella*), pyridovericin and pyridomacrolidin from *Beauveria bassiana*, torrubiellones A–D from the genus *Torrubiella*, oosporein from *Lecanicillium aphanocladii*, whereas anthraquinone-related compounds are reported from *Cordyceps farinosa* (formerly known as *Isaria farinosa*) (Figure 9a) [41,73–75,125]. Similarly, the pigments erythrostrominone, 4-O-methyl erythrostrominone, deoxyerythrostrominone, deoxyerythrostrominol, epierythrostrominol, and 3,5,8-TMON (3,5,8-trihydroxy-6-methoxy-2-(5-oxohexa-1,3-dienyl)-1,4-naphthoquinone) have been reported from *Ophiocordyceps unilateralis* (formerly known as *Cordyceps unilateralis*), and skyrin from *Hyperdermium bertonii* (Figure 9a) [25].



**Figure 9.** Pigments from the fungi of the Cordycipitaceae family and some other group. **(a)** Pigments from members of the families Cordycipitaceae (species of *Beauveria*, *Torrubiella*, *Cordyceps*, *Hyperdermium*, and *Lecanicillium*) and Ophiocordycipitaceae (*Ophiocordyceps* sp.), re-drawn from [25,41,73–75,125]. **(b)** Pigments known from other groups of fungi (species of *Chlorociboria*, *Scytalidium*, and *Epicoccum*), re-drawn from [37,41].

Apart from this, studies have reported the production of the pigment xylindein from *Chlorociboria aeruginosa* and *Chlorociboria aeruginascens*, draconin red from *Scytalidium cuboideum*, and a yellow pigment from *Scytalidium ganodermorphothorum* and *Scytalidium lignicola*. Other pigments, such as oreovactaene produced from *Epicoccum nigrum*, emodin,  $\omega$ -hydroxyemodin, and emodic acid from *Hamigera avellanea* (formerly known as *Talaromyces avellaneus*) are also known (Figure 3b, Figure 9b) [33,36,37,39,41,109]. Recently, fungi such as *Sanghuangporus baumii* and *Clonostachys intermedia* have been found to produce a yellow pigment under submerged fermentation [71]. Production of melanin was reported from different groups of fungi such as *Phyllosticta capitalensis*, *Xylaria polymorpha*, *Trametes versicolor*, *Inonotus hispidus*, *Oxyporus populinus*, *Fomes fomentarius*, *Exophiala dermatitidis*, *Tuber melanosporum*, *Sporothrix schenckii*, and *Cryptococcus neoformans* [29,34,35,44,80,81,83]. Similarly, a study has shown the possible industrial application of the red pigment produced by *Paecilomyces sinclairii* [126]. Besides filamentous fungi, certain genera of yeasts (*Rhodotorula*, *Sporidiobolus*, *Sporobolomyces* and *Xanthophyllomyces*) have also been known as pigment producers. Different species of *Rhodotorula* (*Rhodotorula glutinis*, *Rhodotorula mucilaginosa* (syn. *Rhodotorula rubra*), *Rhodotorula babjevae*, *Rhodotorula toruloides* *Rhodotorula graminis*), *Sporidiobolus* (*Sporidiobolus pararoseus*, *Sporidiobolus johnsonii*), and *Sporobolomyces* (*Sporobolomyces uberrimus*, *Sporobolomyces salmonicolor*) have been reported to be prolific producers of torulin and torularhodin [127]. Researchers have discovered pigments such as  $\beta$ -carotene, torulene, and torularhodin from *Rhodotorula glutini* and multi-hydroxy carotenoids (4,4'-dihydroxy-nostoxanthin and 4-hydroxy-nostoxanthin) from *Xanthophyllomyces dendrorhous* (Figure 10) [13,128].



**Figure 10.** Pigments reported from yeasts such as *Rhodotorula glutini* and *Xanthophyllomyces dendrorhous*, re-drawn from [13,128].

In addition to terrestrial fungi, marine fungi are also very good producers of a variety of unique pigments having promising therapeutic and industrial applications [129,130]. Studies on marine fungi by many researchers have reported a wide range of pigments and hues, e.g., a variety of anthraquinone pigments [asperflavin, 2-O-methyleurotinone, questin, eurorubrin, 2-O-methyl-9-dehydroxyeurotinone, 2-O-methyl-4-O-( $\alpha$ -D-ribofuranosyl)-9-dehydroxyeurotinone, and 6,3-O-( $\alpha$ -D-ribofuranosyl)-questin] from the mangrove endophytic fungus *A. ruber* (formerly known as *Eurotium rubrum*), fusarnaphthoquinones B and fusarnaphthoquinones C from the sea fan-derived fungi *Fusarium* species, and bianthraquinone derivatives (alterporriol K, alterporriol L, and alterporriol M) from mangrove endophytic *Alternaria* sp. (Figure 11) [69,79,117]. Researchers have also investigated the red pigment production from mangrove fungus *Penicillium* sp. and a yellow pigment production from the marine sponge-associated fungus *Trichoderma parareesei* [70,99].




Figure 11. Pigments produced by marine fungal isolates, re-drawn from [69,72,79,82,117].

Also, many studies have revealed the production of polyketide pigments (*N*-threonine rubropunctamine) and chlorinated azaphilone pigments (chaephilone-C, chaetoviridides-A, chaetoviridides-B, chaetoviridides-C) from marine fungal isolates of *Talaromyces* spp. and *Chaetomium* sp., respectively (Figure 11) [72,82]. A recent study has reported a novel pigment, *N*-GABA-PP-V (6-[(Z)-2-Carboxyvinyl]-*N*-GABA-PP-V), along with *N*-threonine-monascorubramine, *N*-glutaryl-rubropunctamine, and PP-O from the marine-derived fungus *Talaromyces albobiverticillius* (Figure 11) [131]. Many antarctic fungi have also been discovered to produce pigments of different

chemical classes and characteristics. A number of yeast and filamentous fungi isolated from the different samples collected from Antarctic regions have been reported to produce a variety of pigments with different colors [86].

#### 4. Optimization for Enhancement of Pigment Production

Most of the investigators have focused their study on the enhancement of pigment production from different fungal strains such as *Monascus*, *Penicillium*, *Talaromyces*, *Fusarium*, etc., by optimizing various fermentation parameters such as media, media composition, pH, temperature, light intensity, orbital speed, etc. [26,132–135]. Some studies have reported about the assessment of the pigment production potential of different fungi on natural substrates (rice, corn, wheat, cassava, whole sorghum grain, dehulled sorghum grain, and sorghum bran) and on different agro-industrial residues (feather meal, fish meal, cheese whey, grape waste, soybean protein, soybean meal, chicken feather and rice husk, orange processing waste) [134,136–138]. Enhancement in xylindein production was reported in *Chlorociboria aeruginascens* upon addition of test woods (*Acer saccharum*, *Populus tremuloides*, spalted *P. tremuloides*, and *Ailanthus altissima*) in agar-based media [33].

Some studies have also evaluated the effect of different sugar sources such as glucose, fructose, lactose, sucrose, and maltose on pigment production by the species of *Monascus*. Results of these studies have shown that maximum pigment production was achieved in media with fructose as a carbon source for *M. purpureus*, and lactose as a carbon source for *M. ruber* [132,139]. Studies have also discovered that the addition of different nitrogen sources such as ammonium, peptone, sodium nitrate, glutamic acid, monosodium glutamate, 6-furtyrylaminopurine, and tryptophan could enhance the yield of pigment, alter the hue of the fermentation liquid, and also improve light stability of the pigments of *Monascus* species [132,140–143]. NaCl has been proved to be a very good enhancer that stimulates pigment production and inhibits citrinin production in *M. purpureus* without affecting the growth of the fungus [144]. A study on the effect of nutrients on pigment production of *C. aeruginascens* shows that high biomass but no pigment production was observed in media with high nutrient concentration, whereas low biomass and high pigmentation was observed in media with low nitrogen concentration [145]. Investigators have also found variations in the yield, color characteristics (hue and chroma values), and structure of the pigments of *Monascus* species with respect to the type of amino acids in the media [146,147]. Beside this, the pH of the media also plays an important role in pigment production. In the case of *Monascus* species (*M. purpureus*, *M. major*, and *M. rubiginosus*), pH optimization studies have shown that a low pH of the media increases pigment production [140,146,148]. Another study has revealed that the pH of the substrate plays an important role in melanin production by *X. polymorpha*, *T. versicolor*, *Ceriporus squamosus* (formerly known as *Polyporus squamosus*), *Lentinus brumalis* (formerly known as *Polyporus brumalis*), *F. fomentarius* and *I. hispidus*. The maximum pigment production was observed in the pH range from 4.5 to 5.5 [35]. Similar studies in other fungi such as *Penicillium purpurogenum*, *P. aculeatum*, *A. niger*, *Altemaria* sp., *Fusarium* sp., *C. aeruginascens*, have shown that the optimum pH for maximum pigment production varies with the fungal species in submerged fermentation [35,149–152].

Along with chemical parameters, physical parameters such as temperature, light intensity, color of light, agitation speed, and oxygen supply have an impact on pigment production. Studies have also been reported showing the influence of temperature on the biosynthesis of pigments by certain fungal isolates such as *M. ruber*, *T. purpureogenus* (formerly known as *P. purpurogenum*), *C. aeruginascens*, etc. [150,152,153]. Enhancement of yellow pigment production in a *Monascus anka* mutant strain under submerged fermentation using a two-stage agitation speed control strategy (400 rpm followed by 300 rpm) has been successfully reported [154]. A study has also revealed that a sufficient supply of oxygen is necessary for xylindein production by *C. aeruginascens* [152]. The impact of darkness and different color light on the yield of extracellular and intracellular pigment and biomass has been assessed by various investigators. Most of the studies have shown that incubation in total darkness resulted in enhanced biomass and pigment production [152,155,156]. Studies have also reported that

there is an enhancement in the pigment production in the case of *A. alternata* and *M. ruber* when exposed to blue and red light, respectively [156,157], and in *F. oxysporum* when exposed to blue and green light [158]. In contrast, reduction in biomass and pigment yield has been observed in *I. farinosa*, *E. nidulans*, *F. verticillioides*, *P. purpurogenum* (currently known as *C. farinosa*, *A. nidulans*, *F. fujikuroi*, *T. purpureogenus*, respectively), and *M. purpureus* when exposed to green and yellow light [155]. Light intensity has also been found to influence the growth and pigment production of *M. ruber* under submerged fermentation [156]. Another study on the influence of moisture content of wood substrate on fungal pigment production in spalted wood was described. Based on the results, low moisture content stimulates the pigmentation in *T. versicolor* and *X. polymorpha*, while enhanced pigment production was observed at higher moisture content in the case of *I. hispidus*, *L. brumalis* (formerly known as *P. brumalis*), *C. squamosus* (formerly known as *P. squamosus*), and *S. cuboideum* [34,159]. Optimization of pigment production by simultaneously altering the physical and chemical parameters has been explored by many investigators. Several studies have reported an enhancement of the yield of pigment and biomass from different fungal genera such as *Monascus*, *Penicillium*, *Fusarium*, *Alternaria*, etc., when the physical and chemical parameters were simultaneously altered [104,133,135,158,160–167].

Nowadays, co-culturing has been found to be an effective method for the activation of cryptic pathways via cell–cell interactions, which ultimately results in the production of novel secondary metabolites such as pigments from the fungi [168,169]. Studies have reported that the induction or enhancement in pigment production was possible using co-culturing of fungi with bacteria or yeast, but it was species-specific. In case of *Monascus* and *A. chevalieri*, co-culturing was found to be effective, whereas in case of *F. oxysporum*, the results were negative [158,170]. Co-culturing of *C. neoformans* with *Klebsiella aerogenes* led to synthesis of melanin by the fungus, using dopamine synthesized by bacteria [171]. Researchers have also found that many fungi produce different types of zone lines when co-cultured with other fungi. Zone lines are narrow, dark marks composed of pigments (primarily melanin) produced in decaying wood by fungi in response to other fungi, to self-isolate from other decaying fungi and protect their resources [172]. It has been observed that many white rot fungi such as *T. versicolor*, *Stereum gausapatum*, *Bjerkandera adusta*, *X. polymorpha*, and few brown rot fungi (*Poria weirii*, *Piptoporus betulinus*) produce zone lines upon detection of another fungus in their territory [173]. *T. versicolor* and *B. adusta* were found to be the best fungal pair which produce zone lines upon co-culturing, whereas *X. polymorpha* produces zone lines individually in the absence of other fungi [174]. This clearly reveals that the method of co-culturing of these fungi has a significant impact on their pigment production which supplies pigments used for coloring different types of woods in order to enhance their market value.

Various modes of cultivation and various methods and techniques of pigment extraction were investigated by several researchers to enhance fungal pigment production and recovery. Different strategies such as the use of different surfactants (Tween 80, Span 20, Triton X-100, and polyethylene glycol polymer 8000), different solvents (acetone, acetonitrile, chloroform, cyclohexane, chloramphenicol, dichloromethane, dimethyl sulfoxide, hexane, isooctane, methanol, methyl sulfoxide, pyridine, tetrahydrofuran, and water), and potential extraction techniques (pressurized liquid extraction technique) have also been assessed, compared, and confirmed by researchers for the rapid extraction and enhanced recovery of pigments from submerged fermentation [72,134,175–177]. Researchers also suggested the use of shake culture methods using water as a carrier instead of using wood-based malt–agar media for pigment production from wood-degrading fungi [178].

Genetic engineering techniques for enhanced pigment production in fungi have been reported [1,20,179]. Certain genetic approaches such as alteration or modifications of genes, cloning of genes, or elimination of non-essential genes (mycotoxins) have been investigated for increasing pigment production and reducing mycotoxins production in fungi [180–182]. The manipulation of biosynthetic pathways has also been investigated by researchers for boosting fungal pigment production. A study on *F. graminearum* has shown that the transcription factor AurR1 has a positive regulatory effect on the aurofusarin gene cluster, enhancing the production of aurofusarin [183].

A recent study on *Monascus* strains, revealed that transcription factors play an important regulatory role in pigment diversity [184]. More research on this aspect may lead to enhanced pigment production.

## 5. Applications or Biological Activities of Fungal Pigments

Many fungal pigments have been reported to have a variety of biological applications because of their different properties such as antimicrobial, antioxidant, anticancer, and cytotoxic activities in addition to coloring property [1,20,25,179]; however, the degree of purity of pigments investigated in the various studies is not always known.

### 5.1. Fungal Pigments as Food Colorants

The majority of work done on fungal pigments is related to their use as food colorants. The possibility of the use of fungal pigments in different industries, particularly in the food industry, has been revealed long ago by many researchers [9,25,46,48,179,185–187]. The potential of fungal pigments to be used as food colorants or as food additives in different food products has been assessed by many researchers [51,188]. Some of the fungal pigments have already entered into the market as food colorants such as *Monascus* pigments, arpink red from *P. oxalicum*, riboflavin from *Ashbya gossypii*, and  $\beta$ -carotene from *B. trispora* [12,25,189].

### 5.2. Fungal Pigments as Antimicrobial Agents

Numerous microbial pigments have been reported to possess many health benefits over synthetic pigments [8,14]. Several studies have proved that the pigments or pigment extracts of certain species of fungal genera (*Monascus*, *Fusarium*, *Talaromyces*, *Trichoderma*, *Penicillium*, and *Aspergillus*) and yeast *R. glutinis* possess antimicrobial activity against different pathogenic bacteria as well as yeast and fungi. All these studies suggest the potential use of bioactive pigments as food preservatives or as antibacterial ingredients in the food and pharmaceutical industries [19,66,70,82,135,166,189–194]. Similarly, the antimicrobial potential against selected pathogenic bacteria of different types of fabrics (cotton, silk, etc.) dyed with pigments of fungi (*A. alternata* and *Thermomyces* spp.) has also been evaluated, and positive results of these studies suggest their possible use in producing specific products for medical application, such as bandages, suture threads, face masks, etc. [195–197].

### 5.3. Fungal Pigments as Antioxidant Agents

It has been reported that microbial pigments such as carotenoids, violacein, and naphthoquinones have antioxidant potential. Many review articles mention the antioxidant potential of pigments from certain fungi and yeast [1,17,20,179,198,199]. Studies on assessment of the antioxidant activity of the pigments of certain fungi such as *Penicillium* (*P. miczynskii*, *P. purpureogenum*, *P. purpuroscens*, *Penicillium* sp.), *Fusarium* sp., *Thermomyces* sp., *Chaetomium* sp., *Sanghuangporus baumii*, *Stemphylium lycopersici*, and species of *Trichoderma* (*T. afroharzianum*, *Trichoderma* spp.) confirm the promising antioxidant potential and their possible applications in the healthcare industry [71,97,192,200,201].


### 5.4. Fungal Pigments as Cytotoxic Agents

The cytotoxic activity of pigments of certain fungal isolates (*F. oxysporum*, *T. verruculosus*, and *Chaetomium* spp.) has been assessed by many researchers using different methods such as sour orange seeds toxicity assay or yeast toxicity test (YTT) using *Saccharomyces cerevisiae*, brine shrimp lethality bioassay, or cell counting kit-8 (CCK-8) assay. These studies confirm the possible application of pigments in different industries, especially in health and pharmaceutical ones [47,82,106,202]. A latest study on the evaluation of dermal toxicity of pigments of *Thermomyces* spp. and *P. purpureogenum* in Wistar rats has revealed the nontoxic nature of pigments and suggested its potential application in cosmetics and dyeing [203].



### 5.5. Fungal Pigments as Anticancer Agents

Fungal pigments are known to possess anticancer/antitumor activity. Several studies have revealed the fungal pigments as a potential anticancer drug. Pigments of *Monascus* species (*M. purpureus* and *M. pilosus*) such as monascin, ankaflavin, monasphilone A–B, monasphilone A–B, monapilol A–D, and monapurone A–C have been proved to possess anticancer/antitumor potential against different types of cancers, such as mouse skin carcinoma, human laryngeal carcinoma, human colon adenocarcinoma, human hepatocellular carcinoma, and pulmonary adenocarcinoma (Figure 12) [32,56–58,204,205]. Besides *Monascus*, pigments from other fungi such as norsolorinic acid from *A. nidulans*, shiraiarin from *Shiraia bambusicola*, alterporriol K, alterporriol L, and alterporriol M from *Alternaria* spp., benzoquinone from *Fusarium* spp., and an uncharacterized red pigment from *F. chlamydosporum* have also been reported to have anticancer, antitumor, or antiproliferative activity mainly against human breast cancer cell lines (MCF-7, MDA-MB-435, and MCF-7 b), whereas hypocrellin D from *S. bambusicola* shows anticancer activity against other cancer cell lines (Bel-7721, A-549, and Anip-973) (Figure 12) [62,68,88,89,113].



**Figure 12.** Pigments from different taxonomic groups of fungi having promising anticancer or antitumor potential, re-drawn from [32,56–58,62,68,88,89,113,204,205].

### 5.6. Fungal Pigments in the Cosmetic Industry

As the demand for natural products is increasing in the market, cosmetic industries are also in search of new types of natural pigments to replace synthetic pigments. Among the natural pigments, the use of fungal pigments is also rapidly expanding in cosmetics because of their advantages. Fungal pigments, especially melanin, carotenoids, lycopene, etc., have been reported for their application in cosmetics, sunscreens, sun lotions, sunblocks, face creams, anti-aging facials, etc. [1,206,207]. Excitingly, some of the fungal pigments (*Monascus* pigments and *Monascus*-like pigments) have already entered the market for their application in cosmetics such as skin conditioning and skin care products, lipsticks, etc. [25].

### 5.7. Fungal Pigments in the Textile Industry

The textile industry is the largest industry after agriculture in terms of economic contribution and employment generation. It majorly depends on synthetic dyes for dyeing different types of fabrics (cotton, silk, and wool). Currently, natural pigments from fungi, with their many advantages (eco-friendly, non-toxic, easy degradation, high colorfastness, high staining capability, etc.) over hazardous synthetic pigments, have proven to be a good alternative to the synthetic dyes in the textile industry. Many investigations have shown that organic pigments produced by fungi have extensive applications in the textile industry [1,5,8,18,25,207].

The literature reveals that only a handful of studies have investigated the application of fungal pigments in the textile industry, especially for dyeing different types of fabrics, such as cotton, silk, and wool. Various studies on the dyeing potential of pigments of different species of fungal genera (*Monascus*, *Fusarium*, *Aspergillus*, *Penicillium*, *Talaromyces*, *Trichoderma*, *Alternaria*, *Curvularia*, *Chlorociboria*, *Scytalidium*, *Cordyceps*, *Acrostalagmus*, *Bisporomyces*, *Cunninghamella*, *Thermomyces*, and *Phymatotrichum*) for different types of fabrics such as wool, cotton yarn, silk, polyester, and nylon have been reported [37,42,47,106,108,124,195,196,208–211]. Studies on the dyeing potential of pigments from wood spalling fungi (red pigment from *S. cuboideum*, yellow pigment from *S. ganodermophthorum*, and green pigment *C. aeruginosa*) have shown the possible use of these pigments for dyeing bleached cotton, spun polyacrylic, spun polyamide (nylon 6.6), worsted wool, spun polyester (Dacron 54), and garment fabrics, because of their high stability and good colorfastness to washing [37,212]. Another study has revealed that natural oils cannot be used in conjunction with these fungal pigments, as these fungal pigments are unstable in natural oils [42]. Results of all these studies have shown that these fungal pigments have good color stability, colorfastness properties, and dye uptake potential. Moreover, these fungal pigments do not have any adverse effects on fabric and are non-toxic to human skin. Therefore, the scope of applications of fungal pigments has the opportunity to expand into the textile and clothing industry.

### 5.8. Fungal Pigments in Dyeing Woods or as Color Modifiers

Pigment produced by wood-decaying fungi such as *T. versicolor*, *X. polymorpha*, *I. hispidus*, *S. cuboideum*, *B. adusta*, *C. aeruginascens*, and *Arthrographis cuboidea* have been used for dyeing different types of wood samples to increase their commercial importance [173,174,213]. Researchers have successfully used the red, green, and yellow pigments obtained from *S. cuboideum*, *S. ganodermophthorum*, and *C. aeruginosa*, respectively, to attenuate the presence of blue stain on wood samples of *Pinus* spp. [39].

### 5.9. Fungal Pigments in (Opto) Electronics

A recent study of the (opto)electronic properties of blends of the pigment xylindein extracted from *C. aeruginosa* has revealed that this pigment has high photostability and electron mobility in amorphous films, which suggests its possible use for the development of sustainable, organic semiconductor materials [214,215].

## 6. Conclusions

Several advantages of fungal pigments over synthetic pigments have increased the demand for fungal pigments worldwide in recent years. This increased public awareness, eco-safety, and health concerns as well as the application of strict environmental and ecological rules and regulations, have challenged researchers to undertake both qualitative and quantitative research on pigments derived from clean, eco-friendly bio-resources, such as fungi, having minimal ecological negative impacts. Therefore, there is a necessity to explore other novel, safe pigments from the diverse taxonomic group of fungi, to meet the existing demand of eco-friendly pigments. Though several fungal strains are known as pigment producers, a large number of fungi have not been systematically explored for their pigment-producing capability. Therefore, there is a great need to explore the vast fungal diversity for rare, novel, safe pigments, using appropriate tools and techniques. A review of the literature revealed that most of the studies focused on the application of fungal pigments in the food and healthcare industries; however, fungal pigments need to pass toxicity tests and quality tests and receive many regulatory approvals before their final entry into the market as food colorants or as drugs. Therefore, the application of fungal pigments in these areas is quite difficult.

Moreover, meager studies on the applicability of fungal pigments in other areas such as textiles, paints, varnishes, and daily household utensils leave immense possibilities to explore the indigenous diversity of fungi for their pigment production potential and their applications in different sectors, including the textile industry. In addition to the coloring properties, the biological properties of fungal pigments may open new avenues for their use in the production of valuable textiles for medical use. This provides an extensive area of exploration to identify natural, eco-friendly fungal pigments and develop their diverse applications to satisfy the public interest and market demand.

**Author Contributions:** Writing—original draft preparation, A.C.L.; writing—review and editing, S.K.S., L.D., P.N.S.

**Funding:** This research received no external funding.

**Acknowledgments:** We thank Prashant Dhakephalkar, Director, MACS' Agharkar Research Institute, Pune, for encouragement and providing necessary facilities to carry out the research work. Ajay C. Lagashetti acknowledges CSIR (Council of Scientific and Industrial Research), New Delhi, for granting Junior Research Fellowship (JRF) and S. P. Pune University, Pune, for granting permission to register for Ph.D. degree. Ajay C. Lagashetti also thanks Siddharam Bagale for his major help in re-drawing chemical structures in manuscript. Laurent Dufossé deeply thanks the Conseil Régional de La Réunion, Réunion island, Indian Ocean, for continuous financial support of research activities dedicated to microbial pigments.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Rao, M.P.N.; Xiao, M.; Li, W.J. Fungal and bacterial pigments: Secondary metabolites with wide application. *Front. Microbiol.* **2017**, *8*, 1113.
2. Downham, A.; Collins, P. Coloring our foods in the last and next millennium. *Int. J. Food Sci. Technol.* **2000**, *35*, 5–22. [[CrossRef](#)]
3. Osman, M.Y.; Sharaf, I.A.; Osman, H.M.Y.; El-Khouly, Z.A.; Ahmed, E.I. Synthetic organic food coloring agents and their degraded products: Effects on human and rat cholinesterases. *Br. J. Biomed. Sci.* **2004**, *61*, 128–132. [[CrossRef](#)] [[PubMed](#)]
4. Babitha, S. Microbial pigments. In *Biotechnology for Agro-Industrial Residues Utilization*; Nigam, P.S., Pandey, A., Eds.; Springer: Dordrecht, The Netherlands, 2009; pp. 147–162.
5. Samanta, A.K.; Agarwal, P. Application of natural dyes on textiles. *Indian J. Fibre Text. Res.* **2009**, *34*, 384–399.
6. Ratna, P.B.S. Pollution due to synthetic dyes toxicity and carcinogenicity studies and remediation. *Int. J. Environ. Sci.* **2012**, *3*, 940–955.
7. Arora, S. Textile dyes: Its impact on the environment and its treatment. *J. Bioremediat. Biodegrad.* **2014**, *5*, 1. [[CrossRef](#)]
8. Akilandeswari, P.; Pradeep, B.V. Exploration of industrially important pigments from soil fungi. *Appl. Microbiol. Biotechnol.* **2016**, *100*, 1631–1643. [[CrossRef](#)]

9. Chattopadhyay, P.; Chatterjee, S.; Sen, S.K. Biotechnological potential of natural food grade biocolorants. *Afr. J. Biotechnol.* **2008**, *7*, 2972–2985.
10. Joshi, V.K.; Attri, D.; Bala, A.; Bhushan, S. Microbial pigments. *Indian J. Biotechnol.* **2003**, *2*, 362–369.
11. Aberoumand, A. A review article on edible pigments properties and sources as natural biocolorants in foodstuff and food industry. *World J. Dairy Food Sci.* **2011**, *6*, 71–78.
12. Dufossé, L. Microbial production of food grade pigments. *Food Technol. Biotechnol.* **2006**, *44*, 313–321.
13. Latha, B.V.; Jeevaratnam, K. Purification and characterization of the pigments from *Rhodotorula glutinis* DFR-PDY isolated from a natural source. *Glob. J. Biotechnol. Biochem.* **2010**, *5*, 166–174.
14. Nagpal, N.; Munjal, N.; Chatterjee, S. Microbial pigments with health benefits—A mini review. *Trends Biosci.* **2011**, *4*, 157–160.
15. Ahmad, W.A.; Ahmad, W.Y.W.; Zakaria, Z.A.; Yusof, N.Z. Isolation of pigment-producing bacteria and characterization of the extracted pigments. In *Application of Bacterial Pigments as a Colorant*; SpringerBriefs in Molecular Science; Springer: Berlin/Heidelberg, Germany, 2012; pp. 25–44.
16. Kirti, K.; Amita, S.; Priti, S.; Kumar, A.M.; Jyoti, S. Colorful world of microbes: Carotenoids and their applications. *Adv. Biol.* **2014**, *1*, 1–13. [[CrossRef](#)]
17. Mata-Gomez, L.C.; Montanez, J.C.; Mendez-Zavala, A.; Aguilar, C.N. Biotechnological production of carotenoids by yeasts: An overview. *Microb. Cell Fact.* **2014**, *13*, 12. [[CrossRef](#)]
18. Kumar, A.; Vishwakarma, H.S.; Singh, J.; Dwivedi, S.; Kumar, M. Microbial pigments: Production and their applications in various industries. *Int. J. Phram. Chem. Biol. Sci.* **2015**, *5*, 203–212.
19. Sarkar, S.L.; Saha, P.; Sultana, N.; Akter, S. Exploring textile dye from microorganisms, an eco-friendly alternative. *Microbiol. Res. J. Int.* **2017**, *18*, 1–9. [[CrossRef](#)]
20. Ramesh, C.; Vinithkumar, N.V.; Kirubakaran, R.; Venil, C.K.; Dufossé, L. Multifaceted applications of microbial pigments: Current knowledge, challenges and future directions for public health implications. *Microorganisms* **2019**, *7*, 186. [[CrossRef](#)]
21. Venil, C.K.; Zakaria, Z.A.; Ahmad, W.A. Bacterial pigments and their applications. *Process. Biochem.* **2013**, *48*, 1065–1079. [[CrossRef](#)]
22. Gupta, C.; Prakash, D.; Gupta, S. Natural useful therapeutic products from microbes. *J. Microbiol. Exp.* **2014**, *1*, 30–37. [[CrossRef](#)]
23. Numan, M.; Bashir, S.; Mumtaz, R.; Tayyab, S.; Rehman, N.U.; Khan, A.L.; Shinwari, Z.K.; Al-Harrasi, A. Therapeutic applications of bacterial pigments: A review of current status and future opportunities. *3 Biotech* **2018**, *8*, 207. [[CrossRef](#)]
24. Indra Arulselvi, P.; Umamaheswari, S.; Ranandkumar Sharma, G.; Karthik, C.; Jayakrishna, C. Screening of yellow pigment producing bacterial isolates from various eco-climatic areas and analysis of the carotenoid produced by the isolate. *J. Food Process. Technol.* **2014**, *5*, 292.
25. Caro, Y.; Venkatachalam, M.; Lebeau, J.; Fouillaud, M.; Dufossé, L. Pigments and colorants from filamentous fungi. In *Fungal Metabolites*; Merillon, J.-M., Ramawat, K.G., Eds.; Springer International Publishing: Cham, Switzerland, 2017; pp. 499–568.
26. Gmoser, R.; Ferreira, J.A.; Lennartsson, P.R.; Taherzadeh, M.J. Filamentous ascomycetes fungi as a source of natural pigments. *Fungal Biol. Biotechnol.* **2017**, *4*, 4. [[CrossRef](#)] [[PubMed](#)]
27. Manikprabhu, D.; Lingappa, K.  $\gamma$  Actinorhodin a natural and attorney source for the synthetic dye to detect acid production of fungi. *Saudi J. Biol. Sci.* **2013**, *20*, 163–168. [[CrossRef](#)] [[PubMed](#)]
28. Blanchette, R.A.; Wilmering, A.M.; Baumeister, M. The use of green-stained wood caused by the fungus *Chlorociboria* in intarsia masterpieces from the 15th century. *Holzforsch Int. J. Biol. Chem. Phys. Technol. Wood* **1992**, *46*, 225–232.
29. Butler, M.J.; Day, A.W. Fungal melanins: A review. *Can. J. Microbiol.* **1998**, *44*, 1115–1136. [[CrossRef](#)]
30. Sakaki, T.; Shibata, M.; Mukai, K.; Sakai, M.; Wakamatsu, K.; Miyauchi, S. *Chlorociboria aeruginosa* pigment as algicide. *Jpn. Kokai Tokkyo Koho JP* **2002**, 2002291493.
31. Carvalho, J.C.; Pandey, A.; Babitha, S.; Soccol, C.R. Production of *Monascus* biopigments: An overview. *Agro Food Ind. Hi-Tech* **2003**, *14*, 37–42.
32. Feng, Y.; Shao, Y.; Chen, F. *Monascus* pigments. *J. Appl. Microbiol. Biotechnol.* **2012**, *96*, 1421–1440. [[CrossRef](#)]
33. Robinson, S.C.; Tudor, D.; Snider, H.; Cooper, P.A. Stimulating growth and xylindein production of *Chlorociboria aeruginascens* in agar-based systems. *AMB Express* **2012**, *2*, 15. [[CrossRef](#)]

34. Tudor, D. Fungal Pigment Formation in Wood Substrate. Ph.D. Thesis, University of Toronto, Toronto, ON, Canada, 2013.
35. Tudor, D.; Robinson, S.C.; Cooper, P.A. The influence of pH on pigment formation by lignicolous fungi. *Int. Biodeterior. Biodegrad.* **2013**, *80*, 22–28. [[CrossRef](#)]
36. Robinson, S.C.; Tudor, D.; Zhang, W.R.; Ng, S.; Cooper, P.A. Ability of three yellow pigment producing fungi to colour wood under controlled conditions. *Int. Wood Prod. J.* **2014**, *5*, 103–107. [[CrossRef](#)]
37. Hinsch, E.M.; Chen, H.L.; Weber, G.; Robinson, S.C. Colorfastness of extracted wood-staining fungal pigments on fabrics: A new potential for textile dyes. *J. Text. Appar. Technol. Manag.* **2015**, *9*, 1–11.
38. Tam, W.T.E.; Tsang, C.C.; Lau, K.P.S.; Woo, C.Y.P. Polyketides, toxins, and pigments in *Penicillium marneffei*. *Toxins* **2015**, *7*, 4421–4436. [[CrossRef](#)] [[PubMed](#)]
39. Hernandez, V.A.; Galleguillos, F.; Robinson, S. Fungal pigments from spalting fungi attenuating blue stain in *Pinus* spp. *Int. Biodeterior. Biodegrad.* **2016**, *107*, 154–157. [[CrossRef](#)]
40. Robinson, S.C.; Michaelsen, H.; Robinson, J.C. *Spalted Wood, History, Science and Art of an Unique Material*; Schiffer Publishing: Atglen, PA, USA, 2016; pp. 1–288.
41. Souza, P.N.C.; Grigoletto, T.L.B.; Moraes, L.A.B.; Abreu, L.M.; Souza, L.H.; Santos, C.; Galvao, L.R.; Cardoso, P.G. Production and chemical characterization of pigments in filamentous fungi. *Microbiology* **2016**, *162*, 12–22. [[CrossRef](#)] [[PubMed](#)]
42. Agurto, M.E.P.; Gutierrez, S.M.V.; Chen, H.-L.; Robinson, S.C. Wood-rotting fungal pigments as colorant coatings on oil-based textile dyes. *Coatings* **2017**, *7*, 152. [[CrossRef](#)]
43. Avalos, J.; Pardo-Medina, J.; Parra-Rivero, O.; Ruger-Herreros, M.; Rodriguez-Ortiz, R.; Hornero-Mendez, D.; Limon, M.C. Carotenoid biosynthesis in *Fusarium*. *J. Fungi* **2017**, *3*, 39. [[CrossRef](#)]
44. Pombeiro-Sponchiado, S.R.; Sousa, G.S.; Andrade, J.C.; Lisboa, H.F.; Gonçalves, R.C. Production of melanin pigment by fungi and its biotechnological applications. In *Melanin*; Blumenberg, M., Ed.; InTechOpen: London, UK, 2017; pp. 45–75.
45. Vega Gutierrez, P.; Robinson, S. Determining the presence of spalted wood in spanish marquetry woodworks of the 1500s through the 1800s. *Coatings* **2017**, *7*, 188. [[CrossRef](#)]
46. Mortensen, A. Carotenoids and other pigments as natural colorants. *Pure Appl. Chem.* **2006**, *78*, 1477–1491. [[CrossRef](#)]
47. Nagia, F.A.; El-Mohamedy, R.S.R. Dyeing of wool with natural anthraquinone dyes from *Fusarium oxysporum*. *Dyes Pigm.* **2007**, *75*, 550–555. [[CrossRef](#)]
48. Mapari, S.A.S.; Thrane, A.U.; Meyer, A.S. Fungal polyketide azaphilone pigments as future natural food colorants? *Trends Biotechnol.* **2010**, *28*, 300–307. [[CrossRef](#)] [[PubMed](#)]
49. Dufossé, L.; Fouillaud, M.; Caro, Y.; Mapari, S.A.S.; Sutthiwong, N. Filamentous fungi are large-scale producers of pigments and colorants for the food industry. *Curr. Opin. Biotechnol.* **2014**, *26*, 56–61. [[CrossRef](#)] [[PubMed](#)]
50. Went, F.A.F.C. *Monascus purpureus* le champignon de l'ang-quac une nouvelle thelebole. *Ann. Des. Sci. Nat. Bot. Biol. Veg.* **1895**, *8*, 1–18.
51. Fabre, C.E.; Santerre, A.L.; Loret, M.O.; Baberian, R.; Pareilleux, A.; Goma, G.; Blanc, P.J. Production and food applications of the red pigments of *Monascus ruber*. *J. Food Sci.* **1993**, *58*, 1099–1102. [[CrossRef](#)]
52. Hajjaj, H.; Klaebe, A.; Loret, M.O.; Tzedakis, T.; Goma, G.; Blanc, P.J. Production and identification of N-glucosylrubropunctamine and N-glucosylmonascorubramine from *Monascus ruber* and occurrence of electron donor-acceptor complexes in these red pigments. *Appl. Environ. Microbiol.* **1997**, *63*, 2671–2678.
53. Lian, X.; Wang, C.; Guo, K. Identification of new red pigments produced by *Monascus ruber*. *Dyes Pigm.* **2007**, *73*, 121–125. [[CrossRef](#)]
54. Loret, M.O.; Morel, S. Isolation and structural characterization of two new metabolites from *Monascus*. *J. Agric. Food Chem.* **2010**, *58*, 1800–1803. [[CrossRef](#)]
55. Chen, W.; Chen, R.; Liu, O.; He, Y.; He, K.; Ding, X.; Kang, L.; Guo, X.; Xie, N.; Zhou, Y.; et al. Orange, red, yellow: Biosynthesis of azaphilone pigments in *Monascus* fungi. *Chem. Sci.* **2017**, *8*, 4917–4925. [[CrossRef](#)]
56. Hsu, Y.-W.; Hsu, L.-C.; Liang, Y.-H.; Kuo, Y.-H.; Pan, T.-M. Monaphilones A–C, three new antiproliferative azaphilone derivatives from *Monascus purpureus* NTU 568. *J. Agric. Food Chem.* **2010**, *58*, 8211–8216. [[CrossRef](#)]
57. Li, J.-J.; Shang, X.-Y.; Li, L.-L.; Liu, M.-T.; Zheng, J.-Q.; Jin, Z.L. New cytotoxic azaphilones from *Monascus purpureus*-fermented rice (red yeast rice). *Molecules* **2010**, *15*, 1958–1966. [[CrossRef](#)] [[PubMed](#)]

58. Hsu, Y.-W.; Hsu, L.-C.; Liang, Y.-H.; Kuo, Y.-H.; Pan, T.-M. New bioactive orange pigments with yellow fluorescence from *Monascus*-fermented Dioscorea. *J. Agric. Food Chem.* **2011**, *59*, 4512–4518. [[CrossRef](#)] [[PubMed](#)]
59. Mukherjee, G.; Singh, S.K. Purification and characterization of a new red pigment from *Monascus purpureus* in submerged fermentation. *Process Biochem.* **2011**, *46*, 188–192. [[CrossRef](#)]
60. Kaur, B.; Chakraborty, D.; Kaur, H. Production and evaluation of physicochemical properties of red pigment from *Monascus purpureus* MTCC 410. *Internet J. Microbiol.* **2008**, *7*, 1–6.
61. Mapari, S.A.S.; Meyer, A.S.; Thrane, U.; Frisvad, J.C. Identification of potentially safe promising fungal cell factories for the production of polyketide natural food colorants using chemotaxonomic rationale. *Microb. Cell Fact.* **2009**, *8*, 24. [[CrossRef](#)]
62. Soumya, K.; Narasimha Murthy, K.; Sreelatha, G.L.; Tirumale, S. Characterization of a red pigment from *Fusarium chlamydosporum* exhibiting selective cytotoxicity against human breast cancer MCF-7 cell lines. *J. Appl. Microbiol.* **2018**, *125*, 148–158. [[CrossRef](#)]
63. Steyn, P.S.; Wessels, P.L.; Marasas, W.F.O. Pigments from *Fusarium moniliforme* Sheldon: Structure and <sup>13</sup>C nuclear magnetic resonance assignments of an azaanthraquinone and three naphthoquinones. *Tetrahedron* **1979**, *35*, 1551–1555. [[CrossRef](#)]
64. Medenstev, A.G.; Arinbasarova, A.; Akimenko, V.K. Biosynthesis of naphthoquinone pigments by fungi of the genus *Fusarium*. *Prikl. Biokhim. Mikrobiol.* **2005**, *41*, 573–577.
65. Pradeep, F.S.; Palaniswamy, M.; Ravib, S.; Thangamanib, A.; Pradeep, B.V. Larvicidal activity of a novel isoquinoline type pigment from *Fusarium moniliforme* KUMBF1201 against *Aedes aegypti* and *Anopheles stephensi*. *Process Biochem.* **2015**, *50*, 1479–1486. [[CrossRef](#)]
66. Frandsen, R.J.N.; Rasmussen, S.A.; Knudsen, P.B.; Uhlig, S.; Petersen, D.; Lysoe, E.; Gotfredsen, C.H.; Giese, H.; Larsen, T.O. Black perithecial pigmentation in *Fusarium* species is due to the accumulation of 5-deoxybostrycoidin-based melanin. *Sci. Rep.* **2016**, *6*, 26206. [[CrossRef](#)]
67. Lebeau, J.; Petit, T.; Clerc, P.; Dufossé, L.; Caro, Y. Isolation of two novel purple naphthoquinone pigments concomitant with the bioactive red bikaverin and derivatives thereof produced by *Fusarium oxysporum*. *Biotechnol. Prog.* **2019**, *35*, e2738. [[CrossRef](#)] [[PubMed](#)]
68. Zheng, L.; Cai, Y.; Zhou, L.; Huang, P.; Ren, X.; Zuo, A.; Meng, X.; Xu, M.; Liao, X. Benzoquinone from *Fusarium* pigment inhibits the proliferation of estrogen receptor-positive MCF-7 cells through the NF-κB pathway via estrogen receptor signalling. *Int. J. Mol. Med.* **2017**, *39*, 39–46. [[CrossRef](#)] [[PubMed](#)]
69. Trisuwan, K.; Khamthong, N.; Rukachaisirikul, V.; Phongpaichit, S.; Preedanon, S.; Sakayaroj, J. Anthraquinone, cyclopentanone, and naphthoquinone derivatives from the sea fan-derived fungi *Fusarium* spp. PSU-F14 and PSU-F135. *J. Nat. Prod.* **2010**, *73*, 1507–1511. [[CrossRef](#)] [[PubMed](#)]
70. Sibero, M.T.; Triningsih, D.W.; Radjasa, O.K.; Sabdono, A.; Trianto, A. Evaluation of antimicrobial activity and identification of yellow pigmented marine sponge-associated fungi from Teluk Awur, Jepara, Central Java. *Indones. J. Biotechnol.* **2016**, *21*, 1–11. [[CrossRef](#)]
71. Heo, Y.M.; Kim, K.; Kwon, S.L.; Na, J.; Lee, H.; Jang, S.; Kim, C.H.; Jung, J.; Kim, J.J. Investigation of filamentous fungi producing safe, functional water-soluble pigments. *Mycobiology* **2018**, *46*, 269–277. [[CrossRef](#)]
72. Lebeau, J.; Venkatachalam, M.; Fouillaud, M.; Petit, T.; Vinale, F.; Dufossé, L.; Caro, Y. Production and new extraction method of polyketide red pigments produced by ascomycetous fungi from terrestrial and marine habitats. *J. Fungi* **2017**, *3*, 34. [[CrossRef](#)]
73. Velmurugan, P.; Lee, Y.H.; Nanthakumar, K.; Kamala-Kannan, S.; Dufossé, L.; Mapari, S.A.S.; Oh, B.T. Water-soluble red pigments from *Isaria farinosa* and structural characterization of the main colored component. *J. Basic Microbiol.* **2010**, *50*, 581–590. [[CrossRef](#)]
74. Wat, C.-K.; McInnes, A.G.; Smith, D.G.; Wright, J.L.C.; Vining, L.C. The yellow pigments of *Beauveria* species. Structures of tenellin and bassianin. *Can. J. Chem.* **1977**, *55*, 4090–4098. [[CrossRef](#)]
75. Isaka, M.; Chinthanom, P.; Supothina, S.; Tobwor, P. Pyridone and tetramic acid alkaloids from the spider pathogenic fungus *Torrubiella* sp. BCC 2165. *J. Nat. Prod.* **2010**, *73*, 2057–2060. [[CrossRef](#)]
76. Devi, S.; Kumar, H.A.K.; Ramachandran, G.; Subramanian, C.; Karuppan, P. Growth and mass spectrometry profile of *Alternaria alternata* pigment grown in maize grain extract. *J. Microbiol. Biotechnol. Food Sci.* **2014**, *4*, 179–184. [[CrossRef](#)]
77. Suemitsu, R.; Nakamura, A.; Isono, F.; Sano, T. Isolation and identification of Dactylariol from the culture liquid of *Alternaria porri* (Ellis) Ciferri. *Agric. Biol. Chem.* **1982**, *46*, 1693–1694. [[CrossRef](#)]

78. Okuno, T.; Natsume, I.; Sawai, K.; Sawamura, K.; Furusaki, A.; Matsumoto, T. Structure of antifungal and phytotoxic pigments produced by *Alternaria* sps. *Tetrahedron Lett.* **1983**, *24*, 5653–5656. [[CrossRef](#)]
79. Huang, C.H.; Pan, J.H.; Chen, B.; Yu, M.; Huang, H.B.; Zhu, X.; Lu, Y.J.; She, Z.G.; Lin, Y.C. Three bianthraquinone derivatives from the mangrove endophytic fungus *Alternaria* sp. ZJ9–6B from the South China Sea. *Mar. Drugs* **2011**, *9*, 832–843. [[CrossRef](#)] [[PubMed](#)]
80. Williamson, P.R.; Wakamatsu, K.; Ito, S. Melanin biosynthesis in *Cryptococcus neoformans*. *J. Bacteriol.* **1998**, *180*, 1570–1572.
81. Harki, E.; Talou, T.; Dargent, R. Purification, characterization and analysis of melanin extracted from *Tuber melanosporum* vitt. *Food Chem.* **1997**, *58*, 69–73. [[CrossRef](#)]
82. Wang, W.; Liao, Y.; Chen, R.; Hou, Y.; Ke, W.; Zhang, B.; Gao, M.; Shao, Z.; Chen, J.; Li, F. Chlorinated azaphilone pigments with antimicrobial and cytotoxic activities isolated from the deep sea-derived fungus *Chaetomium* sp. NA–S01–R1. *Mar. Drugs* **2018**, *16*, 61. [[CrossRef](#)]
83. Suryanarayanan, T.S.; Ravishankar, J.P.; Venkatesan, G.; Murali, T.S. Characterization of the melanin pigment of a cosmopolitan fungal endophyte. *Mycol. Res.* **2004**, *108*, 974–978. [[CrossRef](#)]
84. O’Leary, M.A.; Hanson, J.R.; Yeoh, B.L. The structure and biosynthesis of hinnuliquinone, a pigment from *Nodulisporium hinnuleum*. *J. Chem. Soc. Perkin Trans. 1* **1984**, *1*, 567–569. [[CrossRef](#)]
85. Wang, L.; Dong, J.Y.; Song, H.C.; Shen, K.Z.; Wang, L.M.; Sun, R.; Wang, C.R.; Gao, Y.X.; Li, G.H.; Li, L.; et al. Three new naphthoquinone pigments isolated from the freshwater fungus, *Astrosphaeriella papuana*. *Planta Med.* **2009**, *75*, 1339–1343. [[CrossRef](#)]
86. Duarte, A.W.F.; de Menezes, G.C.A.; e Silva, T.R.; Bicas, J.L.; Oliveira, V.M.; Rosa, L.H. Antarctic fungi as producers of pigments. In *Fungi of Antarctica*; Rosa, L., Ed.; Springer: Cham, Switzerland, 2019; pp. 305–318.
87. Singh, S.M.; Singh, P.N.; Singh, S.K.; Sharma, P.K. Pigment, fatty acid and extracellular enzyme analysis of a fungal strain *Thelebolus microsporus* from Larsemann Hills, Antarctica. *Polar Rec.* **2014**, *50*, 31–36. [[CrossRef](#)]
88. Fang, L.Z.; Qing, C.; Shao, H.J.; Yang, Y.D.; Dong, Z.J.; Wang, F.; Zhao, W.; Yang, W.Q.; Liu, J.K. Hypocrellin D, a cytotoxic fungal pigment from fruiting bodies of the ascomycete *Shiraia bambusicola*. *J. Antibiot.* **2006**, *59*, 351–354. [[CrossRef](#)] [[PubMed](#)]
89. Cai, Y.; Ding, Y.; Tao, G.; Liao, X. Production of 1, 5-dihydroxy-3-methoxy-7-methylanthracene-9, 10-dione by submerged culture of *Shiraia bambusicola*. *J. Microbiol. Biotechnol.* **2008**, *18*, 322–327. [[PubMed](#)]
90. Avalos, J.; Prado-Cabrero, A.; Estrada, A.F. Neurosporaxanthin production by *Neurospora* and *Fusarium*. In *Microbial Carotenoids from Fungi: Methods and Protocols*; Barredo, J.-L., Ed.; Springer Protocols: Totowa, NJ, USA, 2012; pp. 263–274.
91. Teixeira, M.F.S.; Martins, M.S.; Da Silva, J.C.; Kirsch, L.S.; Fernandes, O.C.C.; Carneiro, A.L.B.; De Conti, R.; Durrn, N. Amazonian biodiversity: Pigments from *Aspergillus* and *Penicillium*-characterizations, antibacterial activities and their toxicities. *Curr. Trends Biotechnol. Pharm.* **2012**, *6*, 300–311.
92. Viggiano, A.; Salo, O.; Ali, H.; Szymanski, W.; Lankhorst, P.P.; Nygard, Y.; Bovenberg, R.A.L.; Driessena, A.J.M. Pathway for the biosynthesis of the pigment Chrysogine by *Penicillium chrysogenum*. *Appl. Environ. Microbiol.* **2018**, *84*, 1–11. [[CrossRef](#)] [[PubMed](#)]
93. Suzuki, S.; Hosoe, T.; Nozawa, K.; Kawai, K.; Yaguchi, T.; Udagawa, S. Antifungal substances against pathogenic fungi, Talaroconvolutins, from *Talaromyces convolutus*. *J. Nat. Prod.* **2000**, *63*, 768–772. [[CrossRef](#)]
94. Santos, P.O.; Ferraz, C.G.; Soares, A.C.F.; Miranda, F.M.; da Silva, F.; de Abreu Roque, M.R. Sclerotiorin, a novel pigment from *Penicillium mallochii*. In Proceedings of the 6th Brazilian Conference on Natural Products, Federal University of Espirito Santo Victoria, Vitoria, Brazil, 5–8 November 2017.
95. Ogihara, J.; Kato, J.; Oishi, K.; Fujimoto, Y. PP-R, 7-(2-Hydroxyethyl)-Monascorubramine, a red pigment produced in the mycelia of *Penicillium* sp. *AZ. J. Biosci. Bioeng.* **2001**, *91*, 44–47. [[CrossRef](#)]
96. Pandey, N.; Jain, R.; Pandey, A.; Tamta, S. Optimisation and characterization of the orange pigment produced by a cold-adapted strain of *Penicillium* sp. (GBPI\_P155) isolated from mountain ecosystem. *Mycology* **2018**, *9*, 81–92. [[CrossRef](#)]
97. Dhale, M.A.; Vijay-Raj, A.S. Pigment and amylase production in *Penicillium* sp. NIOM-02 and its radical scavenging activity. *Int. J. Food Sci. Technol.* **2009**, *44*, 2424–2430. [[CrossRef](#)]
98. Jiang, Y.; Li, H.B.; Chen, F.; Hyde, K.D. Production potential of water-soluble *Monascus* red pigment by a newly isolated *Penicillium* sp. *J. Agric. Technol.* **2005**, *1*, 113–126.
99. Chintapenta, L.K.; Rath, C.C.; Maringinti, B.; Ozbay, G. Pigment production from a mangrove *Penicillium*. *Afr. J. Biotechnol.* **2014**, *13*, 2668–2674.

100. Frisvad, J.C.; Yilmaz, N.; Thrane, U.; Rasmussen, K.B.; Houbraken, J.; Samson, R.A. *Talaromyces atroroseus*, a new species efficiently producing industrially relevant red pigments. *PLoS ONE* **2013**, *8*, e84102. [[CrossRef](#)] [[PubMed](#)]
101. Bhardwaj, S.; Shukla, A.; Mukherjee, S.; Sharma, S.; Guptasarma, P.; Chakraborti, A.K.; Chakrabarti, A. Putative structure and characteristics of red water-soluble pigment secreted by *Penicillium marneffei*. *Med. Mycol.* **2007**, *45*, 419–427. [[CrossRef](#)] [[PubMed](#)]
102. Mendez-Zavala, A.; Contreras-Esquivel, J.C.; Lara-Victoriano, F.; Rodriguez-Herrera, R.; Aguilar, C.N. Fungal production of the red pigment using a xerophilic strain *Penicillium purpurogenum* GH-2. *Rev. Mex. Ing. Quim.* **2007**, *6*, 267–273.
103. Arai, T.; Koganei, K.; Umemura, S.; Kojima, R.; Kato, J.; Kasumi, T.; Ogihara, J. Importance of the ammonia assimilation by *Penicillium purpurogenum* in amino derivative *Monascus* pigment, PP-V production. *AMB Express* **2013**, *3*, 19. [[CrossRef](#)]
104. Sethi, B.K.; Parida, P.; Sahoo, S.L.; Dikshit, B.; Pradhan, C.; Sena, S.; Behera, B.C. Extracellular production and characterization of red pigment from *Penicillium purpurogenum* BKS9. *Alger. J. Nat. Prod.* **2016**, *4*, 379–392.
105. Ogbonna, C.N.; Aoyagi, H.; Ogbonna, J.C. Isolation and identification of *Talaromyces purpurogenus* and preliminary studies on its pigment production potentials in solid-state cultures. *Afr. J. Biotechnol.* **2017**, *16*, 672–682.
106. Chadni, Z.; Rahaman, M.H.; Jerin, I.; Hoque, K.M.F.; Reza, M.A. Extraction and optimization of red pigment production as secondary metabolites from *Talaromyces verruculosus* and its potential use in textile industries. *Mycology* **2017**, *8*, 48–57. [[CrossRef](#)]
107. Koolen, H.H.F.; Menezes, L.S.; Souza, M.P.; Silva, F.M.A.; Almeida, F.G.O.; de Souza, A.Q.L.; Nepel, A.; Barison, A.; da Silva, F.H.; Evangelistae, D.E.; et al. Talaroxanthone, a novel xanthone dimer from the endophytic fungus *Talaromyces* sp. associated with *Duguetia stelechantha* (Diels) R. E. Fries. *J. Braz. Chem. Soc.* **2013**, *24*, 880–883.
108. Morales-Oyervides, L.; Oliveira, J.; Sousa-Gallagher, M.; Mendez-Zavala, A.; Montanez, J.C. Assessment of the dyeing properties of the pigments produced by *Talaromyces* spp. *J. Fungi* **2017**, *3*, 38. [[CrossRef](#)]
109. Zhai, M.-M.; Li, J.; Jiang, C.-X.; Shi, Y.P.; Di, D.L.; Crews, P.; Wu, Q.-X. The bioactive secondary metabolites from *Talaromyces* species. *Nat. Prod. Bioprospect.* **2016**, *6*, 1–24. [[CrossRef](#)]
110. Pattenden, G. Synthesis of Asperenone, a new pigment from *Aspergillus niger* and *Aspergillus awamori*. *Tetrahedron Lett.* **1969**, *10*, 4049–4052. [[CrossRef](#)]
111. Youngchim, S.; Morris-Jones, R.; Hay, R.J.; Hamilton, A.J. Production of melanin by *Aspergillus fumigatus*. *J. Med. Microbiol.* **2004**, *53*, 175–181. [[CrossRef](#)] [[PubMed](#)]
112. Brown, D.W.; Solvo, J.J. Isolation and characterization of sexual spore pigments from *Aspergillus nidulans*. *Appl. Environ. Microbiol.* **1994**, *60*, 979–983. [[PubMed](#)]
113. Wang, C.C.; Chiang, Y.M.; Kuo, P.L.; Chang, J.K.; Hsu, Y.L. Norsolorinic acid from *Aspergillus nidulans* inhibits the proliferation of human breast adenocarcinoma MCF-7 cells via Fas-mediated pathway. *Basic Clin. Pharmacol. Toxicol.* **2008**, *102*, 491–497. [[CrossRef](#)] [[PubMed](#)]
114. Ray, A.C.; Eakin, R.E. Studies on the biosynthesis of Aspergillin by *Aspergillus niger*. *J. Appl. Microbiol.* **1975**, *30*, 909–915.
115. Zabala, A.O.; Xu, W.; Chooi, Y.-H.; Tang, Y. Discovery and characterization of a silent gene cluster that produces azaphilones from *Aspergillus niger* ATCC 1015 reveal a hydroxylation-mediated pyran-ring formation. *Chem. Biol.* **2012**, *19*, 1049–1059. [[CrossRef](#)] [[PubMed](#)]
116. Hosoe, T.; Mori, N.; Kamano, K.; Itabashi, T.; Yaguchi, T.; Kawai, K. A new antifungal yellow pigment from *Aspergillus nishimurae*. *J. Antibiot.* **2011**, *64*, 211–212. [[CrossRef](#)] [[PubMed](#)]
117. Li, D.-L.; Li, X.-M.; Wang, B.-G. Natural anthraquinone derivatives from a marine mangrove plant-derived endophytic fungus *Eurotium rubrum*: Structural elucidation and DPPH radical scavenging activity. *J. Biotechnol.* **2009**, *19*, 675–680.
118. Akilandeswari, P.; Pradeep, B.V. *Aspergillus terreus* KMBF1501 a potential pigment producer under submerged fermentation. *Int. J. Pharm. Pharm. Sci.* **2017**, *9*, 38–43. [[CrossRef](#)]
119. Assante, G.; Camarda, L.; Locci, R.; Merlini, L. Isolation and structure of red pigments from *Aspergillus flavus* and related species, grown on a differential medium. *J. Agric. Food Chem.* **1981**, *29*, 785–787. [[CrossRef](#)]
120. Narendrababu, B.N.; Shishupala, S. Spectrophotometric detection of pigments from *Aspergillus* and *Penicillium* isolates. *J. Appl. Biol. Biotechnol.* **2017**, *5*, 53–58. [[CrossRef](#)]



121. Cambaza, E. Comprehensive description of *Fusarium graminearum* pigments and related compounds. *Foods* **2018**, *7*, 165. [[CrossRef](#)] [[PubMed](#)]
122. Wollenberg, R.D.; Saei, W.; Westphal, K.R.; Klitgaard, C.S.; Nielsen, K.L.; Lysøe, E.; Gardiner, D.M.; Wimmer, R.; Sondergaard, T.S.; Sørensen, J.L. Chrysogine biosynthesis is mediated by a two-module nonribosomal peptide synthetase. *J. Nat. Prod.* **2017**, *80*, 2131–2135. [[CrossRef](#)] [[PubMed](#)]
123. Sardaryan, E. Strain of the Microorganism *Penicillium oxalicum* var. *Armeniaca* and Its. Application. Patent EP1070136B1, 4 August 2004.
124. Gupta, C.; Sharma, D.; Aggarwal, S.; Nagpal, N. Pigment production from *Trichoderma* spp. for dyeing of silk and wool. *Int. J. Sci. Nat.* **2013**, *4*, 351–355.
125. Takahashi, S.; Uchida, K.; Kakinuma, N.; Hashimoto, R.; Yanagisawa, T.; Nakagawa, A. The structures of Pyridovericin and Pyridomacrolidin, new metabolites from the entomopathogenic fungus, *Beauveria bassiana*. *J. Antibiot.* **1998**, *51*, 1051–1054. [[CrossRef](#)] [[PubMed](#)]
126. Cho, Y.J.; Park, J.P.; Hwang, H.J.; Kim, S.W.; Choi, J.W.; Yun, J.W. Production of red pigment by submerged culture of *Paecilomyces sinclairii*. *Lett. Appl. Microbiol.* **2002**, *35*, 195–202. [[CrossRef](#)] [[PubMed](#)]
127. Kot, A.M.; Błażejczak, S.; Gientka, I.; Kieliszek, M.; Bryś, J. Torulene and torularhodin: “new” fungal carotenoids for industry? *Microb. Cell Fact.* **2018**, *17*, 49. [[CrossRef](#)]
128. Pollmann, H.; Breitenbach, J.; Wolff, H.; Bode, H.B.; Sandmann, G. Combinatorial biosynthesis of novel multi-hydroxy carotenoids in the red yeast *Xanthophyllomyces dendrorhous*. *J. Fungi* **2017**, *3*, 9. [[CrossRef](#)]
129. Fouillaud, M.; Venkatachalam, M.; Girard-Valenciennes, E.; Caro, Y.; Dufossé, L. Anthraquinones and derivatives from marine-derived fungi: Structural diversity and selected biological activities. *Mar. Drugs* **2016**, *14*, 64. [[CrossRef](#)]
130. Fouillaud, M.; Venkatachalam, M.; Llorente, M.; Magalon, H.; Cuet, P.; Dufossé, L. Biodiversity of pigmented fungi isolated from the marine environment in La Reunion Island, Indian Ocean: New resources for colored metabolites. *J. Fungi* **2017**, *3*, 36. [[CrossRef](#)]
131. Venkatachalam, M.; Zelena, M.; Cacciola, F.; Ceslova, L.; Girard-Valenciennes, E.; Clerc, P.; Dugo, P.; Mondello, L.; Fouillaud, M.; Rotondo, A.; et al. Partial characterization of the pigments produced by the marine-derived fungus *Talaromyces albobiverticillius* 30548. Towards a new fungal red colorant for the food industry. *J. Food Compos. Anal.* **2018**, *67*, 38–47. [[CrossRef](#)]
132. Tseng, Y.Y.; Chen, M.T.; Lin, C.F. Growth, pigment production and protease activity of *Monascus purpureus* as affected by salt, sodium nitrite, polyphosphate, and various sugars. *J. Appl. Microbiol.* **2000**, *88*, 3–37. [[CrossRef](#)] [[PubMed](#)]
133. Chatterjee, S.; Maity, S.; Chattopadhyay, P.; Sarkar, A.; Laskar, S.; Sen, S.K. Characterization of red pigment from *Monascus* in submerged cultured pigment from *Monascus purpureus*. *J. Appl. Sci. Res.* **2009**, *5*, 2102–2108.
134. Carvalho, J.C.; Oishi, B.O.; Woiciechowski, A.L.; Pandey, A.; Babitha, S.; Soccol, C.R. Effect of substrates on the production of *Monascus* biopigments by solid-state fermentation and pigment extraction using different solvents. *Indian J. Biotechnol.* **2007**, *6*, 194–199.
135. Patil, S.A.; Sivanandhan, G.; Thakare, D.B. Effect of physical and chemical parameters on the production of red exopigment from *Penicillium purpurogenum* isolated from spoiled onion and study of its antimicrobial activity. *Int. J. Curr. Microbiol. Appl. Sci.* **2015**, *4*, 599–609.
136. Lopes, F.C.; Tichota, D.M.; Pereira, J.Q.; Segalin, J.; Rios, A.D.O. Pigment production by filamentous fungi on agro-industrial byproducts: An eco-friendly alternative. *Appl. Biochem. Biotechnol.* **2013**, *171*, 616–625. [[CrossRef](#)]
137. Srianta, I.; Zubaidah, E.; Estiasih, T.; Yamada, M. Comparison of *Monascus purpureus* growth, pigment production and composition on different cereal substrates with solid state fermentation. *Biocatal. Agric. Biotechnol.* **2016**, *7*, 181–186. [[CrossRef](#)]
138. Kantifedaki, A.; Kachrimanidou, V.; Mallouchos, A.; Papanikolaou, S.; Koutinas, A.A. Orange processing waste valorisation for the production of bio-based pigments using the fungal strains *Monascus purpureus* and *Penicillium purpurogenum*. *J. Clean. Prod.* **2018**, *185*, 882–890. [[CrossRef](#)]
139. Costa, J.P.V.; Vendruscolo, F. Production of red pigments by *Monascus ruber* CCT 3802 using lactose as a substrate. *Biocatal. Agric. Biotechnol.* **2017**, *11*, 50–55. [[CrossRef](#)]
140. Chen, M.; Johns, M.R. Effect of pH and nitrogen source on pigment production by *Monascus purpureus*. *Appl. Microbiol. Biotechnol.* **1993**, *40*, 132–138. [[CrossRef](#)]

141. Blanc, P.J.; Loret, M.O.; Santerre, A.L.; Pareilleux, A.; Prome, D.; Prome, J.C.; Laussac, J.P.; Goma, G. Pigments of *Monascus*. *J. Food Sci.* **1994**, *59*, 862–865. [[CrossRef](#)]
142. Pastrana, L.; Blanc, P.J.; Santerre, A.L.; Loret, M.O.; Goma, G. Production of red pigments by *Monascus ruber* in synthetic media with a strictly controlled nitrogen source. *Process Biochem.* **1995**, *30*, 333–341. [[CrossRef](#)]
143. Zhang, X.; Wang, J.; Chen, M.; Wang, C. Effect of nitrogen sources on production and photostability of *Monascus* pigments in liquid fermentation. *IERI Procedia* **2013**, *5*, 344–350. [[CrossRef](#)]
144. Zhen, Z.; Xiong, X.; Liu, Y.; Zhang, J.; Wang, S.; Li, L.; Gao, M. NaCl inhibits citrinin and stimulates *Monascus* pigments and monacolin K production. *Toxins* **2019**, *11*, 118. [[CrossRef](#)]
145. Stange, S.; Steudler, S.; Delenk, H.; Werner, A.; Walther, T.; Wagenführ, A. Influence of the nutrients on the biomass and pigment production of *Chlorociboria aeruginascens*. *J. Fungi* **2019**, *5*, 40. [[CrossRef](#)]
146. Carels, M.; Shepherd, D. The effect of pH and amino acids on conidiation and pigment production of *Monascus major* ATCC 16362 and *Monascus rubiginosus* ATCC 16367 in submerged shaken culture. *Can. J. Microbiol.* **1978**, *24*, 1346–1357. [[CrossRef](#)]
147. Jung, H.; Kim, C.; Kim, K.; Shin, C.S. Color characteristics of *Monascus* pigments derived by fermentation with various amino acids. *J. Agric. Food Chem.* **2003**, *51*, 1302–1306. [[CrossRef](#)]
148. Li, L.; Chen, S.; Gao, M.; Ding, B.; Zhang, J.; Zhou, Y.; Liu, Y.; Yang, H.; Wu, Q.; Chen, F. Acidic conditions induce the accumulation of orange *Monascus* pigments during liquid-state fermentation of *Monascus ruber* M7. *Appl. Microbiol. Biotech.* **2019**, *103*, 8393–8402. [[CrossRef](#)]
149. Mawthols, K.R.; Deshpande, R.; Ware, D.; Mahajan, M. Effect of pH on pigment production of fungi and their toxicity on seed germination. *Ecol. Environ. Conserv.* **2005**, *11*, 325–326.
150. Mendez, A.; Perez, C.; Montanez, J.C.; Martinez, G.; Aguilar, C.N. Red pigment production by *Penicillium purpurogenum* GH2 is influenced by pH and temperature. *J. Zhejiang Univ. Sci. B (Biomed. Biotechnol.)* **2011**, *12*, 961–968. [[CrossRef](#)]
151. Afsharia, M.; Shahidia, F.; Mortazavia, S.A.; Tabatabaia, F.; Es'haghib, Z. Investigating the influence of pH, temperature and agitation speed on yellow pigment production by *Penicillium aculeatum* ATCC 10409. *Nat. Prod. Res.* **2015**, *29*, 1300–1306. [[CrossRef](#)] [[PubMed](#)]
152. Stange, S.; Steudler, S.; Delenk, H.; Werner, A.; Walther, T.; Wagenführ, A. Influence of environmental growth factors on the biomass and pigment production of *Chlorociboria aeruginascens*. *J. Fungi* **2019**, *5*, 46. [[CrossRef](#)] [[PubMed](#)]
153. Huang, T.; Tan, H.; Chen, G.; Wang, L.; Wu, Z. Rising temperature stimulates the biosynthesis of water-soluble fluorescent yellow pigments and gene expression in *Monascus ruber* CGMCC10910. *AMB Express* **2017**, *7*, 134. [[CrossRef](#)] [[PubMed](#)]
154. Zhou, B.; Tian, Y.; Zhong, H. Application of a two-stage agitation speed control strategy to enhance yellow pigments production by *Monascus anka* mutant. *J. Microbiol. Biotechnol. Food Sci.* **2019**, *8*, 1260–1264. [[CrossRef](#)]
155. Velmurugan, P.; Lee, Y.H.; Venil, C.K.; Lakshmanaperumalsamy, P.; Chae, J.C.; Oh, B.T. Effect of light on growth, intracellular and extracellular pigment production by five pigment-producing filamentous fungi in synthetic medium. *J. Biosci. Bioeng.* **2010**, *109*, 346–350. [[CrossRef](#)] [[PubMed](#)]
156. Buhler, R.M.M.; Muller, B.L.; Moritz, D.E.; Vendruscolo, F.; Oliveira, D.; Ninow, J.L. Influence of light intensity on growth and pigment production by *Monascus ruber* in submerged fermentation. *Appl. Biochem. Biotechnol.* **2015**, *176*, 1277–1289. [[CrossRef](#)]
157. Haggblom, P.; Unestam, T. Blue light inhibits mycotoxin production and increases total lipids and pigmentation in *Alternaria alternata*. *Appl. Environ. Microbiol.* **1979**, *38*, 1074–1077.
158. Palacio-Barrera, A.M.; Areiza, D.; Zapata, P.; Atehortúa, L.; Correa, C.; Peñuela-Vásquez, M. Induction of pigment production through media composition, abiotic and biotic factors in two filamentous fungi. *Biotechnol. Rep.* **2019**, *21*, e00308. [[CrossRef](#)]
159. Tudor, D.; Robinson, S.C.; Cooper, P.A. The influence of moisture content variation on fungal pigment formation in spalted wood. *AMB Express* **2012**, *2*, 69. [[CrossRef](#)]
160. Gunsekaran, S.; Poorniammal, R. Optimization of fermentation conditions for red pigment production from *Penicillium* sp. under submerged cultivation. *Afr. J. Biotechnol.* **2008**, *7*, 1894–1898. [[CrossRef](#)]
161. Chutia, M.; Ahmed, G.U. Optimization of biomass and pigment production by *Penicillium* species isolated from virgin forest floor. *Biotechnol.* **2012**, *6*, 61–69.

162. Pradeep, F.S.; Pradeep, B.V. Optimization of pigment and biomass production from *Fusarium moniliforme* under submerged fermentation conditions. *Int. J. Pharm. Pharm. Sci.* **2013**, *5*, 526–535.
163. Ahmad, M.; Panda, B.P. Optimization of red pigment production by *Monascus purpureus* MTCC 369 under solid-state fermentation using response surface methodology. *Songklanakarin J. Sci. Technol.* **2014**, *36*, 439–444.
164. Devi, S.; Karuppan, P. Influence of culture condition and pH on growth and production of brown pigment from *Alternaria alternata*. *Int. J. Sci. Res.* **2014**, *3*, 458–461.
165. Santos-Ebinuma, V.C.; Roberto, I.C.; Teixeira, M.F.S.; Pessoa, J., Jr. Improvement of submerged culture conditions to produce colorants by *Penicillium purpurogenum*. *Braz. J. Microbiol.* **2014**, *45*, 731–742. [[CrossRef](#)] [[PubMed](#)]
166. Seyedin, A.; Yazdian, F.; Hatamian-Zarmi, A.; Rasekh, B.; Mir-derikvand, M. Natural pigment production by *Monascus purpureus*: Bioreactor yield improvement through statistical analysis. *Appl. Food Biotechnol.* **2015**, *2*, 23–30.
167. Patrovsky, M.; Sinovska, K.; Branska, B.; Patakova, P. Effect of initial pH, different nitrogen sources, and cultivation time on the production of yellow or orange *Monascus purpureus* pigments and the mycotoxin citrinin. *Food Sci. Nutr.* **2019**, 1–7. [[CrossRef](#)]
168. Serrano, R.; González-Menéndez, V.; Rodríguez, L.; Martín, J.; Tormo, J.R.; Genilloud, O. Co-culturing of fungal strains against *Botrytis cinerea* as a model for the induction of chemical diversity and therapeutic agents. *Front. Microbiol.* **2017**, *8*, 649. [[CrossRef](#)]
169. Tan, Z.Q.; Leow, H.Y.; Lee, D.C.W.; Karisnan, K.; Song, A.A.L.; Mai, C.W.; Yap, W.S.; Lim, S.H.E.; Lai, K.S. Co-culture Systems for the production of secondary metabolites: Current and future prospects. *Open Biotechnol. J.* **2019**, *13*, 18–26. [[CrossRef](#)]
170. Shin, C.S.; Kim, H.J.; Kim, M.J.; Ju, J.Y. Morphological change and enhanced pigment production of *Monascus* when cocultured with *Saccharomyces cerevisiae* or *Aspergillus oryzae*. *Biotechnol. Bioeng.* **1998**, *59*, 576–581. [[CrossRef](#)]
171. Frases, S.; Chaskes, S.; Dadachova, E.; Casadevall, A. Induction by *Klebsiella aerogenes* of a melanin-like pigment in *Cryptococcus neoformans*. *Appl. Environ. Microbiol.* **2006**, *72*, 1542–1550. [[CrossRef](#)] [[PubMed](#)]
172. Smith, K.T. Zone lines. In *Encyclopedia of Plant Pathology*; Malloy, O.C., Murray, T.D., Eds.; John Wiley & Sons: Hoboken, NJ, USA, 2001; Volume 2, pp. 1217–1218.
173. Robinson, S. The fine art of decay. *Am. Sci.* **2014**, *102*, 206–213. [[CrossRef](#)]
174. Robinson, S.C. Developing fungal pigments for “painting” vascular plants. *Appl. Microbiol. Biotechnol.* **2012**, *93*, 1389–1394. [[CrossRef](#)] [[PubMed](#)]
175. Robinson, S.C.; Hinsic, E.; Weber, G.L.; Freitas, S. Method of extraction and resolubilisation of pigments from *Chlorociboria aeruginosa* and *Scytalidium cuboideum*, two prolific spalling fungi. *Coloration Technol.* **2014**, *130*, 221–225. [[CrossRef](#)]
176. Morales-Oyervides, L.; Oliveira, J.; Sousa-Gallagher, M.; Mendez-Zavala, A.; Montanez, J.C. Perstraction of intracellular pigments through submerged fermentation of *Talaromyces* spp. in a surfactant-rich media: A novel approach for enhanced pigment recovery. *J. Fungi* **2017**, *3*, 33. [[CrossRef](#)]
177. Kaur, S.; Panesar, P.S.; Gurumayum, S.; Rasane, P.; Kumar, V. Optimization of aqueous extraction of oreveactaene and flavanoid pigments produced by *Epicoccum nigrum*. *Pigment Resin Technol.* **2019**, *48*, 301–308. [[CrossRef](#)]
178. Weber, G.L.; Boonloed, A.; Naas, K.M.; Koesdjojo, M.T.; Remcho, V.T.; Robinson, S.C. A method to stimulate production of extracellular pigments from wood-degrading fungi using a water carrier. *Curr. Res. Environ. Appl. Mycol.* **2016**, *6*, 218–230. [[CrossRef](#)]
179. Sen, T.; Barrow, C.J.; Deshmukh, S.K. Microbial pigments in the food industry—Challenges and the way forward. *Front. Nutr.* **2019**, *6*, 7. [[CrossRef](#)]
180. Jones, J.D.; Hohn, T.M.; Leathers, T.D. Genetically modified strains of *Fusarium sporotrichioides* for production of lycopene and  $\beta$ -carotene. In Proceedings of the Society of Industrial Microbiology Annual Meeting, San Diego, CA, USA, 29 July 2004; p. 91.
181. Fu, G.; Xu, Y.; Li, Y.; Tan, W. Construction of a replacement vector to disrupt pksCT gene for the mycotoxin citrinin biosynthesis in *Monascus aurantiacus* and maintain food red pigment production. *Asia Pacif. J. Clin. Nutr.* **2007**, *16*, 137–142.
182. Jia, X.Q.; Xu, Z.N.; Zhou, L.P.; Sung, C.K. Elimination of the mycotoxin citrinin production in the industrial important strain *Monascus purpureus* SM001. *Metab. Eng.* **2010**, *12*, 1–7. [[CrossRef](#)]

183. Westphal, K.; Wollenberg, R.; Herbst, F.A.; Sørensen, J.; Sondergaard, T.; Wimmer, R. Enhancing the production of the fungal pigment aurofusarin in *Fusarium graminearum*. *Toxins* **2018**, *10*, 485. [[CrossRef](#)] [[PubMed](#)]
184. Guo, X.; Li, Y.; Zhang, R.; Yu, J.; Ma, X.; Chen, M.; Wang, Y. Transcriptional regulation contributes more to *Monascus* pigments diversity in different strains than to DNA sequence variation. *World J. Microbiol. Biotechnol.* **2019**, *35*, 138. [[CrossRef](#)] [[PubMed](#)]
185. Dufossé, L.; Galaup, P.; Yaron, A.; Arad, S.M.; Blanc, P.; Murthy, K.N.C.; Ravishankar, G.A. Microorganisms and microalgae as sources of pigments for food use: A scientific oddity or an industrial reality? *Trends Food Sci. Technol.* **2005**, *16*, 389–406. [[CrossRef](#)]
186. Mapari, S.A.S.; Meyer, A.S.; Thrane, U. Colorimetric characterization for comparative analysis of fungal pigments and natural food colorants. *J. Agric. Food Chem.* **2006**, *54*, 7027–7035. [[CrossRef](#)]
187. Simpson, B.K.; Benjakul, S.; Klomklao, S. Chapter 37. Natural Food Pigments. In *Food Biochemistry and Food Processing*, 2nd ed.; Simpson, B.K., Ed.; Wiley-Blackwell: Hoboken, NJ, USA, 2012; pp. 704–722.
188. Fink-Gremmels, J.; Leistner, L. Biologische Wirkungen von *Monascus purpureus*. *Fleischwutsch* **1989**, *69*, 116–122.
189. Kim, D.; Ku, S. Beneficial effects of *Monascus* sp. KCCM 10093 pigments and derivatives: A mini review. *Molecules* **2018**, *23*, 98. [[CrossRef](#)]
190. Martinkova, L.; Juzlova, P.; Vesely, D. Biological activity of polyketide pigments produced by the fungus *Monascus*. *J. Appl. Bacteriol.* **1995**, *79*, 609–616. [[CrossRef](#)]
191. Vendruscolo, F.; Tosin, I.; Giachini, A.J.; Schmidell, W.; Ninow, J.L. Antimicrobial activity of *Monascus* pigments produced in submerged fermentation. *J. Food Process. Preserv.* **2014**, *38*, 1860–1865. [[CrossRef](#)]
192. Manon Mani, V.; Shanmuga Priya, M.; Dhaylini, S.; Preethi, K. Antioxidant and antimicrobial evaluation of bioactive pigment from *Fusarium* sp. isolated from the stressed environment. *Int. J. Curr. Microbiol. Appl. Sci.* **2015**, *4*, 1147–1158.
193. Saravanan, D.; Radhakrishnan, M. Antimicrobial activity of pigments produced by fungi from the Western Ghats. *J. Chem. Pharm. Res.* **2016**, *8*, 634–638.
194. Yolmeh, M.; Hamed, H.; Khomeiri, M. Antimicrobial activity of pigments extracted from *Rhodotorula glutinis* against some bacteria and fungi. *Zahedan J. Res. Med Sci.* **2016**, *18*, e4954. [[CrossRef](#)]
195. Poorniammal, R.; Parthiban, M.; Gunasekaran, S.; Murugesan, R.; Thilagavathi, G. Natural dye production from *Thermomyces* sp. fungi for textile application. *Indian J. Fibre Text. Res.* **2013**, *38*, 276–279.
196. Devi, S.; Karuppan, P. Reddish brown pigments from *Alternaria alternata* for textile dyeing and printing. *Indian J. Fibre Text. Res.* **2015**, *40*, 315–319.
197. Prathiban, M.; Thilagavathi, G.; Vijju, S. Development of antibacterial silk sutures using the natural fungal extract for healthcare applications. *J. Text. Sci. Eng.* **2016**, *6*, 249.
198. Tuli, H.S.; Chaudhary, P.; Beniwal, V.; Sharma, A.K. Microbial pigments as natural color sources: Current trends and future perspectives. *J. Food Sci. Technol.* **2015**, *52*, 4669–4678. [[CrossRef](#)] [[PubMed](#)]
199. Vendruscolo, F.; Buhler, R.M.M.; de Carvalho, J.C.; de Oliveira, D.; Moritz, D.E.; Schmidell, W.; Ninow, J.L. *Monascus*: A reality on the production and application of microbial pigments. *J. Appl. Biochem. Biotechnol.* **2016**, *178*, 211–223. [[CrossRef](#)]
200. Li, F.; Xue, F.; Yu, X. GC-MS, FTIR and Raman analysis of antioxidant components of red pigments from *Stemphylium lycopersici*. *Curr. Microbiol.* **2017**, *74*, 532–539. [[CrossRef](#)]
201. Poorniammal, R.; Prabhu, S.; Sakthi, A.R. Evaluation of in vitro antioxidant activity of fungal pigments. *Pharma Innov. J.* **2019**, *8*, 326–330.
202. Malik, K.; Tokas, J.; Anand, R.C. Characterization and cytotoxicity assay of pigment-producing microbes. *Int. J. Curr. Microbiol. Appl. Sci.* **2016**, *5*, 370–376. [[CrossRef](#)]
203. Poorniammal, R.; Prabhu, S.; Sakthi, A.R.; Gunasekaran, S. Subacute dermal toxicity of *Thermomyces* sp. and *Penicillium purpurogenum* pigments in wistar rats. *Int. J. Chem. Stud.* **2019**, *7*, 630–634.
204. Akihisa, T.; Tokuda, H.; Ukiya, M.; Kiyota, A.; Yasukawa, K.; Sakamoto, N.; Kimura, Y.; Suzuki, T.; Takayasu, J.; Nishino, H. Anti-tumor-initiating effects of Monascin, an azaphilone pigment from the extract of *Monascus pilosus* fermented rice (red-mold rice). *Chem. Biodivers.* **2005**, *2*, 1305–1309. [[CrossRef](#)] [[PubMed](#)]
205. Su, N.-W.; Lin, Y.-L.; Lee, M.-H.; Ho, C.-Y. Ankaflavin from *Monascus*-fermented red rice exhibits selective cytotoxic effect and induces cell death on Hep G2 cells. *J. Agric. Food Chem.* **2005**, *53*, 1949–1954. [[CrossRef](#)] [[PubMed](#)]

206. Hill, H. The function of melanin or six blind people examine an elephant. *BioEssays* **1992**, *14*, 49–56. [[CrossRef](#)] [[PubMed](#)]
207. Sajid, S.; Akber, N. Applications of fungal pigments in biotechnology. *Pure Appl. Biol.* **2018**, *7*, 922–930. [[CrossRef](#)]
208. Velmurugan, P.; Kim, M.J.; Park, J.S.; Karthikeyan, K.; Lakshmanaperumalsamy, P.; Lee, K.J.; Park, Y.J.; Oh, B.T. Dyeing of cotton yarn with five water-soluble fungal pigments obtained from five fungi. *Fibers Polym.* **2010**, *11*, 598–605. [[CrossRef](#)]
209. Mabrouk, A.M.; El-Kkhrisy, E.A.M.; Youssef, Y.A.; Asem, M.A. Production of textile reddish brown dyes by fungi. *Malays. J. Microbiol.* **2011**, *7*, 33–40.
210. Sharma, D.; Gupta, C.; Aggarwal, S.; Nagpal, N. Pigment extraction from fungus for textile dyeing. *Indian J. Fibre Text. Res.* **2012**, *37*, 68–73.
211. Aishwarya, A.D. Extraction of natural dyes from fungus—An alternate for textile dyeing. *J. Nat. Sci. Res.* **2014**, *4*, 1–6.
212. Weber, G.; Chen, H.L.; Hinsch, E.; Freitas, S.; Robinson, S. Pigments extracted from the wood-staining fungi *Chlorociboria aeruginosa*, *Scytalidium cuboideum*, and *S. ganodermophthorum* show potential for use as textile dyes. *Coloration Technol.* **2014**, *130*, 445–452. [[CrossRef](#)]
213. Robinson, S.C.; Tudor, D.; Cooper, P.A. Utilizing pigment-producing fungi to add commercial value to American beech (*Fagus grandifolia*). *Appl. Microbiol. Biotechnol.* **2012**, *93*, 1041–1048. [[CrossRef](#)]
214. Giesbers, G.; Van Schenck, J.; Gutierrez, S.V.; Robinson, S.; Ostroverkhova, O. Fungi-derived pigments for sustainable organic (opto) electronics. *MRS Adv.* **2018**, *3*, 3459–3464. [[CrossRef](#)]
215. Giesbers, G.; Van Schenck, J.; Quinn, A.; Van Court, R.; Vega Gutierrez, S.M.; Robinson, S.C.; Ostroverkhova, O. Xylindein: Naturally produced fungal compound for sustainable (opto) electronics. *ACS Omega* **2019**, *4*, 13309–13318. [[CrossRef](#)] [[PubMed](#)]



© 2019 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).