



HAL
open science

Trees as net sinks for methane (CH₄) and nitrous oxide (N₂O) in the lowland tropical rain forest on volcanic Reunion Island

Katerina Machacova, Libor Borak, Thomas Agyei, Thomas Schindler, Kaido Soosaar, Ülo Mander, Claudine Ah-Peng

► To cite this version:

Katerina Machacova, Libor Borak, Thomas Agyei, Thomas Schindler, Kaido Soosaar, et al.. Trees as net sinks for methane (CH₄) and nitrous oxide (N₂O) in the lowland tropical rain forest on volcanic Reunion Island. *New Phytologist*, 2020, 229, pp.1993-1994. 10.1111/nph.17002 . hal-03047906

HAL Id: hal-03047906

<https://hal.univ-reunion.fr/hal-03047906v1>

Submitted on 9 Dec 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Trees as net sinks for methane (CH₄) and nitrous oxide (N₂O) in the lowland tropical rain forest on volcanic Réunion Island

Katerina Machacova¹ , Libor Borak¹, Thomas Agyei¹ , Thomas Schindler^{1,2}, Kaido Soosaar^{1,2} , Ülo Mander^{1,2}  and Claudine Ah-Peng³

¹Global Change Research Institute of the Czech Academy of Sciences, Belidla 986/4a, Brno CZ-60300, Czech Republic; ²Department of Geography, Institute of Ecology & Earth Sciences, University of Tartu, 46 Vanemuise, Tartu EST-51014, Estonia; ³UMR PVBMT, Université de La Réunion, 7 chemin de l'IRAT, Saint-Pierre, La Réunion F-97410, France

Summary

Author for correspondence:
Katerina Machacova
Email: machacova.k@czechglobe.cz

Received: 2 July 2020
Accepted: 1 October 2020

New Phytologist (2020)
doi: 10.1111/nph.17002

Key words: basaltic lava flows, cryptogams, methane flux, nitrous oxide flux, soil, tree stem, tropical lowland rain forest, uptake.

- Trees are known to emit methane (CH₄) and nitrous oxide (N₂O), with tropical wetland trees being considerable CH₄ sources. Little is known about CH₄ and especially N₂O exchange of trees growing in tropical rain forests under nonflooded conditions.
- We determined CH₄ and N₂O exchange of stems of six dominant tree species, cryptogamic stem covers, soils and volcanic surfaces at the start of the rainy season in a 400-yr-old tropical lowland rain forest situated on a basaltic lava flow (Réunion Island). We aimed to understand the unknown role in greenhouse gas fluxes of these atypical tropical rain forests on basaltic lava flows.
- The stems studied were net sinks for atmospheric CH₄ and N₂O, as were cryptogams, which seemed to be co-responsible for the stem uptake. In contrast with more commonly studied rain forests, the soil and previously unexplored volcanic surfaces consumed CH₄. Their N₂O fluxes were negligible.
- Greenhouse gas uptake potential by trees and cryptogams constitutes a novel and unique finding, thus showing that plants can serve not only as emitters, but also as consumers of CH₄ and N₂O. The volcanic tropical lowland rain forest appears to be an important CH₄ sink, as well as a possible N₂O sink.

Introduction

Covering an area of *c.* 1730 million ha, tropical forests comprise 45% of the global forested area (D'Annunzio *et al.*, 2017). The most extensive tropical forest type is tropical rain forest, which encompasses almost 60% of all tropical forest area (Shvidenko *et al.*, 2005). These forests are found predominantly in South America, Africa and Asia, and are characterized as closed-canopy evergreen broadleaf forests with minimum annual temperature and precipitation of 25°C and 1500 mm, respectively (Richards, 1996). In general, tropical forests (including all kinds of forest types) constitute important carbon (C) stock (428 Gt-C in vegetation and soil) and C sink (−0.37 t-C ha^{−1} yr^{−1}; Dalal & Allen, 2008). Furthermore, they are considered to be a natural source of nitrous oxide (N₂O), and a natural sink and source of methane (CH₄), both of which are important greenhouse gases (GHG) with global warming potential of 265–298 and 28–36 over 100 yr, respectively (Myhre *et al.*, 2013). Net N₂O emission from tropical forest soils is estimated to be 4.76 kg ha^{−1} yr^{−1}, whereas net soil CH₄ consumption, without considering possible canopy fluxes from trees (Keppler *et al.*, 2006), seems to be −3.86 kg ha^{−1} yr^{−1} (Dalal & Allen, 2008). In the case of CH₄, soils of tropical rain forests growing under submerged conditions (e.g. Amazonia) can be substantial CH₄ sources as well, as a

consequence of the prevailing anaerobic conditions required for CH₄ production (Pangala *et al.*, 2017). Tropical forest soils have the highest N₂O emission potential among natural forest ecosystems (Dalal & Allen, 2008), and therefore tropical forests play an essential role in global N₂O inventories.

Nitrous oxide is naturally produced in soils through a wide range of nitrogen (N) turnover processes having different soil water content optima, including aerobic nitrification, anaerobic denitrification, and also dissimilatory nitrate reduction to ammonium in suboxic conditions (Espenberg *et al.*, 2018). The denitrification processes are the only processes known to reduce N₂O to dinitrogen (N₂) (Smith *et al.*, 2003). By contrast, CH₄ is produced by strictly anaerobic methanogenesis in water-saturated soils and is oxidized by methanotrophic bacteria (Smith *et al.*, 2003).

Both gases can be released into the atmosphere by gas diffusion at the soil surface and by ebullition in the case of flooded areas. Plants, moreover, can contribute to ecosystem N₂O and CH₄ exchange by: (1) taking up both gases from the soil via the root system and transporting them into the atmosphere through the transpiration stream or aerenchyma system and enlarged intercellular spaces (Rusch & Rennenberg, 1998; Machacova *et al.*, 2013); (2) producing N₂O and CH₄ directly in plant tissues (Smart & Bloom, 2001; Keppler *et al.*, 2006); (3) consuming

N_2O and CH_4 from the atmosphere by a nonspecified mechanism (Sundqvist *et al.*, 2012; Machacova *et al.*, 2016b, 2017, 2019); and (4) altering the N and C turnover processes in adjacent soil (Menyailo & Hungate, 2005; Yu & Chen, 2009). Moreover, cryptogamic stem covers (i.e. photoautotrophic organisms associated with tree bark, such as lichens, liverworts, mosses or ferns) also might contribute to N_2O and CH_4 exchange of trees and forest ecosystems (Lenhart *et al.*, 2015; Machacova *et al.*, 2017). Especially (but not solely) in tropical rain forests, cryptogams often grow on as much as 100% of the tree bark surface and all the way up to the crowns. These organisms remain overlooked as potential players in the trace gas exchange of forests and trees, even though they are present in the majority of tree stem chamber measurements and can, therefore, contribute to the gas exchange between the bark's surface and the chamber headspace. In summary, the net exchange of trace gases at the soil–plant–atmosphere interfaces results from a balance of simultaneously ongoing processes of gas production and consumption, gas transport within the relevant system, gas emission into the atmosphere, and gas uptake from the atmosphere, all of which together determine whether the forest compartment will be a source or sink of CH_4 and N_2O (Barba *et al.*, 2019).

Recent research in various climatic zones has revealed that not only soils and herbaceous plants but also woody plants can be significant sources of N_2O and CH_4 to the atmosphere (Machacova *et al.*, 2013, 2016a,b, 2019; Pangala *et al.*, 2013, 2017; Maier *et al.*, 2018; Welch *et al.*, 2018; Schindler *et al.*, 2020). The trace gas exchange capacity of trees and their contributions to ecosystem N_2O and CH_4 exchange seem, however, to vary considerably among tree individuals, tree species, forest ecosystem types and climatic zones, and to depend on many aspects, such as soil and site parameters, tree size, age and health conditions, environmental conditions and seasonal dynamics (Barba *et al.*, 2019; Covey & Megonigal, 2019; Machacova *et al.*, 2019). Even though the interactions between soil, vegetation and atmosphere exert a crucial role in controlling the ecosystem budget of N_2O and CH_4 , our current – still limited – knowledge on tree exchange of CH_4 , and especially of N_2O , does not allow us to clearly identify common characteristics, processes, pathways, and mechanisms of N_2O and CH_4 exchange in the soil–tree–atmosphere continuum, and to constrain the magnitudes and patterns of N_2O and CH_4 emissions.

Wetlands and floodplains are the largest natural sources of atmospheric CH_4 in the tropics (Saunois *et al.*, 2016). To date, calculations of N_2O and CH_4 fluxes between wetlands and the atmosphere have been based mostly upon GHG exchange at the soil–atmosphere interface only, thus excluding the exchange activity of such other ecosystem compartments as trees and other vegetation. This approach can lead to underestimating the ecosystem fluxes (Barba *et al.*, 2019). In general, trees growing in wetlands and floodplain forests seem to be stronger emitters of CH_4 than trees in upland forests (Pangala *et al.*, 2013, 2017; Machacova *et al.*, 2016a,b; Covey & Megonigal, 2019; Jeffrey *et al.*, 2019; Sjögersten *et al.*, 2020), whereas trees in riparian forests show emission values in between (Schindler *et al.*, 2020). The giants for their CH_4 emission potential are angiosperms in

tropical rain forests of the Amazon basin, with CH_4 emissions two to three orders of magnitude greater than those of trees growing in other tropical and temperate floodplain forests (Pangala *et al.*, 2017; Covey & Megonigal, 2019). Pangala *et al.* (2017) showed that these trees adapted to high soil water level are responsible for as much as half of the CH_4 emissions from the Amazon floodplain, which is the largest natural CH_4 source in the tropics.

With the exception of detailed, in-depth studies ongoing in the Amazon basin, which are today often used in estimating the overall CH_4 exchange of tropical forests, the widely distributed rain forests in other tropical continental and insular areas of Africa and Asia remain understudied. Likewise, mangroves and tropical forests on upland soils also are wholly understudied in relation to the CH_4 exchange of their woody vegetation (Pangala *et al.*, 2013; Welch *et al.*, 2018; Jeffrey *et al.*, 2019; Sjögersten *et al.*, 2020). Moreover, the N_2O exchange of mature trees growing under their natural field conditions is rarely investigated world-wide (Díaz-Pinés *et al.*, 2016; Machacova *et al.*, 2017, 2019; Wen *et al.*, 2017), and information for tropical regions, including tropical rain forests, is scarce (Welch *et al.*, 2018). Without ecologically relevant studies of tree and ecosystem exchange of CH_4 and N_2O covering the broad mosaic of tropical forest ecosystems, it is more than challenging to understand the role of woody plants in the GHG balance of tropical regions, to correctly estimate the tropical forest CH_4 and N_2O budgets, and to predict their future development in relation to global climate change.

Accordingly, the objective of our case study was to quantify the N_2O , CH_4 and, additionally, carbon dioxide (CO_2 , an indicator of physiological activity) exchange of dominant tree species in a tropical lowland rain forest on volcanic Réunion Island (southwestern Indian Ocean). We studied trace gas fluxes from two trees endemic for Réunion and Mauritius islands (*Doratoxylon apetalum*, *Antirhea borbonica*) and four regional endemics of Madagascar, Mauritius and Réunion (*Syzygium borbonicum*, *Homalium paniculatum*, *Mimusops balata* and *Labourdonnaisia calophylloides*). These measurements were accompanied by the investigation of GHG exchange from adjacent soil and volcanic surfaces (basaltic lava flows) and from widespread cryptogamic (bryophytic) stem covers (*Pyrrhobryum spiniforme*, *Leucoloma capillifolium*). The volcanic Réunion Island belongs to one of 36 world hotspots of biodiversity with high level of endemism (Kreft *et al.*, 2008), and its National Park, covering 42% of the island, provides an exceptional setup for scientific experimentations in primary vegetation along a major elevational gradient (3069 m above sea level (asl), Piton des Neiges) and with proximity to a regularly active volcano (2632 m asl, Piton de La Fournaise).

Still relatively young, the studied forest has developed on a 400-yr-old basaltic lava flow. Because young lava flows lack nutrients and organic matter, the microorganisms colonizing these newly created ecosystems fix not only N_2 , CO_2 and ammonium (NH_4^+), but also other trace gases from the atmosphere, such as carbon monoxide (CO), hydrogen (H_2) and CH_4 , and drive the sequestration of N, C and other nutrients needed for further ecosystem development (King, 2003; Gomez-Alvarez *et al.*, 2007;

Byloos *et al.*, 2018). The further ecological succession is connected to the accumulation of nutrients and leads to the gradual development of various plant communities starting with algae, mosses, lichens and ferns, followed by woody shrubs, shrubs growing to trees and later mature trees, forming a dense canopy forest *c.* 400 yr after the destruction of all past vegetation by new lava substrate formation (Potgieter *et al.*, 2014). This development is accompanied by the formation of a soil layer, with associated changes in soil properties and microbial community. In the case of the studied area, an unique native lowland rain forest has developed (Kirman *et al.*, 2007).

Even though the island has a tropical climate with high annual precipitation (>4000 mm), the studied forest is atypical compared to well-studied Amazon basin forests. It is specific for its porous volcanic bedrocks with the presence of lava holes and tubes, thin and irregular soil layer with weak water holding capacity, and steep slopes. It hosts high biodiversity, including mostly tree species endemic at the archipelago or regional levels. The aforementioned characteristics have resulted in an unique and atypical tropical rain forest ecosystem without standing water even under heavy rains. Therefore, we aimed to investigate: (1) whether – and, if so, to what extent – the tree stems growing on a lava flow exchange CH₄ and N₂O with the atmosphere; (2) how the tree fluxes contribute to the forest GHG exchange; and (3) whether the tropical rain forest is a source or sink for CH₄ and N₂O at the beginning of the rainy season.

Materials and Methods

Site description and study design

The experiment was conducted within the Mare Longue Nature Reserve (lat. –21°21′28.2024″N, long. 55°44′37.554″E), situated in the southeast of Réunion Island, a tropical volcanic island located in the southwestern Indian Ocean. The studied mixed forest is a tropical lowland rain forest characterized by endemic vegetation typical for the volcanic islands in the Mascarene Archipelago. The studied forest site (size 1 ha, established in 1999 by the University of Réunion Island) is located on sloping terrain at 180–200 m asl and situated on a pahoehoe basaltic lava flow *c.* 400 yr old (Kirman *et al.*, 2007). The soil cover is irregular and thin, and consists mostly of organic matter, parent-rock fragments and iron oxyhydroxides (Kirman, 2003; Meunier *et al.*, 2010). The thin A horizon (average 1 cm depth) is accompanied by thin and irregularly occurring eluvial deposits (weathered surface of the basaltic lava bedrocks; *i.e.* the C horizon). The chemical composition of the basalt (composed of feldspar (anorthite), olivine (forsterite) and pyroxene (augite)) and of the soil in the studied forest site can be found in Meunier *et al.* (2010). In the experimental plots, we distinguished soil-covered lava flow spots (further referred to as ‘soil’) from bare lava flow surfaces without soil coverage (further referred to as ‘basaltic lava flows’ or ‘volcanic rocks’). As a consequence of the roughness of the soil surface, the plant roots colonize deep volumes of lava bedrocks (Meunier *et al.*, 2010). The aboveground biomass and litterfall of the studied forest were measured in the previous study of Kirman

et al. (2007). The total standing biomass (535 t ha⁻¹) and annual litterfall (7.6 t ha⁻¹), as well the chemical composition of the biomass (stored major elements), are similar to other tropical rain forests (Kirman *et al.*, 2007; Meunier *et al.*, 2010).

The long-term mean annual precipitation of the southeast of Réunion Island is 4256 mm; the mean minimum and maximum temperatures are 19.9 and 26.4°C, respectively (data from Le Baril Météo-France station, 1981–2010). The study period (8 October–7 November 2018) was characterized by mean daily air and soil temperature (50 cm soil depth) of 20.6 ± 2.0°C and 19.3 ± 0.5°C (mean ± SD), respectively, and mean daily relative humidity of 97.4% ± 7.1%. All daily parameters were measured in the studied forest site using an RHTemp1000IS relative humidity and temperature data logger (MadgeTech, Warner, NH, USA), and Soil Matric Potential Sensors 253-L and 257-L (Campbell Scientific, Logan, UT, USA). The measurement period was a time of transition between a cool and dry season (May–October) and a warm and humid season (November–April).

Within the studied forest site area of 1 ha, we randomly selected 17 experimental plots (each 100 m²) out of 100 available experimental plots. The exchange of nitrous oxide (N₂O), methane (CH₄) and carbon dioxide (CO₂) from stems was studied on six dominant tree species (in total 24 mature trees representatively selected within these plots): *Syzygium borbonicum* J. Guého et A.J. Scott (*n* = 5); *Doratoxylon apetalum* (Poir.) Radlk. var. *apetalum* (*n* = 5); *Antirhea borbonica* J.F. Gmel (*n* = 5); *Homalium paniculatum* (Lam.) Benth. (*n* = 3); *Mimusops balata* (Aubl.) C.F. Gaertn. (*n* = 3); and *Labourdonnaisia calophylloides* Bojer (*n* = 3). The biometric parameters of the measured trees can be found in Table 1. Moreover, cryptogamic (bryophytic) stem covers (*i.e.* photoautotrophic organisms growing on tree bark) typical for the selected trees (*Pyrrhobryum spiniforme* (Hedw.) Mitt., *Leucoloma capillifolium* Renaud; *n* = 4) were collected for further trace gas flux measurements under laboratory conditions. The exchange of greenhouse gases (GHG) from the adjacent soil was measured close to each selected individual tree (total *n* = 24 soil positions). Finally, basaltic lava flows as volcanic surfaces without soil cover (*n* = 8) were studied for their N₂O, CH₄ and CO₂ exchange potential.

The measurements of N₂O, CH₄ and CO₂ fluxes from stems and soil were made in pairs, in the sense that the stem gas flux measurement for an individual tree directly followed the gas flux measurement of the adjacent soil to ensure measurements under similar environmental and climatic conditions. The measurements on cryptogams and basaltic lava flows were carried out in blocks. All fluxes were determined between 09:00 h and 18:00 h. One measurement set from all selected trees, cryptogams, soils and basaltic lava flows required *c.* 1.5–2 wk. All of the forest components were measured twice.

Gas sampling from stems, soil and basaltic lava flows

Stem fluxes of N₂O, CH₄ and CO₂ were measured at the bottom part of the stems (*c.* 0.4 m aboveground) for all selected tree species and tree individuals. The vertical profile of the stem fluxes

Table 1 Biometric parameters of studied trees and forest stand characteristics (mean \pm SD).

| Tree | DBH (m) | Tree height* (m) | Stem surface area (m ²) | Tree density** (trees ha ⁻¹) |
|--------|-----------------|------------------|-------------------------------------|--|
| Syzbor | 0.33 \pm 0.08 | 18.8 \pm 4.4 | 10.2 \pm 5.0 | 30 |
| Dorape | 0.25 \pm 0.05 | 13.9 \pm 2.7 | 5.5 \pm 2.2 | 91 |
| Antbor | 0.16 \pm 0.02 | 13.1 \pm 1.9 | 3.4 \pm 1.0 | 86 |
| Hompan | 0.43 \pm 0.23 | 16.0 \pm 7.8 | 12.8 \pm 12.6 | 42 |
| Mimbal | 0.42 \pm 0.17 | 23.7 \pm 9.4 | 17.3 \pm 14.1 | 65 |
| Labcal | 0.54 \pm 0.21 | 28.5 \pm 10.6 | 26.2 \pm 19.3 | 58 |

Syzbor, *Syzygium borbonicum* ($n = 5$); Dorape, *Doratoxylon apetalum* ($n = 5$); Antbor, *Antirhea borbonica* ($n = 5$); Hompan, *Homalium paniculatum* ($n = 3$); Mimbal, *Mimusops balata* ($n = 3$); Labcal, *Labourdonnaisia calophylloides* ($n = 3$); DBH, stem diameter at breast height.

*Tree height of studied trees was estimated using allometric relationships for Mare Longue Nature Reserve kindly provided by Olivier Flores (University of Réunion Island).

**Tree density is calculated for each tree species separately and for trees with DBH > 0.10 m to include only trees relevant for flux measurements. The tree density of all six tree species studied is 372 trees ha⁻¹. The overall forest density, including all c. 80 tree species present, is estimated to be 2150 trees ha⁻¹ (Kirman *et al.*, 2007).

(measurements at three stem heights of c. 0.4, 1.1 and 1.8 m aboveground; Fig. 1a–c) was studied in three individual trees of each of the following three species: *S. borbonicum*, *D. apetalum* and *A. borbonica*. The stem fluxes were measured using static stem chamber systems installed at the beginning of the measurement campaign. The chambers consisted of transparent plastic containers with removable airtight lids (Lock & Lock, Seoul, South Korea) and a neoprene sealing frame. They were gas-tightly affixed to the carefully smoothed bark surface c. 2 wk before measurements. Control measurements were performed to ensure that the observed fluxes did not originate from the chamber materials used. On each tree other than *S. borbonicum* individuals, two chambers were installed at one stem height on opposite sides of the stem and interconnected with polyurethane tubes into a single flow-through chamber system. The total enclosed stem area was 0.0108 m² and total internal system volume was 0.0021 m³. (For more details, see Machacova *et al.*, 2015, 2017, 2019.) Owing to the large stem diameter of *S. borbonicum*, three chambers (total area 0.0162 m², total internal volume 0.0028 m³) were installed at one stem height in the same manner on those trees. Gas-tightness of all the chambers was tested regularly using CO₂ signal and a portable gas analyzer (see the 'Flux measurements and calculations' section below).

Soil N₂O, CH₄ and CO₂ fluxes were measured using static soil chamber systems (Fig. 1b,d). The soil collars were installed in the vicinity of the investigated trees at the beginning of the research stay to reduce soil disturbances. As a result of the shallow soil layer on the lava flow, the bottom part of the collars was imbedded to a maximum 2.5 cm deep into the soil. The soil chambers were made of sewage pipes (total enclosed soil area 0.0083 m², total internal system volume c. 0.0015 m³ depending on soil depth). During measurements, the soil chambers were closed by

a lid and sealed with water between the body of the chamber and the lid (Machacova *et al.*, 2013).

The fluxes of N₂O, CH₄ and CO₂ from basaltic lava flows were determined using the stem chambers as described above but installed directly onto the volcanic surfaces (Fig. 1c). One flow-through chamber system consisted of one chamber (total enclosed rock area 0.0054 m², total internal system volume 0.0013 m³) connected to a gas analyzer (see the 'Flux measurements and calculations' section below).

Collection and incubation of cryptogamic stem covers

Pyrrhobryum spiniforme and *Leucoloma capillifolium* mosses were collected from the bark of *S. borbonicum* trees. To avoid disruption to the bark microcosm within the stem chambers, the cryptogams were collected from bark outside the measurement chambers. After collection, fresh samples were placed into plastic gas-tight incubation chamber (0.0039 m³ volume; Lock & Lock; Fig. 1e,f) connected to a gas analyzer and incubated in the laboratory under dark conditions and low light intensity of 10–15 $\mu\text{mol m}^{-2} \text{s}^{-1}$ corresponding to the low light conditions of forest understories.

The stem, soil, lava flow and incubation chamber systems were left open during the time between the individual measurements.

Flux measurements and calculations

For GHG flux measurements, the measurement chambers were closed with lids, and the concentration changes of N₂O, CH₄ and CO₂ in chamber headspace were measured using a Gasmeter DX-4015 portable Fourier transform infrared (FTIR) gas analyzer (Gasmeter Technologies Oy, Vantaa, Finland; Warlo *et al.*, 2018). A single measurement run of stem, soil and lava flow gas fluxes lasted c. 45 min. The measurement time for cryptogams was 180–240 min, depending on flux rates. The internal pump (0.002 m³ min⁻¹) of the analyzer ensured mixing of air inside the chamber systems. Every morning, zero-point calibration of the analyzer was made using N₂ (99.9992% purity).

The gas exchanges of stems, soil, lava flows and cryptogams were quantified based upon the linear changes in N₂O, CH₄ and CO₂ concentrations in the chamber headspace over time (for examples, see Supporting Information Fig. S1; for equations used, see Machacova *et al.*, 2016a). Decrease of gas concentration over time indicated gas uptake (i.e. negative flux), increase of gas concentration indicated gas emission (i.e. positive flux). The fluxes from tree stems and soil were further roughly scaled up to the ecosystem level based on tree and forest characteristics (stem diameter at breast height (DBH), tree height, tree density for individual tree species with trees of DBH > 0.10 m; Table 1; Machacova *et al.*, 2016a). The scaling up of trace gas exchange of volcanic surfaces of lava flows was not possible because the exact extent of volcanic bedrock surfaces without soil cover within the studied forest was unknown. The contribution of cryptogams to the stem GHG fluxes was estimated based on upscaling of the fluxes related to g DW (dry weight) to the unit of stem surface area (Machacova *et al.*, 2017).




Fig. 1 View of study set-up and chamber systems used to determine greenhouse gas fluxes in the lowland tropical rain forest on volcanic Réunion Island. Overall view of the site: stem chambers installed in the vertical profile of tree stems (a–c), soil chambers in the vicinity of studied trees (a, b; marked with white circle; and d), volcanic rock chambers fixed on basaltic lava flow (c, marked with white asterisk), cryptogamic stem covers in incubation chambers (e, f), and portable greenhouse gas analyzer used for gas concentration measurements (a, b).

Statistics

The flux data were checked for normal distribution (Shapiro–Wilk test) and equality of variances in the different subpopulations. Student's *t*-test and one-way ANOVA for multiple comparisons were applied for normally distributed data. The nonparametric Mann–Whitney rank-sum test and Kruskal–Wallis one-way ANOVA on ranks for multiple comparisons were applied for non-normally distributed data or data with unequal variances. The exact applied tests and *n* values for statistical analyses are stated in the figure legends and in the 'Site description and study design' section above, respectively. The study was performed in 17 randomly selected experimental plots within the studied 1 ha forest site, and the studied trees together with the soil positions and basaltic lava flow surfaces were representatively chosen within these plots. Statistical significance for all tests was defined as $P < 0.05$. The statistics were run using SIGMAPLOT 11.0 (Systat Software, San Jose, CA, USA).

Results

Methane exchange of forest compartments

The soil of the tropical lowland rain forest consistently consumed CH_4 from the atmosphere ($-83.3 \pm 9.8 \mu\text{g-CH}_4 \text{ m}^{-2} \text{ h}^{-1}$ (soil area), mean \pm SE) at the beginning of the rainy season. Uniform CH_4 consumption was observed at all subsites with low, nonsignificant spatial flux heterogeneity (Fig. 2a). The volcanic surfaces of basaltic lava flows also showed consistent uptake of CH_4 from the atmosphere ($-107 \pm 21.6 \mu\text{g-CH}_4 \text{ m}^{-2} \text{ h}^{-1}$ (rock area), Fig. 2a). As the similar consumption rates of irregularly spread soil layers and of volcanic rocks are expressed per 'ground' area, and both surface types seem to be equally distributed in the studied forest, we can confidently say that the detected CH_4 consumption of both forest compartments equally and significantly contributes to CH_4 exchange of the forest ground area (Fig. 2a). The




Fig. 2 Fluxes of methane (CH_4) (a), nitrous oxide (N_2O) (b), and carbon dioxide (CO_2) (c) from adjacent soil measured close to each studied tree and from volcanic rock surfaces of basaltic lava flows. The fluxes are expressed as medians (solid lines) and means (dashed lines) of measurements from soil positions in the vicinity of trees of each individual tree species (Syzbor, *Syzygium borbonicum* ($n=5$); Dorape, *Doratoxylon apetalum* ($n=5$); Antbor, *Antirhea borbonica* ($n=5$); Hompan, *Homalium paniculatum* ($n=3$); Mimbal, *Mimusops balata* ($n=3$); Labcal, *Labourdonnaisia calophylloides* ($n=3$)), from all studied soil positions ('All soil positions', $n=24$), and from volcanic rocks ($n=8$). The forest floor in the studied forest is approximately equally covered with soil and volcanic surfaces without soil cover (proportion roughly 50 : 50). Fluxes are expressed per m^2 of soil and volcanic rock area. Positive fluxes indicate trace gas emission; and negative fluxes trace gas uptake. The box boundaries mark the 25th and 75th percentiles and whiskers the 10th and 90th percentiles. Dots mark outliers. As indicated by the letter 'a' above each bar, there were no statistically significant differences among fluxes in adjacent soil of individual tree species at $P < 0.05$. One-way ANOVA was applied for CH_4 fluxes and Kruskal–Wallis one-way ANOVA on ranks was used for N_2O and CO_2 fluxes.

forest floor CH_4 fluxes can therefore be estimated using 50:50 proportion of soil and volcanic rock coverage.

All mature studied tree stems were net sinks of CH_4 from the atmosphere ($-15.6 \pm 2.0 \mu\text{g-CH}_4 \text{ m}^{-2} \text{ h}^{-1}$ (stem area)). The CH_4 consumption was uniform across the various tree species and did not show significant species-specific variability (Fig. 3a). Moreover, no significant changes in stem CH_4 consumption were observed with respect to stem height aboveground (Fig. 4a). The uptake of CH_4 by tree stems was not closely connected to stem CO_2 efflux (Fig. 5a). To better understand the mechanisms behind the observed CH_4 uptake by trees, the GHG exchange of widespread cryptogamic stem cover was measured. We estimated how much CH_4 is taken up by the active cryptogams per stem area unit when fully covered with these organisms. As a result, cryptogams were identified as net sinks of CH_4 ($-8.3 \pm 3.0 \mu\text{g-CH}_4 \text{ m}^{-2} \text{ h}^{-1}$ (stem area); $-0.047 \pm 0.016 \mu\text{g g}^{-1} \text{ h}^{-1}$ (DW); Fig. 3a). All main forest compartments studied therefore consumed CH_4 from the atmosphere.

In order to estimate the contribution of the tree stems to the tropical rain forest CH_4 exchange, the detected stem and soil

consumption rates were scaled up to ecosystem level (Fig. 6a,b). The six tree species studied (joint tree density of $c. 370$ trees ha^{-1}) consumed in total $-59.0 \pm 17.9 \text{ mg-CH}_4 \text{ ha}^{-1} \text{ h}^{-1}$ (ground area) (i.e. the sum of mean tree species consumption rates and SEs) and contributed 7.1% to the soil CH_4 uptake ($-833.1 \pm 97.5 \text{ mg-CH}_4 \text{ ha}^{-1} \text{ h}^{-1}$; Fig. 6b). As there are > 80 tree species present in the studied forest (total tree density of $c. 2150$ trees ha^{-1} ; Kirman *et al.*, 2007), we estimate that the overall contribution of all tree stems present to the soil CH_4 uptake might be 41% when assuming similar CH_4 uptake potential by all tree species present. This tropical rain forest with its > 80 tree species seems to be a strong sink for CH_4 at the beginning of the rainy season ($-1176 \text{ mg-CH}_4 \text{ ha}^{-1} \text{ h}^{-1}$).

Nitrous oxide exchange of forest compartments

The soil was on average a weak source of N_2O ($1.8 \pm 2.5 \mu\text{g-N}_2\text{O m}^{-2} \text{ h}^{-1}$; Fig. 2b). The detected exchange potential of volcanic surfaces of basaltic lava flows was negligible ($0.024 \pm 0.871 \mu\text{g-N}_2\text{O m}^{-2} \text{ h}^{-1}$). The exchange of soil and




Fig. 3 Fluxes of methane (CH_4) (a), nitrous oxide (N_2O) (b), and carbon dioxide (CO_2) (c) from tree stems and cryptogamic stem covers. The fluxes are expressed as medians (solid lines) and means (dashed lines) of measurements from stems of six individual tree species (Syzbor, *Syzygium borbonicum* ($n = 5$); Dorape, *Doratoxylon apetalum* ($n = 5$); Antbor, *Antirhea borbonica* ($n = 5$); Hompan, *Homalium paniculatum* ($n = 3$); Mimbal, *Mimusops balata* ($n = 3$); Labcal, *Labourdonnaisia calophylloides* ($n = 3$)), from all studied trees and tree species ('All trees', $n = 24$), and from cryptogams (*Pyrrhobryum spiniforme*, *Leucocoloma capillifolium*; $n = 4$). The trace gas exchange of cryptogams is presented as mean and median of all measurements under low light and dark conditions, because CH_4 and N_2O fluxes did not differ in relation to light conditions. The mean and median fluxes of CO_2 in cryptogams include both CO_2 emission and uptake measured under dark and light conditions, resulting in low gas exchange (more details in manuscript text). All fluxes, including fluxes from cryptogams, are expressed per m^2 of stem area. Positive fluxes indicate trace gas emission and negative fluxes trace gas uptake. The box boundaries mark the 25th and 75th percentiles and whiskers the 10th and 90th percentiles. Dots mark outliers. Statistically significant differences among fluxes in individual tree species at $P < 0.05$ are indicated by different letters above bars. One-way ANOVA was applied for CH_4 and N_2O fluxes and Kruskal–Wallis one-way ANOVA on ranks was used for CO_2 fluxes.

volcanic surfaces showed high spatial heterogeneity without any clear trend, including both N_2O consumption and emission. Moreover, the majority of the exchange rates were very low.

In contrast, clear uptake of N_2O by tree stems was observed. Thus, these are net sinks of N_2O from the atmosphere ($-3.0 \pm 0.8 \mu\text{g-N}_2\text{O m}^{-2} \text{h}^{-1}$; Fig. 3b). The lowest N_2O consumption was detected for *Syzygium borbonicum* and the highest for *Homalium paniculatum*. The uptake of N_2O by tree stems was negatively correlated with stem CO_2 efflux (Fig. 5b). Similar to the situation for CH_4 , no significant changes in stem N_2O exchange were observed within the vertical stem profile (Fig. 4b). The cryptogams were a clear sink not only of CH_4 , but also of N_2O ($-1.9 \pm 0.7 \mu\text{g-N}_2\text{O m}^{-2} \text{h}^{-1}$ (stem area); $-0.011 \pm 0.004 \mu\text{g g}^{-1} \text{h}^{-1}$ (DW); Fig. 3b). Our test measurements confirmed that the observed cryptogamic N_2O and CH_4 uptake could not be explained by gas dilution in air humidity within the incubation chambers (Machacova *et al.*, 2017).

The scaled-up rates of N_2O consumption by tree stems of the six dominant tree species studied ($-11.9 \pm 6.3 \text{ mg-N}_2\text{O ha}^{-1} \text{h}^{-1}$) represented -64% of the soil N_2O emissions ($18.4 \pm 25.2 \text{ mg-N}_2\text{O ha}^{-1} \text{h}^{-1}$; Fig. 6c,d), thus decreasing the weak source strength of the soil. (Please note that the usage of the negative percentage in case of N_2O is due to comparison of bidirectional fluxes – negative fluxes (i.e. N_2O uptake by stems) with positive fluxes (i.e. N_2O emission by soil; equal to 100%). In the case of CH_4 , the percentage contribution of the stem fluxes to the soil fluxes is positive, as both – tree stems and soil – are sinks for CH_4 .) Under the assumption of all 80 tree species in the forest being N_2O consumers at similar uptake rates, however, we expect that the overall N_2O uptake potential of all trees ($-68.9 \text{ mg-N}_2\text{O ha}^{-1} \text{h}^{-1}$) might represent even -374% of the soil N_2O exchange, thus turning the tropical rain forest into a sink of N_2O ($-50.45 \text{ mg-N}_2\text{O ha}^{-1} \text{h}^{-1}$).




Fig. 4 Fluxes of methane (CH_4) (a), nitrous oxide (N_2O) (b), and carbon dioxide (CO_2) (c) from tree stem vertical profiles. The fluxes are expressed as medians (solid lines) and means (dashed lines) of measurements from trees of three tree species: *Syzygium borbonicum* ($n = 3$), *Doratoxylon apetalum* ($n = 3$) and *Antirhea borbonica* ($n = 3$). The measurements were performed at three stem heights of c. 0.4, 1.1 and 1.8 m aboveground. All fluxes are expressed per m^2 of stem area. Positive fluxes indicate trace gas emission and negative fluxes trace gas uptake. The box boundaries mark the 25th and 75th percentiles and whiskers the 10th and 90th percentiles. As indicated by the letter 'a' above each bar, there were no statistically significant differences in fluxes among stem heights at $P < 0.05$. One-way ANOVA was applied for CH_4 and N_2O fluxes and Kruskal–Wallis one-way ANOVA on ranks was used for CO_2 fluxes.

Carbon dioxide exchange of forest compartments

The measured CO_2 exchange of forest compartments is an ancillary parameter (i.e. an indicator of physiological activity) helping to understand the CH_4 and N_2O exchange in the soil–tree–ecosystem–atmosphere continuum. The CO_2 emissions from soil substantially exceeded those from volcanic surfaces of lava flows (650 ± 95 and 35.5 ± 4.7 $\text{mg-CO}_2 \text{ m}^{-2} \text{ h}^{-1}$, respectively; Fig. 2c). Tree stems emitted 295 ± 73 $\text{mg-CO}_2 \text{ m}^{-2} \text{ h}^{-1}$, with *Homalium paniculatum* being the strongest emitter (Fig. 3c). The stem CO_2 efflux was uniform across the vertical stem profile (Fig. 4c). As expected, photoautotrophic cryptogamic stem covers consumed CO_2 under light conditions (-15.6 ± 2.5 $\text{mg-CO}_2 \text{ m}^{-2} \text{ h}^{-1}$ (stem area); -0.094 ± 0.027 $\text{mg g}^{-1} \text{ h}^{-1}$ (DW)) and emitted CO_2 under dark conditions (14.8 ± 2.9 $\text{mg-CO}_2 \text{ m}^{-2} \text{ h}^{-1}$ (stem area); 0.087 ± 0.010 $\text{mg g}^{-1} \text{ h}^{-1}$ (DW); NB Fig. 3c shows total mean and median values from both incubation conditions).

Discussion

All studied forest compartments are net methane sinks

Soil and basaltic lava flows Soil (i.e. soil on soil-covered lava flow spots) and basaltic lava flows (i.e. bare lava flow surfaces




Fig. 5 Relationships between methane (CH_4) vs carbon dioxide (CO_2) stem fluxes (a) and nitrous oxide (N_2O) vs CO_2 stem fluxes (b). All six tree species (24 trees in total) are included. All fluxes are expressed per m^2 of stem area. Positive flux values indicate gas emission and negative values indicate gas uptake.

without soil coverage) consistently consumed methane (CH_4) from the atmosphere, even though the measurements were done at the start of the rainy season. The forest floor is covered by irregular and thin soil (Kirman *et al.*, 2007), likely preventing the formation and preservation of anaerobic soil conditions for methanogenic CH_4 production and thus supporting methanotrophic CH_4 oxidation (Smith *et al.*, 2003). The steep slope further promotes rapid rainwater runoff from the forest into the ocean. Moreover, the high porosity typical for the volcanics forming the underlying rocks of the forests and their typical irregular distribution with the presence of hollows and lava tubes also contributes to rapid rainwater runoff and oxygenation of the whole system.

To the best of our knowledge, the ability of basaltic lava flows to exchange CH_4 with the atmosphere has never been reported. We speculate that the CH_4 oxidation occurs in volcanic pores covered with a thin layer of biofilm consisting of a broad spectrum of microbial communities, including (*inter alia*) methanotrophs (e.g. Gomez-Alvarez *et al.*, 2007; Byloos *et al.*, 2018). Methanotrophs seem to colonize the newly formed lava already within 40 yr, but the colonization is closely related to the development of plant and microbial communities and local climatic conditions (King & Nanba, 2008). We expect that volcanic surfaces' low water absorption and retention capacity (Gomez-Alvarez *et al.*, 2007; Byloos *et al.*, 2018) promote aerobic conditions and support, *inter alia*, CH_4 -oxidizing microorganisms leading to the observed CH_4 uptake by volcanic surfaces.

Soil CH_4 flux measurements on lava flows are rare. The only study conducted on relatively fresh volcanic deposits detected soil CH_4 uptake in a tropical rain forest (c. 300 yr old) on the islands of Hawaii (King, 2003). The CH_4 consumption rate in that study (reaching -75 $\mu\text{g-CH}_4 \text{ m}^{-2} \text{ h}^{-1}$) is in accordance with the findings

of our study ($-83.3 \mu\text{g-CH}_4 \text{ m}^{-2} \text{ h}^{-1}$). The similar CH_4 consumption potentials of soil and volcanic surfaces were accompanied by soil carbon dioxide (CO_2) emissions substantially exceeding those from volcanic surfaces (Fig. 2c). By comparison, the volcanic deposits in Hawaii produced twice as much CO_2 ($58.8 \text{ mg-CO}_2 \text{ m}^{-2} \text{ h}^{-1}$; King, 2003). The very low CO_2 emissions of volcanic surfaces in Réunion might be explained by the low availability of organic carbon (C) and resulting low respiration rates (King, 2003).

Tree stems and cryptogamic stem covers Our study clearly determined all mature tree stems of the dominant tree species to be net sinks of CH_4 (Fig. 3a). To the best of our knowledge, such consistent consumption of CH_4 by tree stems constitutes an unique finding, because trees are generally known as CH_4 sources with high individual variability in CH_4 emission rates, depending on tree species, climatic zone, forest ecosystem type, environmental and meteorological conditions, and seasonal dynamics (Barba *et al.*, 2019; Covey & Megonigal, 2019). In particular, stems of Amazonia trees that are well adapted to inundation are regarded as extraordinarily strong CH_4 sources (Pangala *et al.*, 2017). Trees from other tropical areas seem also to play important roles in forest CH_4 budgets, even with much lower detected emissions than in Amazonia (Pangala *et al.*, 2013; Welch *et al.*, 2018; Jeffrey *et al.*, 2019; Sjögersten *et al.*, 2020). It is expected that CH_4 emitted from these trees mostly originates from the deeper soil layers, as reported for the only study done on tropical upland soils (Welch *et al.*, 2018). In that case, trees were observed to serve as ‘chimneys’ for soil-produced CH_4 to reach the atmosphere without being oxidized in well-aerated soil surface layers.

Only boreal tree species are known to be capable of taking up CH_4 from the atmosphere by their leaves and stems (Sundqvist *et al.*, 2012; Machacova *et al.*, 2016b), although the stem uptake potential was detected only in the dormant season and was rather negligible (Machacova *et al.*, 2016b). Welch *et al.* (2018) detected irregular CH_4 uptake by stems of *Simarouba amara* in a tropical upland forest in Panama during the transition from dry to wet season. Those authors assumed a diffusion gradient from the atmosphere via tree stems into the soil with predominant CH_4 oxidation, which might result in tree stem uptake of trace gases.

Our incubation experiments with the most abundant cryptogams on studied trees consistently revealed that all freshly collected cryptogams (bryophytes) were net CH_4 sinks independently of light conditions (Fig. 3a). The estimated CH_4 uptake rates per area unit were of the same order of magnitude as stem uptake rates measured under field conditions (Fig. 3a). Therefore, it seems that the epiphytes are co-responsible for the observed CH_4 uptake by tropical trees on Réunion. The only known study on cryptogamic CH_4 exchange presents various cryptogamic species as small CH_4 sources ($0.28 \text{ ng-CH}_4 \text{ g}^{-1} \text{ h}^{-1}$, Lenhart *et al.*, 2015; vs $-47 \pm 16 \text{ ng-CH}_4 \text{ g}^{-1} \text{ h}^{-1}$, this study). We hypothesize that CH_4 -oxidising microorganisms (i.e. methanotrophs) can be involved in the CH_4 consumption observed by trees and cryptogams. Future analyses of wood and cryptogams samples for the *pmoA* gene controlling CH_4 oxidation could bring more explanation to this topic.

In conclusion, the studied tropical rain forest located on a basaltic lava flow seems to be a strong net sink of CH_4 , with soil

and volcanic surfaces predominantly being responsible for the CH_4 uptake. We dare to say that, owing to the reasons stated above, no elevated water table levels should be expected during the rainy season, and the studied tropical rain forest on lava flow might be a substantial sink for CH_4 all through the year. The stems of the six tree species studied contributed 7.1% to the forest floor CH_4 uptake (Fig. 6b). As there are > 80 tree species present with high tree density, we estimate that the overall contribution of all tree stems to the forest floor CH_4 uptake might be as much as 41% and the overall forest uptake potential can reach to $-1176 \text{ mg-CH}_4 \text{ ha}^{-1} \text{ h}^{-1}$. The trees growing in the tropical rain forest on lava flows, therefore, are significant contributors to ecosystem CH_4 uptake. Theoretically, the contribution can be even greater, as foliage, known from the first available studies as important contributors to ecosystem CH_4 exchange (Machacova *et al.*, 2016a), is not included in this case study.

Tree stems and cryptogamic stem covers are net nitrous oxide sinks

By contrast with their relationship to CH_4 , soil and basaltic lava flows do not significantly exchange nitrous oxide (N_2O) with the atmosphere (Fig. 2b). The reason might be nitrogen (N) limitation in the thin soil layer, thus resulting in very limited biomass increment in the studied forest over the past 10 yr (O. Flores, pers. comm.; Bennici, 2015). Moreover, the young lava flow is probably still poor in N-rich nutrients and organic compounds (King, 2003; Gomez-Alvarez *et al.*, 2007).

Nevertheless, the studied trees were identified as net sinks of N_2O (Fig. 3b). Their uptake potential was weakly and negatively correlated with stem CO_2 efflux (Fig. 5b), which means that low N_2O consumption by tree stems was associated with low stem CO_2 efflux. The net stem CO_2 efflux, an indicator of tree physiological activity, is a result of stem respiration, radial CO_2 diffusion from the transpiration stream and CO_2 re-fixation on the stem surface (Aubrey & Teskey, 2009; Hölttä & Kolari, 2009; Bloemen *et al.*, 2013). A physiological dependence of N_2O exchange has been observed in boreal trees, whose seasonality in stem N_2O release followed the tree physiological activity, particularly processes of CO_2 uptake and release (Machacova *et al.*, 2019). A closer connection between plant N_2O and CO_2 fluxes was detected early under a wide range of controlled environmental conditions in species belonging to cryptogamic covers (Lenhart *et al.*, 2015; Machacova *et al.*, 2017) and Spermatophyta (Machacova *et al.*, 2017; Lenhart *et al.*, 2019).

Similar to tree stems, cryptogams covering the studied tree stems up to the crowns also were detected as net N_2O sinks, and their N_2O uptake rates were of the same order of magnitude (Fig. 3b). These findings are unique, as the majority of studies on mature trees under natural field conditions present trees only as weak N_2O emitters (Machacova *et al.*, 2019). The only known study presenting trees (European beech, temperate mountain forest) as substantial net sinks of N_2O showed cryptogamic stem covers to be co-responsible for the observed N_2O uptake by tree stems (Machacova *et al.*, 2017). We hypothesize that the N_2O , which is taken up by the cryptogams, might be reduced to N_2 in




Fig. 6 Fluxes of methane (CH_4) (a, b), nitrous oxide (N_2O) (c, d) and carbon dioxide (CO_2) (e, f) from tree stems and adjacent soil expressed per stem or soil surface area unit (a, c, e) and scaled up to unit ground area of the tropical rain forest (b, d, f). The fluxes are expressed as medians (solid lines) and means (dashed lines) of measurements from all studied trees of six tree species ($n = 24$) and all studied soil positions ($n = 24$). Positive fluxes indicate trace gas emission; negative fluxes trace gas uptake. The box boundaries mark the 25th and 75th percentiles and whiskers the 10th and 90th percentiles. Statistically significant differences in fluxes at tree stem and soil level at $P < 0.05$ are indicated by different letters above the bars. Mann–Whitney rank-sum test was applied for the gas flux pairs. The contributions of stem fluxes of six tree species to the soil fluxes (equal to 100%) are expressed as percentages of the soil flux.

the final step of the denitrification pathway. Some irregular N_2O uptake by trees also has been observed by boreal trees (Machacova *et al.*, 2019) and tropical upland trees (Welch *et al.*, 2018). In general, trees seem to exchange N_2O (and CH_4) with the atmosphere in both directions under certain conditions. However, the processes and mechanisms behind and fate of the N_2O (and CH_4) taken up by the tree stems and cryptogams are still unknown and warrant further investigation.

The studied tree species contributed -64% to soil N_2O exchange and, therefore, reduced the weak soil N_2O emission potential (Fig. 6d). We expect an overall N_2O uptake potential of all tree species in the forest might represent as much as -374% of the soil exchange, thus turning the tropical rain forest into a strong sink of N_2O ($-50.45 \text{ mg-N}_2\text{O ha}^{-1} \text{ h}^{-1}$).

Conclusion and future perspectives

This study shows for the first time that native trees in tropical lowland rain forest growing on a basaltic lava flow can be important sinks of atmospheric CH_4 and N_2O and can contribute as much as 41% and -374% to the soil CH_4 and N_2O exchange, respectively. Also, cryptogamic stem covers were identified as substantial CH_4 and N_2O sinks. This ‘plant’ uptake was accompanied by strong CH_4 uptake by soil and volcanic surfaces of the basaltic lava flow.

In relation to ecosystem CH_4 exchange, the lowland rain forests growing on lava flows seem to behave differently from more commonly studied rain forests, not acting as strong CH_4 sources but rather as significant CH_4 sinks. More studies in different tropical

volcanic regions are needed to understand the trace gas exchange potential and mechanisms of these specific ecosystems.

Because our case study was limited by time, the results need to be extended by examining the seasonal variability in CH₄ and N₂O exchange for a larger set of tree species, further including fluxes from leaves, and simultaneous measurements of soil and plant chemical and hydrological properties following the previous work of Meunier *et al.* (2010), and seeking to understand the role of cryptogams and basaltic lava flows in the CH₄ and N₂O exchange under different climatic conditions.

Our results highlight the importance of previously unexplored lowland tropical rain forests on basaltic lava flows in CH₄ and N₂O uptake and, therefore, can be used to improve national greenhouse gas (GHG) inventories. As the studied lowland tropical rain forest in Mare Longue Nature Reserve falls within the range of tropical rain forests (biogeochemistry, biomass increment, litterfall; Kirman *et al.*, 2007; Meunier *et al.*, 2010), this study can be justified as a model ecosystem of lowland tropical rain forests. These forests are the most species-rich ecosystems in the world and are of high importance in regulating C and water cycling in the global climate system. However, they are under the greatest pressures from anthropogenic disturbances (acceleration of deforestation of original lowland tropical rain forests, impacts of global warming and drought events) and start to occur only as isolated forest fragments with decreasing biodiversity (Turner & Corlett, 1996; Corlett, 2011; Corlett & Primack, 2011). Confirming the unequivocal CH₄ and N₂O uptake potential by future studies would bolster the argument that the protection of such dynamic ecosystems in volcanic regions around the world should be strengthened not only because of their great biodiversity but also for their contribution to reducing GHGs in the atmosphere.

Acknowledgements





This research was supported by the Czech Science Foundation (17-18112Y), transnational Environmental Research Infrastructure (ENVRplus) access supported by the European Commission under the Horizon 2020 – Research and Innovation Action (H2020-INFRADEV-1-2014-1, Grant Agreement: 654182), SustES – Adaptation strategies for sustainable ecosystem services and food security under adverse environmental conditions (CZ.02.1.01/0.0/0.0/16_019/0000797). Also, financial support was provided by the Estonian Research Council (PRG-352 and MOBERC20) and by the EU through the European Regional Development Fund (EcolChange Centre of Excellence). Logistical support was provided by the Feder project (DIVINES), Réunion National Park, Office National des Forêts and OSU-Réunion. We thank Olivier Flores for providing allometric relationships in Mare Longue, as well as Yoan Benoit, Pierre Staménoff and Leszek Darjusz Laptaszynski for technical and organizational support.

Author contributions

KM had the idea for the study; KM and CA designed the study; KM, TA and LB carried out the field measurements and analyzed

the data; and KM, CA, TS, ÜM, KS and LB contributed to writing the manuscript.

ORCID

Thomas Agyei  <https://orcid.org/0000-0002-4606-3117>
Katerina Machacova  <https://orcid.org/0000-0002-8289-169X>
Ülo Mander  <https://orcid.org/0000-0003-2340-6989>
Kaido Soosaar  <https://orcid.org/0000-0002-1261-6326>

Data availability

The datasets generated and analyzed during this study are available from the corresponding author upon reasonable request.

References

- Aubrey DP, Teskey RO. 2009. Root-derived CO₂ efflux via xylem stream rivals soil CO₂ efflux. *New Phytologist* **184**: 35–40.
- Barba J, Bradford MA, Brewer PE, Bruhn D, Covey K, van Haren J, Megonigal JP, Mikkelsen TN, Pangala SR, Pihlatie M *et al.* 2019. Methane emissions from tree stems: a new frontier in the global carbon cycle. *New Phytologist* **222**: 18–28.
- Bennici A. 2015. *Modélisation de la dynamique d'une forêt tropicale humide insulaire: Mare-Longue, La Réunion*. MSc thesis, Université de la Réunion, Réunion, France.
- Bloemen J, McGuire MA, Aubrey DP, Teskey RO, Steppe K. 2013. Transport of root-respired CO₂ via the transpiration stream affects aboveground carbon assimilation and CO₂ efflux in trees. *New Phytologist* **197**: 555–565.
- Byloos B, Monsieurs P, Mysara M, Leys N, Boon N, Van Houdt R. 2018. Characterization of the bacterial communities on recent Icelandic volcanic deposits of different ages. *BMC Microbiology* **18**: 122.
- Corlett RT. 2011. Impacts of warming on tropical lowland rainforests. *Trends in Ecology and Evolution* **26**: 606–613.
- Corlett RT, Primack RB. 2011. *Tropical rain forests: an ecological and biogeographical comparison*. Hoboken, NJ, USA: Wiley-Blackwell.
- Covey KR, Megonigal JP. 2019. Methane production and emissions in trees and forests. *New Phytologist* **222**: 35–51.
- D'Annunzio R, Lindquist EJ, MacDicken KG, eds. 2017. *Global forest land-use change from 1990 to 2010: an update to a global remote sensing survey of forests. Forest Resource Assessment Working Paper 187*. Rome, Italy: Food and Agriculture Organization of the United Nations.
- Dalal RC, Allen DE. 2008. Turner review no. 18: Greenhouse gas fluxes from natural ecosystems. *Australian Journal of Botany* **56**: 369–407.
- Díaz-Pinés E, Heras P, Gasche R, Rubio A, Rennenberg H, Butterbach-Bahl K, Kiese R. 2016. Nitrous oxide emissions from stems of ash (*Fraxinus angustifolia* Vahl) and European beech (*Fagus sylvatica* L.). *Plant and Soil* **398**: 35–45.
- Espenberg M, Truu M, Mander Ü, Kasak K, Nõlvak H, Ligi T, Oopkaup K, Maddison M, Truu J. 2018. Differences in microbial community structure and nitrogen cycling in natural and drained tropical peatland soils. *Scientific Reports* **8**: 4742.
- Gomez-Alvarez V, King GM, Nüsslein K. 2007. Comparative bacterial diversity in recent Hawaiian volcanic deposits of different ages. *FEMS Microbiology Ecology* **60**: 60–73.
- Hölttä T, Kolari P. 2009. Interpretation of stem CO₂ efflux measurements. *Tree Physiology* **29**: 1447–1456.
- Jeffrey LC, Reithmaier G, Sippo JZ, Johnston SG, Tait DR, Harada Y, Maher DT. 2019. Are methane emissions from mangrove stems a cryptic carbon loss pathway? Insights from a catastrophic forest mortality. *New Phytologist* **224**: 146–154.
- Keppeler F, Hamilton JTG, Brass M, Röckmann T. 2006. Methane emissions from terrestrial plants under aerobic conditions. *Nature* **439**: 187–191.

- King GM. 2003. Contributions of atmospheric CO and hydrogen uptake to microbial dynamics on recent Hawaiian volcanic deposits. *Applied and Environmental Microbiology* 69: 4067–4075.
- King GM, Nanba K. 2008. Distribution of atmospheric methane oxidation and methanotrophic communities on Hawaiian volcanic deposits and soils. *Microbes and Environments* 23: 326–330.
- Kirman S. 2003. *Cycles biogéochimiques et biodiversité en forêt tropicale humide: étude d'une succession primaire sur coulées basaltiques (La Réunion, Ocean Indien)*. PhD thesis, Aix-Marseille University, Marseille, France.
- Kirman S, Strasberg D, Grondin V, Colin F, Gilles J, Meunier JD. 2007. Biomass and litterfall in a native lowland rainforest: Mare Longue Reserve, La Réunion Island, Indian Ocean. *Forest Ecology and Management* 252: 257–266.
- Kreft H, Jetz W, Mutke J, Kier G, Barthlott W. 2008. Global diversity of island floras from a macroecological perspective. *Ecology Letters* 11: 116–127.
- Lenhart K, Behrendt T, Greiner S, Steinkamp J, Well R, Giesemann A, Keppler F. 2019. Nitrous oxide effluxes from plants as a potentially important source to the atmosphere. *New Phytologist* 221: 1398–1408.
- Lenhart K, Weber B, Elbert W, Steinkamp J, Clough T, Crutzen P, Pöschl U, Keppler F. 2015. Nitrous oxide and methane emissions from cryptogamic covers. *Global Change Biology* 21: 3889–3900.
- Machacova K, Bäck J, Vanhatalo A, Halmeenmäki E, Kolari P, Mammarella I, Pumpanen J, Acosta M, Urban O, Pihlatie M. 2016a. *Pinus sylvestris* as a missing source of nitrous oxide and methane in boreal forest. *Scientific Reports* 6: 23410.
- Machacova K, Halmeenmäki E, Pihlatie M, Urban O. 2016b. Seasonal courses of methane fluxes in boreal trees. *Report Series in Aerosol Science* 189: 308–311.
- Machacova K, Maier M, Svobodova K, Lang F, Urban O. 2017. Cryptogamic stem covers may contribute to nitrous oxide consumption by mature beech trees. *Scientific Reports* 7: 13243.
- Machacova K, Papen H, Kreuzwieser J, Rennenberg H. 2013. Inundation strongly stimulates nitrous oxide emissions from stems of the upland tree *Fagus sylvatica* and the riparian tree *Alnus glutinosa*. *Plant and Soil* 364: 287–301.
- Machacova K, Pihlatie M, Halmeenmäki E, Pavelka M, Dušek J, Bäck J, Urban O. 2015. Summer fluxes of nitrous oxide from boreal forest. In: Urban O, Šprtová M, Klem K, eds. *Global change: a complex challenge, conference proceedings*. Brno, Czech Republic: Global Change Research Center, 78–81.
- Machacova K, Vainio E, Urban O, Pihlatie M. 2019. Seasonal dynamics of stem N₂O exchange follow the physiological activity of boreal trees. *Nature Communications* 10: 4989.
- Maier M, Machacova K, Lang F, Svobodova K, Urban O. 2018. Combining soil and tree-stem flux measurements and soil gas profiles to understand CH₄ pathways in *Fagus sylvatica* forests. *Journal of Plant Nutrition and Soil Science* 181: 31–35.
- Menyailo OV, Hungate BA. 2005. Tree species effects on potential production and consumption of carbon dioxide, methane, and nitrous oxide: the Siberian afforestation experiment. In: Binkley D, Menyailo OV, eds. *Tree species effects on soils: implications for global change*. NATO Science Series. Dordrecht, the Netherlands: Kluwer Academic, 293–305.
- Meunier JD, Kirman S, Strasberg D, Nicolini E, Delcher E, Keller C. 2010. The output and bio-cycling of Si in a tropical rain forest developed on young basalt flows (La Reunion Island). *Geoderma* 159: 431–439.
- Mylre G, Shindell D, Bréon F-M, Collins W, Fuglestedt J, Huang J, Koch D, Lamarque J-F, Lee D, Mendoza B *et al.* 2013. Anthropogenic and natural radiative forcing. In: Stocker TF, Qin D, Plattner G-K, Tignor M, Allen SK, Boschung J, Nauels A, Xia Y, Bex V, Midgley PM, eds. *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge, UK & New York, NY, USA: Cambridge University Press, 659–740.
- Pangala SR, Enrich-Prast A, Basso LS, Peixoto RB, Bastviken D, Hornibrook ERC, Gatti LV, Ribeiro H, Calzans LSB, Sakuragai CM *et al.* 2017. Large emissions from floodplain trees close the Amazon methane budget. *Nature* 552: 230–234.
- Pangala SR, Moore S, Hornibrook ERC, Gauci V. 2013. Trees are major conduits for methane egress from tropical forested wetlands. *New Phytologist* 197: 524–531.
- Potgieter LJ, Wilson JR, Strasberg D, Richardson DM. 2014. *Casuarina* invasion alters primary succession on lava flows on La Réunion Island. *Biotropica* 46: 268–275.
- Richards PW. 1996. *The tropical rain forest*. Cambridge, UK: Cambridge University Press.
- Rusch H, Rennenberg H. 1998. Black alder (*Alnus glutinosa* (L.) Gaertn.) trees mediate methane and nitrous oxide emission from the soil to the atmosphere. *Plant and Soil* 201: 1–7.
- Saunois M, Bousquet P, Poulter B, Peregon A, Ciais P, Canadell JG, Dlugokencky EJ, Etiope G, Bastviken D, Houweling S *et al.* 2016. The global methane budget 2000–2012. *Earth System Science Data* 8: 697–751.
- Schindler T, Mander Ü, Machacova K, Espenberg M, Krasnov D, Escuer-Gatius J, Veber G, Pärn J, Soosaar K. 2020. Short-term flooding increases CH₄ and N₂O emissions from trees in a riparian forest soil–stem continuum. *Scientific Reports* 10: 3204.
- Shvidenko A, Barber CV, Persson R, Gonzalez P, Hassan R, Lakyda P, McCallum I, Nilsson S, Pulhin J, van Rosenburg B *et al.* 2005. Forest and woodland systems. In: Hassan R, Scholes R, Ash N, eds. *Ecosystems and human well-being: current state and trends*. Washington, DC, USA: Island Press, 585–621.
- Sjögersten S, Siegenthaler A, Lopez OR, Aplin P, Turner B, Gauci V. 2020. Methane emissions from tree stems in neotropical peatlands. *New Phytologist* 225: 769–781.
- Smart DR, Bloom AJ. 2001. Wheat leaves emit nitrous oxide during nitrate assimilation. *Proceedings of the National Academy of Sciences, USA* 98: 7875–7878.
- Smith KA, Ball T, Conen F, Dobbie KE, Massheder J, Rey A. 2003. Exchange of greenhouse gases between soil and atmosphere: interactions of soil physical factors and biological processes. *European Journal of Soil Science* 54: 779–791.
- Sundqvist E, Crill P, Mölder M, Vestin P, Lindroth A. 2012. Atmospheric methane removal by boreal plants. *Geophysical Research Letters* 39: L21806.
- Turner IM, Corlett RT. 1996. The conservation value of small, isolated fragments of lowland tropical rain forest. *Tree* 11: 330–333.
- Warlo H, Machacova K, Nordstrom N, Maier M, Laemmel T, Roos A, Schack-Kirchner H. 2018. Comparison of portable devices for sub-ambient concentration measurements of methane (CH₄) and nitrous oxide (N₂O) in soil research. *International Journal of Environmental Analytical Chemistry* 98: 1030–1037.
- Welch B, Gauci V, Sayer EJ. 2018. Tree stem bases are sources of CH₄ and N₂O in a tropical forest on upland soil during the dry to wet season transition. *Global Change Biology* 25: 361–372.
- Wen Y, Corre MD, Rachow C, Chen L, Veldkamp E. 2017. Nitrous oxide emissions from stems of alder, beech and spruce in a temperate forest. *Plant and Soil* 420: 423–434.
- Yu K, Chen G. 2009. Nitrous oxide emissions from terrestrial plants: observations, mechanisms and implications. In: Sheldon AI, Barnhart EP, eds. *Nitrous oxide emissions research progress*. Hauppauge, NY, USA: Nova Science Publishers, 85–104.

Supporting Information

Additional Supporting Information may be found online in the Supporting Information section at the end of the article.

Fig. S1 Examples of CH₄, N₂O and CO₂ concentration changes over time in the headspaces of soil chambers, volcanic rock chambers, tree stem chambers and incubation chambers containing cryptogams.

Please note: Wiley Blackwell are not responsible for the content or functionality of any Supporting Information supplied by the authors. Any queries (other than missing material) should be directed to the *New Phytologist* Central Office.