



HAL
open science

Validation and comparison of incoming solar radiation satellite databases on the Atlantic coast of Central Africa

Amine Ouhechou, Nathalie Philippon, Béatrice Morel

► **To cite this version:**

Amine Ouhechou, Nathalie Philippon, Béatrice Morel. Validation and comparison of incoming solar radiation satellite databases on the Atlantic coast of Central Africa. EGU General Assembly 2021, Apr 2021, Vienna, Austria. 10.5194/egusphere-egu21-3493 . hal-03695952

HAL Id: hal-03695952

<https://hal.univ-reunion.fr/hal-03695952v1>

Submitted on 16 Jun 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Validation and comparison of incoming solar radiation satellite databases on the Atlantic coast of Central Africa.

Amine OUHECHOU, Nathalie PHILIPPON and Béatrice MOREL



• Objectives




- ❖ Analyze the mean spatial and temporal variations of solar radiation (light availability for tree photosynthesis)
- ❖ Document the specificity of the Atlantic coast
- ❖ Determine which satellite estimate(s) (in the absence of sufficient in-situ measurements) is the most suitable for the study of solar radiation on the Atlantic coast.

• Data used

Data set	Spatial resolution	Temporal resolution	Temporal range	Satellite	orbit type
CERES-EBAF	1° × 1°	Monthly	2000-03 / 2019-11	Terra + Aqua	Heliosynchronous orbit
CERES-SYN1deg	1° × 1°	Hourly	2000-03 / 2019-11	Terra + Aqua	Heliosynchronous orbit
Ré-analyses ERA5	0.25° × 0.25°	Monthly	1979-01 / 2020-12	---	---
CM SAF-CLARA-A2	0.25° × 0.25°	Monthly	1982-01 / 2015-12	AVHRR sensors on series of NOAA satellites and the METOP satellite.	polar orbit
CAMS-JADE	0.2° × 0.2°	15 minutes	2005-01 / 2018-12	MVIRI/SEVIRI on METEOSAT MSG	geostationary orbit
CM SAF-SARAH 2	0.05° × 0.05°	Hourly	1983-01 / 2020-12	MVIRI/SEVIRI on METEOSAT MSG	geostationary orbit
FAO stations	N/A(station)	Monthly means	---	---	---
Original meteorological stations	N/A(station)	15min to daily	---	---	---

Results and Discussion


Spatial distribution of the annual mean of incoming solar radiation



- The Atlantic coast receives the least quantity of solar radiation
- Satellite products from MSG are the brightest

• Results and Discussion

Hovmöller diagram




☐ The longitudinal extent is 8° - 30°E

- ☐ The minimum of solar radiation is observed in June-August
- ☐ CLARA and ERA5 have a second minimum during the October-November rainy season


• Results and Discussion

Evaluation of products on sub-regions using FAO's
in situ measurements

Climate regionalization of Central Africa



Location of the FAO stations





Philippon, N et al. (2018). CYCLES DIURNES DE RAYONNEMENT SOLAIRE ET FORETS EN AFRIQUE CENTRALE.

Results and Discussion

Mean annual cycle of incoming solar radiation

Products	Correlation	Bias	Spatial resoluti ^o
CERES-EBAF	0,85	-2,32 W/m ²	1° x 1°
CERES-SYN1deg	0,85	-3,51 W/m ²	1° x 1°
ERA5	0,81	13 W/m ²	0,25° x 0,25°
CMSAF-CLARA	0,76	11 W/m ²	0,25° x 0,25°
CMSAF-SARAH	0,77	15 W/m ²	0,05° x 0,05°
CAMS-JADE	0,75	8 W/m ²	0,2° x 0,2°








- The cycle is well reproduced by all the products (R>0,75)
- CERES underestimates and the other products overestimate
- The maximum in February-March and the minimum in July-August



Results and Discussion







Validation and inter-comparison of (CERES-SYN1deg , CMSAF-SARAH and CAMS-JADE) on a daily scale with original meteorological data.

Stations	Country	Temporal resolution	Availability period
Pokola	Congo	15 min	03/2016 – 12/2020
Lopé	Gabon	Daily	01/2012 – 10/2015
Lékédi	Gabon	15/30 min	07/2018 – 07/2019
Mokabi	Congo	15 min	03/2016 – 12/2020
NSIMI	Cameron	Daily	04/2004 – 03/2008
Kissoko	Congo	30 min	04/2004 – 06/2006
Tchizamalou	Congo	15 min	06/2006 – 12/2009
Edea	Cameroun	15 min	01/2020 – 12/2020
Mbandaka	RDC	15 min	11/2018 – 12/2020
Mfou	Cameron	15 min	09/2018 – 12/2020



Results and Discussion

Monthly variation of bias results.



- ☐ The strongest biases are observed during the 2 dry seasons.
- ☐ Strong biases in January-February when it is in the North of the equator and the opposite when it is in the South.

	CERES-SYN1deg	CAMS-JADE	CMSAF-SARAH
Correlation	[0.55 – 0.91]	[0.58 – 0.87]	[0.6 – 0.9]
Bias	[-20 – 51]	[23.8 – 52]	[34.7 – 86]
RMSE	[8.8 – 83]	[19 – 73]	[21 – 113]




• Results and Discussion

CMSAF-SARAH




CERES-SYN1deg

CAMS-JADE

All years




Dry season JJAS



☐ Products see too many bright days and fewer obscure days

• Conclusion

The incoming solar radiation data from 5 satellite estimates, including CERES-EBAF, CERES-SYN1deg, CMSAF-SARAH, CMSAF-CLARA and CAMS-JADE, as well as the ERA5 reanalyses



The comparison of the six products shows that the Atlantic coast of Central Africa receives the least amount of solar radiation in all products.

Less of Bias for CERES-EBAF and CERES SYN1deg products which underestimate the incident solar radiation while the other satellite products overestimate it. CMSAF-SARAH overestimates the most.



The satellite products represent well the spatial distribution, and reproduce well the mean annual cycle of incident solar radiation