



HAL
open science

Scabies incidence and association with skin and soft tissue infection in Loyalty Islands Province, New Caledonia: A 15-year retrospective observational study using electronic health records

Yves-Marie Ducrot, Enzo Bruno, Jean-Marc Franco, Loïc Raffray, Samuel Beneteau, Antoine Bertolotti

► To cite this version:

Yves-Marie Ducrot, Enzo Bruno, Jean-Marc Franco, Loïc Raffray, Samuel Beneteau, et al.. Scabies incidence and association with skin and soft tissue infection in Loyalty Islands Province, New Caledonia: A 15-year retrospective observational study using electronic health records. *PLoS Neglected Tropical Diseases*, 2022, 16 (9), pp.e0010717. 10.1371/journal.pntd.0010717 . hal-04104411

HAL Id: hal-04104411

<https://hal.univ-reunion.fr/hal-04104411v1>

Submitted on 24 May 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

RESEARCH ARTICLE

Scabies incidence and association with skin and soft tissue infection in Loyalty Islands Province, New Caledonia: A 15-year retrospective observational study using electronic health records

Yves-Marie Ducrot^{1*}, Enzo Bruno², Jean-Marc Franco², Loïc Raffray^{3,4}, Samuel Beneteau⁵, Antoine Bertolotti^{6,7}

1 Centre Médico-Social de Wé, Direction de l'Action Communautaire et de l'Action Sanitaire, Province des Iles Loyauté, Lifou, Nouvelle-Calédonie, **2** Département de médecine générale, Université de La Réunion, Saint Pierre, La Réunion, France, **3** Département de médecine interne, CHU de la Réunion, Saint Denis, La Réunion, France, **4** Unité Mixte de Recherche Processus Infectieux en Milieu Insulaire Tropical (UMR PIMIT), Université de La Réunion, CNRS 9192, INSERM 1187, IRD 249, Plateforme Technologique CYROI, Sainte-Clotilde, France, **5** Unité de soutien méthodologique, CHU de la Réunion, Saint Denis, France, **6** Service des Maladies Infectieuses–Dermatologie, CHU de La Réunion, Saint Pierre, La Réunion, France, **7** Inserm CIC1410, CHU de La Réunion, Saint Pierre, La Réunion, France

* yvesmarie.ducrot@gmail.com



OPEN ACCESS

Citation: Ducrot Y-M, Bruno E, Franco J-M, Raffray L, Beneteau S, Bertolotti A (2022) Scabies incidence and association with skin and soft tissue infection in Loyalty Islands Province, New Caledonia: A 15-year retrospective observational study using electronic health records. *PLoS Negl Trop Dis* 16(9): e0010717. <https://doi.org/10.1371/journal.pntd.0010717>

Editor: Michael Marks, London School of Hygiene and Tropical Medicine, UNITED KINGDOM

Received: April 30, 2022

Accepted: August 3, 2022

Published: September 6, 2022

Copyright: © 2022 Ducrot et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: Data cannot be shared publicly because of restrictions on the database. Data are available upon a reasonable request to the "Direction du Numérique et des Services d'Information" and "Direction de l'Action Communautaire et de l'Action Sanitaire" (contact via Tony PEYRO t-peyro@loyalty.nc and Marie-Rose WAIA mr-waia@loyalty.nc) for researchers who meet the criteria for access to confidential data.

Abstract

Background

Scabies and its complications are a public health problem in the low- and middle-income countries of the Western Pacific region. However, no data are available for the relatively wealthy French territory of New Caledonia. This study aimed to determine the incidence of scabies and its association with skin and soft tissue infection (SSTI) in Loyalty Islands Province (LIP) (20,000 inhabitants), New Caledonia.

Methodology/Principal findings

This retrospective observational study reviewed cases of scabies and SSTI extracted from the electronic health record databases of LIP clinics for the period 2004–2018. Data were validated through double sampling. The overall scabies incidence rate (IR) and scabies IRs by sex and age group were calculated. Scabies seasonality was evaluated. For children <5 years, the presence of SSTI was compared between the 3-month period preceding scabies diagnosis/treatment and the 3-month period preceding the 1-year anniversary of scabies diagnosis/treatment (self-matching).

A total of 16,843 scabies cases were extracted using a detection algorithm with a sensitivity of 96.7% and a specificity of 99.9%. From 2004 to 2018, the average overall scabies IR was 5.9% and the average scabies IR in children <1 year was 18.4%. Almost two-thirds of children aged 14 years had a history of scabies. Females were more affected, especially in the 20–39 age group (sex ratio>2). A strong seasonality was observed, with a 30% increase

Funding: The author(s) received no specific funding for this work.

Competing interests: The authors have declared that no competing interests exist.

in winter. In children <5 years, SSTIs were 4.3 times more frequent in the 3 months preceding the scabies diagnosis than in the 3 months preceding the 1-year anniversary of scabies treatment ($p < 0.001$).

Conclusions

Although health care is much better in New Caledonia than in neighboring countries, scabies is highly endemic in LIP. The disease is especially common in children <2 years and is associated with many SSTIs in children <5 years. Mass drug administration should be considered.

Author summary

Scabies is a known public health problem in the Pacific region. This study aimed to determine the importance of scabies and associated skin infections in the Loyalty Island Province (LIP) of New Caledonia (20,000 inhabitants).

The provincial electronic medical database was searched for all cases of scabies and skin infection that occurred in the LIP population between 2004 and 2018. To evaluate the impact of scabies on skin infections, we measured the decrease in the number of skin infections in children <5 years one year after scabies treatment.

Over the study period, the average number of scabies cases occurring each year was 6 per 100 inhabitants. In children <1 year, this number was 18 per 100 inhabitants. We found that 80% of children aged 15 years had experienced at least one episode of scabies. We also observed a fourfold decrease in the number of skin infections in children <5 years 1 year after treatment. Scabies was more frequent (+30%) in winter.

Introduction

Scabies is a skin infestation caused by *Sarcoptes scabiei var. hominis* that is transmitted through close skin-to-skin contact and is responsible for pruritic skin rashes. Secondary skin and soft tissue infections (SSTIs) associated with this parasitic disease can be severe and are sometimes complicated by acute post-streptococcal glomerulonephritis or acute rheumatic fever (ARF) [1–4]. Scabies is estimated to affect 200 million people worldwide. In 2017, the World Health Organization (WHO) added scabies to the list of neglected tropical diseases [5].

While scabies and its complications are a known public health problem in the low- and middle-income countries of the Western Pacific region [2], no data are currently available for the relatively wealthy French territory of New Caledonia. However, the Melanesian community, which accounts for half of the population of New Caledonia, has generally low standard of living and suffers a heavy burden of health problems, including severe SSTIs and ARF [6]. One can wonder whether scabies plays a role in the occurrence of these diseases in this population.

The aim of this study was to determine the incidence of scabies and its association with SSTI in Loyalty Islands Province (LIP), New Caledonia.

Methods

Ethics statement

The study was approved by the New Caledonia Consultative Ethics Committee (# 202106004) and registered with the French Data Protection Authority (Commission Nationale de

l'Informatique et des Libertés, #2211634v0). The study was advertised via posters in all local clinics of LIP.

Setting

New Caledonia is a French overseas territory (280,000 inhabitants) divided into 3 provinces and located in the Western Pacific between Australia and Fiji. The present study was conducted in LIP, a sparsely populated rural province with 20,000 inhabitants living in 83 small tribes on 3 islands (Ouvéa, Lifou, and Maré). The population of LIP has access to a modern and free public primary and secondary health care system. The density of general practitioners (GPs) is 90 per 100,000 inhabitants, with 93% of the population having access to a GP within a 25-minute drive. The climate is subtropical with 2 marked seasons. The winter is dry and cold and occurs from May to September.

In New Caledonia, scabies is treated in accordance with the guidelines of the French National Authority for Health. Patients receive two doses of: (i) oral ivermectin or (ii) topical benzyl benzoate or (iii) topical permethrin. Simultaneous treatment of all close contacts is also recommended [7,8].

Study design, sampling, and procedures

This retrospective observational study covered the population attending the local clinics of LIP over the period 2004–2018. These clinics offer care to almost 90% of the LIP population and provide systematic prevention consultations to nearly 100% of LIP children <2 years.

Cases of scabies and SSTI were extracted from the electronic health record (EHR) databases of LIP clinics. These open-source relational databases, which are accessible via an EHR platform software, have been in use in LIP for the past 20 years.

Scabies cases were detected using a Structured Query Language (SQL) algorithm (S1 Text). They were defined as: any medical visit (i) coded as “scabies” (B86) according to the 10th International Classification of Diseases (ICD-10) coding system; (ii) AND/OR with a specific anti-scabies drug recorded in the prescription field or in the free text (iii) AND/OR with the keywords “*gale*” (scab in French) or “scab” recorded in the free text. Two successive visits were classified as separate cases if there was an interval of at least 28 days between them. The algorithm excluded negative formulas (e.g., “not scabies”) and false friends (e.g., “éGALEMENT”). Some spelling mistakes were allowed. A similar detection algorithm was used for cases of SSTI (S2 Text).

To evaluate the specificity and sensitivity of the detection algorithms, a random draw of 2,714 visits was manually proofread by a physician. This validation approach is considered the gold standard in case detection studies.

The overall incidence rate (IR) of scabies and scabies IRs by sex, age group, and tribe were calculated. In order to assess the cumulative incidence of scabies from 0 to 15 years of age, we extracted the records of 4,732 children <15 years who had completed at least 5 preventive medical visits (mother and child visits, vaccination visits, or school visits) before the age of 5 years, including 1 visit before the age of 1 year. The incidence rate was calculated and averaged for each birth cohort per age group. Only the first episode of scabies was counted in cases of relapse. For children <5 years, the presence of SSTI was compared between the 3-month period preceding scabies diagnosis/treatment and the 3-month period preceding the 1-year anniversary of scabies diagnosis/treatment (self-matching). The latter analysis was restricted to children <5 years who had completed at least 2 preventive medical visits. Scabies seasonality was also evaluated.

Demographic data were obtained by averaging data from the 2004, 2009, 2014, and 2019 population censuses. Climate data were acquired from the meteorological department of New Caledonia.

Statistical analysis

Incidence rates were calculated after age standardization. Confidence intervals were calculated with a confidence interval of 95% (95% CI). A modified McNemar's test was used for the analysis of the association between scabies and SSTI in children <5 years. Seasonality was assessed using Pearson correlation coefficient. Statistical analyses were performed with R software version R 4.0.4.

Results

Case detection

The detection algorithm for scabies had a sensitivity of 96.7% (95% CI: 90.2–100%) and a specificity of 99.9% (95% CI: 99.8–100%) with a positive predictive value (PPV) of 96.7% and a negative predictive value (NPV) of 99.9%. Overall, 42% of scabies cases were extracted by keyword alone, 13% by prescription alone, and less than 1% by ICD-10 coding alone; moreover, 5% of cases were extracted using the 3 extraction methods (S1 Fig). The performance of the detection algorithm for SSTI was lower, with a sensitivity of 74.6% (95% CI: 69.3–79.9%), a specificity of 99.3% (95% CI: 98.9–99.6%), a PPV of 91.9% and a NPV of 97.4%.

Incidence of scabies

A total of 16,843 cases of scabies were diagnosed in 10,063 individuals over the 2004–2018 period. The recurrence rate was 36.0%. The annual overall average scabies IR was 5.9%. This IR remained relatively stable over the study period (S2 Fig).

The age groups <19 years and >75 years had the highest scabies IRs. The male/female sex ratio (SR) was in favor of females 0.76/1 (95% confidence interval (CI): 0.67, 0.82). In the 20–39 age group, females were twice as affected as males (SR = 0.42/1; 95%CI:0.31,0.53) (Fig 1).

The IR of scabies in children <1 year was 18.4%.




Fig 1. Scabies incidence rate per 1,000 inhabitants by sex and age group. The histograms represent the average scabies incidence rate in females (blue) and males (grey) according to age group.

<https://doi.org/10.1371/journal.pntd.0010717.g001>




Fig 2. Scabies incidence rate by age group and cumulative incidence from 0 to 15 years. The histogram represents the average scabies incidence rate by age group; the curve represents the cumulative incidence of scabies from 0 to 15 years.

<https://doi.org/10.1371/journal.pntd.0010717.g002>

For the entire period of study, no significant differences were observed in the scabies IR between the 3 islands of LIP. However, from 2014 onwards, the scabies IR was higher in Maré than in the other islands (S2 Fig). The scabies IR varied significantly between the different tribes of LIP (S3 Fig).

Cumulative incidence of scabies

The cumulative incidence of scabies was 51.0% in children aged 2 years, 66.3% in children aged 5 years and 80.2% in children aged 15 years (Fig 2 and S1 Table).

Seasonality of scabies

A statistically significant inverse correlation between monthly scabies cases and monthly temperatures was observed ($R = -0.738$; $p = 0.01$), with an increase of nearly 30% between August and January (Fig 3).

Impact of scabies on skin and soft tissue infections

In children <5 years, SSTIs were 4.3 times more frequent in the 3 months preceding the scabies diagnosis than in the 3 months preceding the 1-year anniversary of scabies treatment ($p < 0.001$) (Fig 4).

Discussion

Our study found scabies to be highly endemic in LIP, despite the fact that access to care is much better in New Caledonia than in neighboring countries. Scabies was particularly common in children <2 years and was associated with many SSTIs in children <5 years. It also showed seasonal and geographical variations.

Clinical research based on EHRs is becoming increasingly common in epidemiology [9]. In the absence of systematic IDC coding, cases can be identified by extracting information from the free text of EHRs [10]. While one study in New Zealand did use EHRs from general




Fig 3. Average monthly number of scabies cases in Loyalty Islands Province from 2004 to 2018 and average monthly temperatures (°C) in New Caledonia from 1981 to 2010. The histogram represents the average monthly number of scabies cases in Loyalty Islands Province (left axis); the curve represents average monthly temperatures (°C) in New Caledonia (right axis).

<https://doi.org/10.1371/journal.pntd.0010717.g003>

practices for the detection of SSTI cases [11], to our knowledge our study is the first to use EHRs for the detection of scabies cases.

The overall scabies IR in LIP was 5.9%, a figure 15 to 25 times higher than those reported in European countries [12,13]. Most of the studies conducted in other countries of the Western




Fig 4. Number of skin and soft tissue infections in children <5 years in the 3 months preceding scabies diagnosis/treatment and in the 3 months preceding the 1-year anniversary of scabies diagnosis/treatment.

<https://doi.org/10.1371/journal.pntd.0010717.g004>

Pacific have produced data on scabies prevalence, making it difficult to compare the level of scabies endemicity between these and LIP. However, one study conducted in Fiji [14] did report a scabies IR of 2.87% for the 2018–2019 period. While this figure is slightly lower than the IR observed in our study, the level of scabies endemicity is likely to be similar between LIP and Fiji given that these show important differences in care-seeking behavior and access to care.

In our study, the peak of scabies incidence was observed in children aged 2 years. This finding has been reported in one other study [15] that used a longitudinal methodology as we did. Most studies of scabies have found that the disease peaks at the age of 1 year, a phenomenon that is generally explained by a lack of immunity in very young children. The fact that children in New Caledonia are routinely followed up until the age of 2 years could explain the peak of incidence observed at that age in our study.

The SR was strongly in favor of females in our population, especially in the 20–39 age group. Sex imbalance has rarely been reported in the literature on scabies [16]. It could be linked to local cultural practices, as Melanesian mothers generally sleep with their babies. However, it could also reflect a bias due to our use of medical visit data since mothers accompanying children diagnosed with scabies are systematically screened for the disease in New Caledonia.

Our study found a strong seasonality of scabies, with a marked increase in cases between August and January. A similar pattern has been observed in other countries with well-defined seasons [17–19]—one exception being South Korea, where scabies cases have been shown to peak in fall and spring [16]. The recrudescence of scabies in winter can be explained by the increase in promiscuity in cold weather, which in turn favors transmission. Another explanation is that *Sarcoptes* reproduce and survive more easily in fomites with colder temperatures [20].

Despite a very homogeneous lifestyle, significant differences in scabies incidence were found between the tribes of LIP. This phenomenon, which was also observed in the Solomon Islands, has no clear explanation [21] (S3 Fig).

The link between scabies and SSTIs is well known. It is especially well documented for impetigo caused by *Streptococcus pyogenes*, a pathogen that thrives in grooves and scratch lesions as well as in the digestive tract of *Sarcoptes* [22]. To date, however, few studies have quantified the risk of SSTI in patients with scabies [3]. A study using the same methodology as we did, but with a 15-day interval between the occurrence of SSTI and scabies diagnosis/treatment, found an even stronger association [23]. Our study selected a 3-month interval because this corresponds to the peak of natural scabies infestation in humans [24].

In New Caledonia, ARF is especially common in the Melanesian population, with an incidence of 262/100,000 [25,26]. However, unlike the situation in many tropical countries with a high prevalence of ARF, streptococcal angina is rare in New Caledonia [27,28]. The high prevalence of ARF in our population could be explained by the frequent occurrence of impetigo caused by *Streptococcus pyogenes* [29].

Our study has some limitations. The scabies IR could be underestimated as cases were identified based on a presumptive diagnosis that did not follow precise clinical criteria but depended on the physician's experience. Indeed, GPs working in LIP come mostly from metropolitan France where the scabies IR is low. Given the high GP turnover (the average internship duration is 6 months), it is likely that many cases were missed. Conversely, the scabies IR could be overestimated because asymptomatic contacts are also treated for scabies in New Caledonia. However, the duration and multicentric nature of the study, the large sample size, and the near completeness of data have likely counteracted these biases. We can therefore be fairly confident that the scabies IR in our study is close to the actual IR, particularly among children <2 years. Another limitation is that the 20,000 inhabitants covered in our study represent only

7% of the total population of New Caledonia. Given the variations in socio-cultural and ethnic characteristics across the territory of New Caledonia, our findings cannot be considered representative of other provinces.

Although the population of LIP has access to high-quality healthcare, the scabies IR remained relatively stable in the province over the study period. This finding suggests that more drastic public health measures are needed to reduce the burden of scabies and associated SSTIs in LIP [30–32]. According to a recent WHO framework for scabies control [33,34], ivermectin-based mass drug administration (MDA) should be initiated in areas with a community prevalence $\geq 10\%$ or a prevalence in school children $\geq 15\%$ (as measured using a school mapping strategy) and should be avoided in those with a community prevalence $< 2\%$. In areas with a community prevalence $> 2\%$ and $< 10\%$, either intensive disease management (IDM) or MDA should be initiated based on an assessment of the local context, as no evidence-based recommendations are currently available to decide on the most appropriate control strategy. Unfortunately, recent data on scabies prevalence in LIP are lacking. Nevertheless, we have seen that the level of scabies endemicity in the province is very close to that in neighboring Fiji [14], where a prevalence of 29% has recently been reported [1]. Accordingly, LIP could meet the WHO threshold for MDA implementation. On the other hand, a 1989 cross-sectional study conducted in a rural area of New Caledonia reported a prevalence of 11% in 269 children < 5 years [35]. The relative stability of the scabies IR observed in our study suggests that these prevalence data are still valid today. We can infer from this that either IDM and MDA could be implemented in LIP as per the WHO framework. The first strategy would add little to the management protocol already in place in LIP because it consists mainly in extending treatment to asymptomatic contacts. Mass drug administration, which has shown optimal scabies reduction in small and isolated rural communities with a minimum coverage of 80%, would likely have a greater chance of success as LIP is characterized by insularity, a rural lifestyle, and the presence of small tribes of 50 to 800 inhabitants with a strong political structure. It should be noted, however, that daily aerial exchanges with the large city of Nouméa mitigate this insularity. Moreover, the low vaccination rate against COVID-19 in the province suggests that sufficient MDA coverage would be difficult to achieve. Given these uncertainties, New Caledonia health authorities should consider taking advantage of the ivermectin-based MDA program against filariasis, which is to be launched at the end of 2022 in the smallest island of LIP, to determine the prevalence of scabies and the cost of implementing MDA in all islands of the province.

Despite the fact that access to care is much better in New Caledonia than in neighboring countries, scabies is highly endemic in LIP and is responsible for many SSTIs in children < 5 years. Moreover, the scabies IR has remained relatively stable over time, and the recurrence rate is fairly high. Prevalence studies are needed to determine whether more drastic public health measures, including ivermectin-based MDA, must be implemented in LIP to reduce the burden of scabies and associated SSTIs.

Supporting information

S1 Text. Extraction of scabies cases from an open-source SQL relational database management system, using Structured Language Query.

(DOCX)

S2 Text. Extraction of skin and soft tissue infection cases from an open-source SQL relational database management system, using Structured Language Query.

(DOCX)

S1 Fig. Venn diagram of scabies case extraction. The blue circle represents the proportion of cases extracted by keyword alone, the green circle the proportion of cases extracted by prescription alone, and the orange circle the proportion of cases extracted by ICD-10 coding alone; the intersection of circles represents the proportion of cases extracted using two or more of these extraction methods.

(TIF)

S2 Fig. Evolution of average scabies incidence rates in Loyalty Islands Province and its 3 islands. The histogram represents the average overall incidence rate for the period 2004–2018; the solid curve represents the average incidence rate on the island of Lifou, the large dotted curve represents the average incidence rate on the island of Maré, and the small dotted curve represents the average incidence rate on the island of Ouvéa.

(TIF)

S3 Fig. Differences in average incidence rates between the tribes of Loyalty Islands Province. Map created using qGIS, an open source GIS, with two datasets: a) from open source data <https://georep-dtsi-sgt.opendata.arcgis.com/datasets/dtsi-sgt::limites-administratives-terrestres-1/explore?layer=1&location=-21.196392%2C165.834400%2C7.88> b) tribes limits was provided by the Information System Department of the Loyalty Island Province.

(TIF)

S1 Data. Excel spreadsheet containing, in separate sheets, the underlying numerical data for Figs 1, 2, 3, 4, S1, S2, and S3.

(XLSX)

S1 Table. Incidence rate for each age group of 14 birth cohort, mean incidence rate and cumulative incidence.

(DOCX)

Acknowledgments

We would like to thank Julien Rannou for his computer assistance, Marie Rose Waia, Emile Gaze and Maryline Brun for supporting this work, and Quentin Amic-Desvaud for his participation in the validation of SSTI data. We would also like to thank our copy editor Arianne Dorval.

Author Contributions

Conceptualization: Yves-Marie Ducrot.

Data curation: Yves-Marie Ducrot.

Formal analysis: Yves-Marie Ducrot, Enzo Bruno, Samuel Beneteau, Antoine Bertolotti.

Investigation: Yves-Marie Ducrot, Enzo Bruno.

Methodology: Yves-Marie Ducrot, Samuel Beneteau, Antoine Bertolotti.

Project administration: Yves-Marie Ducrot.

Supervision: Antoine Bertolotti.

Validation: Antoine Bertolotti.

Visualization: Jean-Marc Franco, Loïc Raffray.

Writing – original draft: Yves-Marie Ducrot, Enzo Bruno, Antoine Bertolotti.

Writing – review & editing: Jean-Marc Franco, Loïc Raffray, Samuel Beneteau.

References

1. Romani L, Koroivueta J, Steer AC, Kama M, Kaldor JM, Wand H, et al. Scabies and Impetigo Prevalence and Risk Factors in Fiji: A National Survey. *PLOS Neglected Tropical Diseases*. 2015; 9: e0003452. <https://doi.org/10.1371/journal.pntd.0003452> PMID: 25738499
2. Engelman D, Cantey PT, Marks M, Solomon AW, Chang AY, Chosidow O, et al. The public health control of scabies: priorities for research and action. *The Lancet*. 2019; 394: 81–92. [https://doi.org/10.1016/S0140-6736\(19\)31136-5](https://doi.org/10.1016/S0140-6736(19)31136-5) PMID: 31178154
3. Lynar S, Currie BJ, Baird R. Scabies and mortality. *The Lancet Infectious Diseases*. 2017; 17: 1234. [https://doi.org/10.1016/S1473-3099\(17\)30636-9](https://doi.org/10.1016/S1473-3099(17)30636-9) PMID: 29173877
4. Thornley S, King R, Marshall R, Oakley A, Sundborn G, Harrower J. How strong is the relationship between scabies and acute rheumatic fever? An analysis of neighbourhood factors. *J Paediatr Child Health*. 2019. <https://doi.org/10.1111/jpc.14697> PMID: 31774599
5. Engelman D, Steer A. Control Strategies for Scabies. *TropicalMed*. 2018; 3: 98. <https://doi.org/10.3390/tropicalmed3030098> PMID: 30274494
6. Le Hello S, Doloy A, Baumann F, Roques N, Coudene P, Rouchon B, et al. Clinical and microbial characteristics of invasive *Streptococcus pyogenes* disease in New Caledonia, a region in Oceania with a high incidence of acute rheumatic fever. *J Clin Microbiol*. 2010; 48: 526–530. <https://doi.org/10.1128/JCM.01205-09> PMID: 19955276
7. Salavastru CM, Chosidow O, Boffa MJ, Janier M, Tiplica GS. European guideline for the management of scabies. *J Eur Acad Dermatol Venereol*. 2017; 31: 1248–1253. <https://doi.org/10.1111/jdv.14351> PMID: 28639722
8. Bitar D, Caumes E, Chandre F, Del Guidice P, Gehanno J-F, Le Goaster C, et al. Recommendations relatives à la conduite à tenir devant un ou plusieurs cas de gale. Avis et rapports du Haut Conseil de santé publique. 9 Nov 2012. Available: https://www.hcsp.fr/Explore.cgi/Telecharger?NomFichier=hcsp20122209_conduitegale.pdf. Accessed 30 Jun 2022.
9. Fu S, Carlson LA, Peterson KJ, Wang N, Zhou X, Peng S, et al. Natural Language Processing for the Evaluation of Methodological Standards and Best Practices of EHR-based Clinical Research. *AMIA Jt Summits Transl Sci Proc*. 2020; 2020: 171–180. Available: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7233049/> PMID: 32477636
10. Ford E, Carroll JA, Smith HE, Scott D, Cassell JA. Extracting information from the text of electronic medical records to improve case detection: a systematic review. *Journal of the American Medical Informatics Association*. 2016; 23: 1007–1015. <https://doi.org/10.1093/jamia/ocv180> PMID: 26911811
11. Gu Y, Kennelly J, Warren J, Nathani P, Boyce T. Automatic Detection of Skin and Subcutaneous Tissue Infections from Primary Care Electronic Medical Records. *Stud Health Technol Inform*. 2015; 214: 74–80. <https://doi.org/10.3233/978-1-61499-558-6-74> PMID: 26210421
12. van Deursen B, Hooiveld M, Marks S, Snijdwind I, van den Kerkhof H, Wintermans B, et al. Increasing incidence of reported scabies infestations in the Netherlands, 2011–2021. Mossong J, editor. *PLoS ONE*. 2022; 17: e0268865. <https://doi.org/10.1371/journal.pone.0268865> PMID: 35749401
13. Bitar D, Thiolet J-M, Haeghebaert S, Castor C, Poujol I, Coignard B. La gale en France entre 1999 et 2010: augmentation de l'incidence et implications en santé publique. *Ann Dermatol Vénérologie*. 2012; 139.
14. Thean LJ, Romani L, Engelman D, Jenney A, Wand H, Mani J, et al. Prospective Surveillance of Primary Healthcare Presentations for Scabies and Bacterial Skin Infections in Fiji, 2018–2019. *The American Journal of Tropical Medicine and Hygiene*. 2021; 105: 230–237. <https://doi.org/10.4269/ajtmh.20-1459> PMID: 34029210
15. Clucas DB, Carville KS, Connors C, Currie BJ, Carapetis JR, Andrews RM. Disease burden and healthcare clinic attendances for young children in remote aboriginal communities of northern Australia. *Bull World Health Organ*. 2008; 86: 275–281. <https://doi.org/10.2471/blt.07.043034> PMID: 18438516
16. Kim J-H, Cheong H-K. Epidemiologic Trends and Seasonality of Scabies in South Korea, 2010–2017. *Korean J Parasitol*. 2019; 57: 399–404. <https://doi.org/10.3347/kjp.2019.57.4.399> PMID: 31533406
17. Liu J-M, Wang H-W, Chang F-W, Liu Y-P, Chiu F-H, Lin Y-C, et al. The effects of climate factors on scabies. A 14-year population-based study in Taiwan. *Parasite*. 2016; 23: 54. <https://doi.org/10.1051/parasite/2016065> PMID: 27905271
18. Mimouni D, Ankol OE, Davidovitch N, Gdalevich M, Zangvil E, Grotto I. Seasonality trends of scabies in a young adult population: a 20-year follow-up. *Br J Dermatol*. 2003; 149: 157–159. <https://doi.org/10.1046/j.1365-2133.2003.05329.x> PMID: 12890210

19. Tuzun Y, Kotogyan A, Cenesizoglu E, Baransu O, Ozarmagan G, Ural A, et al. The epidemiology of scabies in Turkey. *Int J Dermatol*. 1980; 19: 41–44. <https://doi.org/10.1111/j.1365-4362.1980.tb01993.x> PMID: 7358440
20. Sokolova TV, Radchenko MI, Lange AB. The seasonality of scabies morbidity and the fertility of the itch mite *Sarcoptes scabiei* de Geer as an index of the activity of a population of the causative agent. *Vestn Dermatol Venerol*. 1989; 11: 12–15.
21. Lake SJ, Engelman D, Sokana O, Nasi T, Boara D, Grobler AC, et al. Defining the need for public health control of scabies in Solomon Islands. *PLOS Neglected Tropical Diseases*. 2021; 15: e0009142. <https://doi.org/10.1371/journal.pntd.0009142> PMID: 33617544
22. McCarthy J, Kemp D, Walton S, Currie B. Scabies: more than just an irritation. *Postgrad Med J*. 2004; 80: 382–387. <https://doi.org/10.1136/pgmj.2003.014563> PMID: 15254301
23. Aung PTZ, Cuningham W, Hwang K, Andrews RM, Carapetis JR, Kearns T, et al. Scabies and risk of skin sores in remote Australian Aboriginal communities: A self-controlled case series study. *PLOS Neglected Tropical Diseases*. 2018; 12: e0006668. <https://doi.org/10.1371/journal.pntd.0006668> PMID: 30044780
24. Mellanby K. The development of symptoms, parasitic infection and immunity in human scabies. *Parasitology*. 1944; 35: 197–206. <https://doi.org/10.1017/S0031182000021612>
25. Corsenac P, Heenan RC, Roth A, Rouchon B, Guillot N, Hoy D. An epidemiological study to assess the true incidence and prevalence of rheumatic heart disease and acute rheumatic fever in New Caledonian school children. *Journal of Paediatrics and Child Health*. 2016; 52: 739–744. <https://doi.org/10.1111/jpc.13185> PMID: 27203400
26. Chatard J, Dubois T, Espinosa F, Kamblock J, Ledos P, Tarpinian E, et al. Screening Rheumatic Heart Disease in 1530 New Caledonian Adolescents. *Journal of the American Heart Association*. 2020; 9: e015017. <https://doi.org/10.1161/JAHA.119.015017> PMID: 32336214
27. McDonald MI, Towers RJ, Andrews RM, Bengner N, Currie BJ, Carapetis JR. Low rates of streptococcal pharyngitis and high rates of pyoderma in Australian aboriginal communities where acute rheumatic fever is hyperendemic. *Clin Infect Dis*. 2006; 43: 683–689. <https://doi.org/10.1086/506938> PMID: 16912939
28. Elazzouzi R. Performances d'un test de diagnostic rapide antigenique dans le diagnostic des angines à streptocoque bêta hémolytique du groupe A dans un dispensaire en Nouvelle-Calédonie en population mélanésienne. Thesis, Bordeaux. 2017. Available: <https://dumas.ccsd.cnrs.fr/dumas-01488605>
29. Parks T, Smeesters PR, Steer AC. Streptococcal skin infection and rheumatic heart disease: Current Opinion in Infectious Diseases. 2012; 25: 145–153. <https://doi.org/10.1097/QCO.0b013e3283511d27> PMID: 22327467
30. Rinaldi G, Porter K. Mass drug administration for endemic scabies: a systematic review. *Trop Dis Travel Med Vaccines*. 2021; 7: 21. <https://doi.org/10.1186/s40794-021-00143-5> PMID: 34193305
31. Thean LJ, Romani L, Engelman D, Wand H, Jenney A, Mani J, et al. Prevention of bacterial complications of scabies using mass drug administration: A population-based, before-after trial in Fiji, 2018–2020. *The Lancet Regional Health—Western Pacific*. 2022; 22: 100433. <https://doi.org/10.1016/j.lanwpc.2022.100433> PMID: 35345391
32. Middleton J. Can ivermectin mass drug administrations to control scabies also reduce skin and soft tissue infections? Hospitalizations and primary care presentations lower after a large-scale trial in Fiji. *The Lancet Regional Health—Western Pacific*. 2022; 22: 100454. <https://doi.org/10.1016/j.lanwpc.2022.100454> PMID: 35462879
33. Engelman D, Marks M, Steer AC, Beshah A, Biswas G, Chosidow O, et al. A framework for scabies control. Taylan Ozkan A, editor. *PLoS Negl Trop Dis*. 2021; 15: e0009661. <https://doi.org/10.1371/journal.pntd.0009661> PMID: 34473725
34. World Health Organization. WHO informal consultation on a framework for scabies control: World Health Organization Regional Office for the Western Pacific: Manila, Philippines, 19–21 February 2019: meeting report. Geneva: World Health Organization; 2020. Available: <https://apps.who.int/iris/handle/10665/333154>
35. Brucker G, Kouchner B, Gentilini M. Rapport sur l'état de santé des populations de Nouvelle Calédonie. Secrétariat d'État auprès du Premier Ministre chargé de l'Action Humanitaire. South Pacific community; 1989 pp. 278–290.