



HAL
open science

Impacts of the antartic ozone hole influence events over southern Brazil in October 2015

Alanna M De Souza, Lucas V Peres, Gabriela D Bittencourt, Damaris K Pinheiro, Bibiana C Lopes, Vagner Anabor, Neusa M.P. Leme, Maria Paulete P. Martins, Rodrigo Da Silva, Gabriela C.G. Dos Reis, et al.

► **To cite this version:**

Alanna M De Souza, Lucas V Peres, Gabriela D Bittencourt, Damaris K Pinheiro, Bibiana C Lopes, et al.. Impacts of the antartic ozone hole influence events over southern Brazil in October 2015. Anais da Academia Brasileira de Ciências, 2023, 95 (suppl 3), 10.1590/0001-3765202320210528 . hal-04237430

HAL Id: hal-04237430

<https://hal.univ-reunion.fr/hal-04237430v1>

Submitted on 11 Oct 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



GEOSCIENCES

Impacts of the antarctic ozone hole influence events over southern Brazil in October 2015

ALANNA M. DE SOUZA, LUCAS V. PERES, GABRIELA D. BITTENCOURT, DAMARIS K. PINHEIRO, BIBIANA C. LOPES, VAGNER ANABOR, NEUSA M.P. LEME, MARIA PAULETE P. MARTINS, RODRIGO DA SILVA, GABRIELA C.G. DOS REIS, MARCO ANTÔNIO G. DOS REIS, JOSÉ V. BAGESTON & HASSAN BENCHERIF

Abstract: The impact of the Antarctic Ozone Hole Influence over Southern Brazil in October 2015 was analyzed using daily mean data of the Total Column Ozone (TCO), Ultraviolet Index (UVI) and Radiative Cloud Fraction (RCF) from the Ozone Monitoring Instrument satellite instrument. Vertical profiles and fields of ozone content and Potential Vorticity available from the European Centre for Medium-Range Weather Forecast reanalysis, air masses backward trajectories from the HYbrid Single-Particle Lagrangian Integrated Trajectory model and channel 3 water vapor images from the Geostationary Operational Environmental Satellite GOES-13 were also analyzed. The five identified events showed an $-7.4 \pm 2.3\%$ average TCO reduction, leading to an $+16.6 \pm 54.6\%$ UVI increase even with a predominance of partly cloudy days. Other impacts were observed in the ozone profiles, where the most significant anomalies occurred from 650 K reaching 1.2 ppmv at the 850 K level. In the ozone fields at 700 K, the presence of a polar origin tongue was observed causing negatives anomalies between -0.2 and 0.4 ppmv in a transient system format forced with eastward-traveling Rossby waves passing through the Southern of Brazil and Uruguay.

Key words: Antarctic Ozone Hole, Southern Brazil, potential vorticity, UV index.

INTRODUCTION

Located approximately between 15 km and 35 km altitude (London 1985), is in the “ozone layer” that photochemical and dynamic processes occur, protecting the Earth and make life possible as it is today (Salby 1996). In the mid-1980s, a massive ozone depletion was discovered during the southern spring over the Antarctic region (Farman et al. 1985), which became known as the “Antarctic Ozone Hole” (Solomon 1999).

The, solar ultraviolet radiation (UVR), corresponds in the electromagnetic spectrum wavelengths from 100 to 400 nm is divided into three groups: UVC (100-280 nm), UVB (280-315 nm) and UVA (315-400 nm) (Liou. 2002; ICNIRP,

2004), being the first completely absorbed in the in the stratospheric ozone layer (Dessler 2000). The UVA type contributes to the synthesis of vitamins (McKenzie et al. 2009), but in excess it can be highly harmful (Oliveira 2014), as well as the type B (UVB), harmful to single-celled organisms and superficial cells of plants and animals, also being partly filtered in the stratosphere (Corrêa 2015). Thus, the UVR is now seen as a public health issue, where different nations carry out actions aimed at reeducating habits of exposure of populations to the sun (WMO 2002).

The ultraviolet index (UVI) has become a measure for determining the intensity of UVR,

first used in Canada in 1992, then adopted as a standard indicator of UVR levels by the World Meteorological Organization (WMO) and the World Health Organization (WHO) in 1994. The total amount of ozone in the atmosphere, cloud cover, snow reflectance, local pollution, and sunrise are the conditions that affect UVI according to a study by Fioletov (2010). Research published by Kirchhoff et al. (2000), Rodriguez (2017) and Sánchez-Pérez et al. (2019) also highlight the importance of this index in cases of sunburn, photodermatoses and even skin cancer, making it possible to determine the sun exposure for each type of skin without suffering strong damage, which can in the long term bring greater effects.

Studies by Kirchhoff et al. (1996) identified from atmospheric balloons (ozone sondes) and surface instruments, such as the Brewer spectrophotometer, that ozone-poor air masses can be ejected from the Antarctic Ozone Hole region (Farman et al. 1985, Solomon 1999, Hassler et al. 2011) and reach mid-latitude regions such as Southern Brazil, in a phenomenon known as the “Antarctic Ozone Hole Secondary Effect” (Pinheiro et al. 2011). This type of phenomenon occurs due to the end of the polar winter, when the increase in temperature causes stratospheric warming and destabilizes the polar vortex, causing an increase in the activity of planetary waves and a break in the Rossby wave (Ndarana et al. 2012), and with that, ejections of these ozone-poor air masses out of the polar vortex, in the form of towards the equator filaments (Schoeberl 1988).

Several studies on a regional scale have begun to observe the occurrence of the Antarctic Ozone Hole Influence events over Southern Brazil, motivated by studies on the air masses isentropic transport in the lower stratosphere and your ability to determine the ozone content in different regions of the planet (Portafaix et al.

2003, Semane et al. 2006, Gettelman et al. 2011). Between 2008 and 2012, was identified fourteen this type events, with an average reduction of $9.3 \pm 2.9\%$ in the TCO (Pinheiro et al. 2011, 2012, Peres et al. 2016). In addition, this events type is one of the explanations for the amphibian species decline over the Southern Brazil (Schuch et al. 2015) and Steffanel et al. (2016) detected these events using Pervasive Computing in order to carry out alerts about the ozone reductions that cause increases in UV radiation.

Dias Nunes et al. (2020), analyzed the impact of ozone content on UVR variability in South America between 2005 and 2014. This analysis proved that only low ozone levels are not determinant for high UVR values. This inverse correlation being seasonally dependent and more intense between July and October over large areas of South America. Cloud cover stands out among the factors that also influence in UVR quantities over South America. This helping to explain the differences between observations obtained by satellite equipment and reconstructed by models using measurements from ground equipment over Chile (Damiani et al. 2014) and the observed UVR attenuation on stations in Peru compared to those compared to numerical simulations carried out for clear sky conditions (Yamamoto et al. 2018).

Peres et al. (2017) analyzed the climate variability in total column ozone (TCO) measurements obtained by Brewer Spectrometer, and Total Ozone Mapping Spectrometer (TOMS) and Ozone Monitoring Instrument (OMI) satellite instruments between 1992 and 2014 over southern Brazil. Seasonal variability presents a minimum in April and a maximum in September and the Quasi-Biennial Oscillation (QBO) is the main mode of interannual variability. This work served as a climatological basis for works such as Bresciani et al. (2018), that using a multi-instrumental and air parcel trajectories from

HYSPLIT (HYbrid Single-Particle Lagrangian Integrated Trajectory) model analysis to observe a large area of critical ozone depletion over Rio Grande do Sul and Uruguay in mid-October 2016. Bittencourt et al. (2018) also used this climatological basis and model trajectories analysis to identified that this event was the most intense recorded in the last 20 years and carried out an analysis of the tropospheric and stratospheric dynamics around your occurrence period.

The long-term monitoring of this phenomenon type conducted by Peres et al. (2019) covered the period between 1979 and 2013 and identified the occurrence of 62 events with -9.35 ± 2.93 % of average reduction in ozone content. Such events were caused by anomalies in the potential vorticity, wind and temperature fields at isentropic levels of the stratosphere, where the observed wave displacement observed in Potential Vorticity (PV) fields, is within of cyclonic circulation area and with predominantly southern winds. The motivation for the present work is that, for the year 2015, events of Influence of the Antarctic Ozone Hole over Southern of Brazil have not yet been identified. The year 2015 have a gap in the identification of this events. Thus, the present work aims to identify and characterize the Influence of the Antarctic Ozone Hole over Southern of Brazil events that occurred in October 2015 and thus collaborate to analyze the impacts in terms of the ozone depletion in multilevels and the ultraviolet index.

MATERIALS AND METHODS

In this work, Total Column Ozone (TCO), Ultraviolet index (UVI) and Radiative Cloud Fraction (RCF) daily data were obtained from the Ozone Monitoring Instrument (OMI) sensor, that was launched in July 2004 aboard NASA (National

Aeronautics and Space Administration) AURA satellite to continued the TOMS (Total Ozone Mapping Spectrometer) instrument recordings that ended yours activities in 2005. The OMI instrument employs hyperspectral imaging in a scanning mode to observe backscattered solar radiation in the electromagnetic spectrum ranges between 0.270nm to 0.314nm and 0.306nm to 0.380nm. The hyperspectral capabilities improve the accuracy of TCO measures and allow radiometric self-calibration of precise and long term wavelengths (Levelt et al. 2006).

The daily TCO, UVI and RCF data acquisition was performed for the central region of the state of Rio Grande do Sul, on a $1^\circ \times 1^\circ$ latitude / longitude grid, comprising the surroundings of the Santa Maria city (29.72°S ; 53.72°O), as shown in Figure 1. The satellite passage is close to local midday and corresponding to days with all sky conditions, so not only clear sky days. Both the daily TCO (OMTO3d v003 product), UVI (OMUVBd v003 product) and RCF(OMTO3d v003 product) data are available on the NASA Giovanni platform (Acker & Leptoukh 2007), which is an online environment for visualization and preliminary analysis of geophysical parameters and data download (<https://giovanni.gsfc.nasa.gov/giovanni/>).

The identification of the Antarctic Ozone Hole Influence events over Southern Brazil start in select days in which the value of TCO was below the value of the mean minus 1.5 of its respective standard deviation in the month of October 2015, following the criterion detailed by Peres et al. (2019). For this, the climatology of the TCO between 1992 and 2014 calculated by Peres et al. (2017) was used, being 290.2 DU (Dobson Unit) the october climatological value, 8.8 DU its respective standard deviation and 277.0 DU the threshold value of climatological mean minus 1.5 of the standard deviation. In addition, the evolution of UVI and RFC daily data were also




Figure 1. Santa Maria city location, in the central area of Rio Grande do Sul State, Brazil.

analyzed in order to verify our impact during the mentioned TCO reductions.

For the selected TCO reduction days, an stratospheric dynamics analysis was performed from meteorological fields of Potential Vorticity (PV), wind magnitude (u and v) and Ozone mass mixing ratio, obtained for 15 potential temperature levels between 265 and 850 Kelvin, as similar used by Dos Reis et al. (2021) in identifying the Antarctic Ozone Hole Influence events over Southern Brazil. The ERA-Interim Pressure and Isentropic (Theta) Levels in Relation to a Standard Atmosphere can be verified in https://rda.ucar.edu/datasets/ds627.1/docs/Pressure_and_isentropic_levels/, as a function of the potential temperature equation, showing that the maximum in the ozone layer occurs at 25Km or 25.492 hPa or 634.5K. These meteorological fields are provided by ERA - Interim reanalysis from European Centre for Medium-Range Weather Forecast (ECMWF), detailed in Dee et al. (2011), for a $1.0^{\circ} \times 1.0^{\circ}$ latitude / longitude resolution grid.

This stratospheric dynamics analysis starts with the observation of the ozone vertical profiles, comparing the 1986 to 2015 October average ozone profile with the daily average ozone profiles and respective anomaly calculations

(1986 to 2015 October average ozone profile minus daily average ozone profile), in order to verify the height at this ozone reductions occur. At the height of observed ozone reduction, PV fields from the previous days for the day in question were analyzed in order to determine the origin of the stratospheric air masses that caused such reductions, similar to that performed by Bittencourt et al. (2018, 2019), in accordance with described in Semane et al. (2006), where a stratospheric air mass has a polar origin when a reduction in PV values is verified.

Complementing this analysis, the ozone mass mixing ratio fields from ERA - Interim reanalysis and air masses back trajectories obtained by the HYSPLIT (HYbrid Single-Particle Lagrangian Integrated Trajectory) model (Rolph et al. 2017) were also analyzed in order to confirm the air masses polar origin that caused the observed ozone reductions, thus determining the occurrence of the Antarctic Ozone Hole Influence events over Southern Brazil, according to Bresciani et al. (2018).

In order to verify the presence and type of cloudiness and your impact on the observed UVI value, channel 3 water vapor images from the Geostationary Operational Environmental Satellite GOES-13, obtained from the Satellites

and Environmental Systems Division of the Center for Weather Forecasting and Climate Studies (CPTEC / INPE 2021) were also analyzed.

Finally, the impact of the October 2015 occurrence events was quantified in relation to the 1986 to 2015 October period behavior average. For this, the average ozone profile for this period (1986 to 2015) was compared with the average ozone profile of the days considered to be events in October 2015 and the difference between them was calculated, presenting the preferred levels of ozone reduction. In these levels, the ozone average fields of the period 1986 to 2015, average ozone field of the events and the anomaly (Wilks 2011) between these were analyzed.

RESULTS AND DISCUSSION

The daily TCO, UVI and RCF data for October 2015 are presented in the Figure 2. In function your 114° viewing angle range of the telescope, that corresponds to a 2600 km wide swath

on the Earth's surface (Levelt et al. 2018), OMI measurements can present systematic observed gaps as observed in Figure 2. The OMI satellite daily TCO values (blue) and the $-1, 5\sigma$ (277 DU) threshold value (black starred line) for October 2015 are presented in the Figure 2a. From thirty-one days, sixteen (51.6%), were observed below the -1.5σ threshold, being considered of TCO reduction, which were verified by the ozone profiles and atmospheric dynamics analysis in order to search of the Antarctic Ozone Hole Influence events over southern Brazil. The search for days in which the TCO was below the -1.5σ threshold resulted in the identification of two Antarctic Ozone Hole Influence events over southern Brazil in 2012 by Peres et al. (2016), 62 events in the 1979 to 2013 period by Peres et al. (2019) including the major event in October 20, 2016, by Bittencourt et al. (2018), which caused 23% TCO reduction.

In the Figure 2b, the daily UVI values (blue) and the 7.6 UVI October 2015 average value (black starred line) are presented. It is observed




Figure 2. (a): Daily TCO values (blue) and the -1.5σ threshold value (black starred line). (b) Daily UVI values (blue) and UVI October 2015 average (starred black line). (c) Daily RCF values (blue) and RCF October 2015 average (starred black line).

that 45.2 % of the days were above the referred monthly average. Analyzes over Southern Brazil detected that 1% ozone variations can cause 0.94% to 1.36% UV variations Guarnieri et al. (2004), being this type of correlation already well established as mentioned by Calbó et al. (2005). The Figure 2c present the daily RCF values (blue) and the 0.44 RCF October 2015 average value (black starred line), on what 38.7 % of the days were above this monthly average. Since clouds can exert great influence on UV radiation, how to reduce its transmittance by 81% in high clouds conditions over Argentina (Cede et al. 2002), analyzing your occurrence as a function of ozone depletion events and UV variations becomes important.

The October 29, 2015, stands out in this analysis for having 271.1 DU TCO value, representing a 7.1% ozone reduction in relation to the 290.2 DU October climatological value (Peres et al. 2017), the highest UVI value (11.8)

for October 2015, representing +53.9 UVI increase and resent clear sky (0 RCF) condition. The sum of these factors makes this event the best example to demonstrate the methodology used to identify the influence of the Antarctic Ozone Hole over Southern Brazil in october 2015.

The vertical ozone profile for the present day (red line), obtained from the ECMWF ERA - Interim reanalysis, was compared whit 1986 to 2015 october average ozone profile (black line) in order to verify what level occur the major ozone reduction (Figure 3a), by calculating the anomaly between them (Figure 3b).

Was observed in the Figure 3a, that on average, the highest values begin to occur from the 450 K potential temperature level, when they begin to exceed 2 ppmv, reaching their maximum at 850 K when it reaches nearly 9 ppmv. These values are in agreement with the mean values observed for 30°S latitude by Davis et al. (2017), using SPARC (Stratosphere-troposphere




Figure 3. (a) 1986 to 2015 october average ozone profile (black line) and October 29, 2015 ozone profile (red line). (b) October 29, 2015 ozone anomaly profile (red line) relative to zero (black line). Both obtained from the ECMWF ERA-INTERIM Reanalysis in ppmv.

Processes and their Role in Climate) Reanalysis data sets. From 650 K potential temperature level to up, ozone reductions were identified, region where the October 29, 2015 ozone profile (red line) has values below of the 1986 to 2015 October average ozone profile (black line). In order to quantify this reduction, the anomaly between them was calculated (Figure 3b) and no significant anomalies are observed from initial to around 400 K potential temperature levels. Between 400 and 500 K levels, ~ 0.2 ppmv negative anomalies are observed and more than 0.4 ppmv positive anomaly is observed in 600 K level.

These levels are in contrast with the levels above 600 K, where intense ozone reductions are evident, with a maximum negative anomaly close to 1.4 ppmv at 850 K level are observed. From the 700K potential temperature level, approximately 24 km and the average ozone layer level at mid-latitudes (Solomon 1999), the potential vorticity maps were analyzed in order

to perform the dynamics analysis and verify the origin of the air masse that caused this ozone reduction (Figure 4). The 700K potential temperature level is often used in studies about ozone isentropic transporte by atmospheric dynamics as performed by Portafaix et al. (2003) and Semane et al. (2006) in the stratospheric South Hemisphere.

With the beginning of SH winter, increase the stratospheric circumpolar winds speed, delineating the barrier between polar and mid-latitude air called "Polar Vortex"(Mizuta & Yoden 2001) with your edge positioned around 60°S (Joseph & Legras 2002). The area of the Antarctic Ozone Hole is characterized by the polar vortex edge positioning (Shepherd 2007), which is determined by the high potential vorticity gradient region (Nash et al. 1996, Serra & Haller 2017). Air masses advection processes, induced by planetary Rossby waves (McIntyre & Palmer 1984), can result in polar filaments, which detach from the edge of the polar vortex




Figure 4. Potential Vorticity and wind fields in 700 K isentropic potential temperature level signaling the air masses stratospheric dynamics from October 27 (a), 28 (b) and 29 (c), 2015 at 12 UTC.

and reach mid-latitudes and reduce the ozone content over these regions (Schoeberl et al. 1995).

So, in order to diagnose the stratospheric dynamics that caused these observed ozone reduction, the 700K PV maps from October 27 to 29, 2015 were analysed in Figure 4. In October 27, 2015 (a), was observed a zonal behavior of PV, with values between -60 and -80 PVU over the central region of the Rio Grande do Sul state (marked with a triangle). The polar origin air masses advection starts to be observed over the Uruguay in October 28, 2015 (b), with lower PV values inside an elongated system as a cyclonic circulation tongue. Southern Brazil begins suffer advection from prevailing southerly winds, reducing their PV values to -80 to -100 PVU.

On October 29, 2015 (c), arrive the polar origin air mass over Southern Brazil on the rear of the cyclonic circulation system, carried by an amplification of anticyclonic circulation system. Thus, the stratospheric fields indicated a PV decrease that reach values between -100 to -120 PVU in the 700 K level of potential temperature, evidencing the polar origin of the air mass in a manner analogous to that verified by Leblanc et al. (2004), that identified large transient displacements of stratospheric air masses causing ozone variability through advection analysis in potential vorticity fields at isentropic levels, accompanied by Rossby wave breaking events.

High degree of correlation between PV and ozone that can be observed from 340 to above 700 K isentropic level, responsible for negative/positive ozone deviations coincident with transport from regions with climatologically low/high ozone values (Koch et al. 2002). Similar behavior of a transient system, whit alternation of wind dynamic system from a cyclonic circulation to an anticyclonic circulation

system, preferentially coming from the south, transported polar origin lower PV and ozone values over Southern Brazil over the 700 K isentropic level as observed in the Figures 4 and 5.

Similar to the PV field in Figure 4, the 700K ozone field and the NOAA's HYSPLIT model backward trajectory at the 20, 24 and 26 km heights in the Figure 5, presents an air masses polar origin advection process that reduced the ozone content over southern Brazil. In October 27, 2015 (a) and (d), a zonal behavior of air masses over Southern Brazil is observed, with 7 to 6.65 ppmv ozone values. A polar tongue air mass is observed in the 700K ozone field over de Uruguay in October 28, 2015 (b), which begins to impact the ozone content over the Santa Maria region, reducing it to between 6.65 and 6.3 ppmv. However for that day, the HYSPLIT backward trajectory (e) still not point to the air mass passing inside the polar circle.

The southern air mass advection process continues and in October 29, 2015 (c) the polar tongue reaching Southern Brazil. This causes the ozone depletion to 6.3 and 5.95 ppmv and is evidenced that the air mass pass over the polar circle in the HYSPLIT backward trajectory (f), configuring the occurrence of Influence of the Antarctic Ozone Hole over Southern Brazil in this date, similarly to observed by Bresciani et al. (2018) and Bittencourt et al. (2018) for other events of this type. These strengthening of ridge-related to negative PV and ozone anomalies in stratospheric levels is associated with Rossby wave breaking events, here diabatic heating reduces the static stability near the tropopause and contributes to this process (Zhang & Wang 2018).

The present ozone reductions observed in the Antarctic Ozone Hole Influence event over Southern Brazil on October 29, 2015 also presented an 11.7 UVI value, representing a




Figure 5. Era Interim 700 K daily ozone average fields (in ppmv) and NOAA's HYSPPLIT model air masses backward trajectory for October 27 (a and d), 28 (b and e) and 29 (c and f) at 12 UTC.

53.9% increase in relation to 7.6 UVI October 2015 average value. This IUUV value is considered very high according to the criteria of the World Health Organization (WHO 2002), which recommends extra protection to the population, such as avoiding being outdoors during the midday, seeking shade and essential use of long sleeve shirt, sunscreen and cap. This high observed UVI value was favored due to clear sky condition in this day, verified by 0 RCF value and in the channel 3 water vapor images from the Geostationary Operational Environmental Satellite GOES-13 (Figure 6), in both hours of high brightness of the day as 12 UTC (a), 15 UTC (b) and 18 UTC (c), not attenuating the present radiation that reached the surface in the central region of the Rio Grande do Sul state.

Analyzing this sequence of satellite images in Figure 6, it is possible to observe that the synoptic characteristics in October 29, 2015 points to a stable post-frontal air mass acting over southern Brazil, without significant cloudiness.

This is related to the presence of subsidence region of the polar jet and the post-frontal high pressure system, which advects cold air, that discouraging the cloudiness formation over this region. This postfrontal synoptic condition pattern has already been observed by Peres et al. 2011 and 2014 in other events of Influence of the Antarctic Ozone Hole over Southern Brazil in 2008 and 2012 respectively.

Table I shows all five events of Influence of the Antarctic Ozone Hole over Southern Brazil in October 2015. On average, this events presented 258.7 ± 6.6 DU TCO values, $7.4 \pm 2.3\%$ ozone rediction in relation to October climatological value, 8.8 ± 4.1 UVI value that representing 16.6% UVI increases in relation to 2015 October average and 0.25 ± 0.42 RCF value, representing that on average the events occur on partially cloudy days.

The cloudiness analysis in each event is presented and observed that clear sky days (0 RCF) predominates in both events with the




Figure 6. Channel 3 Water vapor images from the Geostationary Operational Environmental Satellite GOES-13 of the Environmental Satellites and Systems Division of the Center for Weather Forecasting and Climate Studies for 12 UTC (a), 15 UTC (b) and 18 UTC (c) on October 29, 2015.

highest UVI value in October 29 (11.7 UVI and +53.9% UVI variation) and 24 (10.9 UVI and +43.4% UVI variation) 2015, despite ozone reductions in relation to October climatology are not as deep, -6.54% and 5.89% respectively. Partly Cloudy events as in October 16 and 20, 2015 showed the same 10.1 UVI values despite of different ozone reductions (-11.1% and -5.37%) and cloud cover (0.18 and 0.11 RCF). October 8, 2015, showed Cloudy sky condition (1 RCF), resulting in low UVI value (1.53) despite relatively deep ozone depletion (-8.1%).

In 37 events of Influence of the Antarctic Ozone Hole over Southern Brazil investigated by Bittencourt et al. (2019), the atmospheric behavior was predominant with 70% of cases occurring after the passage of frontal systems and 92% occurred in the presence of the subtropical and/or polar jet stream over the region of study. However, even if the passage of ozone-poor air masses can cause increase in UV

radiation to in relation to expected for a specific time of year, factors such as aerosols and clouds can cause decrease these levels (Palancar & Toselli 2004, Utrillas et al. 2018), corroborating with the results found in this analysis.

The effect of cloudiness on ultraviolet radiation has been subject of several studies and helps to explain the results found here. Mayer et al. (1998) found experimental and theoretical evidence explaining the UV radiation increase due to multiple scattering in clouds, showing that the paths of photons in clouds can be increased by a factor of 10 compared to the cloudless sky, this finding being of great importance in remote sensing applications that take advantage of the measurement of scattered radiation to infer the abundance of atmospheric trace gases. Cumulus clouds, common in frontal systems, can cause not only deep drops, but also remarkable increases in radiation levels, which constitute the so-called broken cloud

Table I. Antarctic Ozone Hole Influence events over Southern Brazil in October 2015. Daily average TCO value, percentage of the TCO variation from the October climatological average, daily UVI value, the percentage of UVI variation from the October 2015 average, daily RCF value and cloudiness.

Events	TCO(UD)	Var TCO (%)	UVI	Var UVI (%)	RCF	Cloudness
08/10/2015	266.7	-8.1	1.53	-79.8	1	Cloudy sky
16/10/2015	258.1	-11.1	10.1	+32.9	0.18	Partly Cloudy
20/10/2015	274.6	-5.37	10.1	+32.9	0.11	Partly Cloud
24/10/2015	273.1	-5.89	10.9	+43.4	0	Clear sky
29/10/2015	271.2	-6.54	11.7	+53.9	0	Clear sky
Mean	258.7±6.6	-7.4±2.3	8.8±4.1	16.6±54.6	0.25±0.42	Partly Cloud

effect, in function that fragmented clouds are often brighter, leading to a radiation gain. (Nack & Green 1974).

Even in high places like Peru, where clear sky conditions register high UVI values, adverse weather conditions such as strong presence of clouds and precipitation, which block part of the solar radiation mainly in summer and autumn, with the highest UVI average only in winter where it presents less cloudiness (Yamamoto et al. 2018). Feister et al. (2015) point out that the scattering of solar radiation from clouds can affect the surface energy balance and obtain solar energy gains for power generation. In addition, in the ultraviolet region, cloudiness can cause an damage increased risk to living organisms, and the role of clouds is the main atmospheric agent that causes UVR variability (McKenzie et al. 2007). They are more transparent to UVR than to visible radiation (400 to 780 nm) meaning that their presence, even covering 100% of the sky, is not a guarantee of no harmless radiation levels (Seckmeyer et al. 1996).

Clouds generally reduce surface UV irradiances, although the magnitude of this effect is highly variable depending on cloud amount and coverage, cloud cell morphology, particle size distributions and phase (water droplets and ice crystals), and possible in-cloud absorbers

(especially tropospheric ozone). It is useful to note that under some conditions, UV irradiances can be higher than for clear sky, as for example when both direct sunlight and light scattered by clouds reach the observer (Madronicha et al. 1998). Jesus (2015) found that UV radiation in the presence of clouds interacts as follows: When clouds are optically thin, it transforms direct irradiance into diffuse irradiance. When clouds are thicker, multiple scattering stands out, extinguishing surface irradiances and increasing upward radiances.

In order to quantify the impact of the Antarctic Ozone Hole influence on ozone profile over Southern Brazil in relation to isentropic levels during October 2015, the average ozone profile of the five events presented in Table I was calculated and analyzed in comparison to the 1986 to 2015 October average profile (Figure 7a) and the anomaly profile of the events was also calculated (Figure 7b).

In Figure 7a, is observed that the levels below 500 K can contain three times less ozone than the levels above 650 K and that in the October 2015 events average, negative anomalies in ozone content were observed between 400 and 550K, with maximum in 475K near -0.4 ppmv and above 650K with maximum in 850K near to -0.9 ppmv (Figure 7b). Below 400 K the ozone




Figure 7. (a) 1986 – 2015 October average ozone profile (black line) and the October 2015 events average ozone profile (red line). (b) October 2015 average events ozone anomaly profile (red line) relative to zero (black line). Both obtained from the ECMWF ERA-INTERIM Reanalysis in ppmv.

reductions are close to 0 ppmv and between 550 K and 650 K positive anomalies are observed with maximum in 600 K near to +0.4 ppmv. This magnitude of ozone anomaly in vertical profiles is consistent with variability and trend studies performed by Kiesewetter et al. (2010).

Since 24 km height is closer to the mean ozone layer level at mid-latitudes, and shows significant negative ozone anomaly, the 700 K potential temperature level was selected to ozone anomalies verification in ozone concentration fields. The isentropic transport of polar origin air masses toward Southern Brazil can be quantified by analyzing the Figure 8, which presents the 1986 to 2015 October average ozone field (a), the October 2015 events average ozone field (b), and the October 2015 events anomalies (c), both for the 700 K isentropic level.

In the 1986 to 2015 October average ozone field (a), the presence of a zonal ozone gradient values is observed, with lower values positioned

toward the pole and higher values toward the equator, with the central region of the state of Rio Grande do Sul having values between 6.65 and 7 ppmv. In the October 2015 events average ozone field, the propagation of a synoptic scale wave disturbance is observed, leading to northward the lines of lower values of this ozone gradient over the Atlantic Ocean, which resulted in the transport of a lower ozone values polar origin tongue toward the central region of Rio Grande do Sul state, which was left with values between 6.3 and 6.65 ppmv (b). This lower ozone values polar origin tongue, caused anomalies in relation to the 1986 – 2015 October average between -0.2 and -0.4 ppmv that extend from Pacific Ocean, passing through to North Chile and Argentina, Southern Brazil and Uruguay and go to the Southeast Atlantic Ocean.

This transient system format extending from the Pacific Ocean to the Atlantic Ocean, passing through Southern Brazil and Uruguay




Figure 8. 1986 to 2015 October average ozone field (a), the October 2015 events average ozone field (b), and the October 2015 events anomalies (c). Both for the 700 K isentropic level.

from polar origin is similar to traveling disturbances forced with eastward-traveling Rossby waves identified by Guha et al. (2016). This transient systems have a live period of 7–8 days and have been observed in the Southern Hemisphere during winter/springtime (Hio & Yoden 2004). This period of time is enough for the air masses to move from polar regions and reach medium latitudes as observed in the studies of ozone and PV simulations on the HS carried out by Marchand et al. (2005) and Hauchecorne et al. (2002). Studies by Semane et al. (2006) also observed that isentropic transport of ozone in the stratosphere caused an unusual ozone reduction in the Southern Hemisphere subtropics, more precisely over Irene (25.5 °S, 28.1 °E) in mid-May 2002 and demonstrate the importance of the stratosphere dynamics impact studies during the Antarctic Ozone Hole Influence over Southern Brazil events occurrence for longer time periods.

CONCLUSIONS

The impact of the Influence of the Antarctic Ozone Hole over Southern Brazil events in October 2015 was analyzed in the present study. Was found that 51.6% of the days has ozone reduction below the -1.5σ limit and with UVI values above the October 2015 average, resulting in five events identified.

As an example of this type phenomenon identification, the event occurred in October 29, 2015 was presented. This event has a 6.54% TCO reduction and a 53.9% UVI increase with a clear sky day, intense reductions in ozone content above 600 K levels which reached close to 1.4 ppmv at the 850 K level, caused by an air mass polar origin tongue, surrounded by a cyclonic circulation with predominantly southerly winds. This type of analysis was effective to identifying five events in October 2015, with $-7.4 \pm 2.3\%$ TCO reduction which impacted an $+16.6 \pm 54.6\%$ UVI

increase with a predominance of partly cloudy days.

In addition, other impacts of these events were observed in ozone profiles that had negative anomalies at all levels, being more significant from 650 K where reached 1.4 ppmv at 850K. In 700K ozone fields, the isentropic transport pointed to the presence of lower ozone values polar origin tongue that caused anomalies between -0.2 and -0.4 ppmv extending from extend from Pacific Ocean, passing through to North Chile and Argentina, Southern Brazil and Uruguay and go to the Southeast Atlantic Ocean, with transient system format similar to traveling disturbances forced with eastward-traveling Rossby waves, demonstrating the importance of conducting this type of study for longer periods of time. With a better understanding of this phenomenon type and your future forecast, will be possible to inform the affected populations in advance, so that they can follow the recommendations established by the World Health Organization.

Acknowledgments

This work is a Bachelor in Atmospheric Sciences course at the Federal University of Western Pará product, with financial support from the Institutional Scientific Initiation Scholarship Program (PIBIC-UFOPA). The authors express their thanks to the MESO Project - Modeling and Prediction of the Secondary Effects of the Antarctic Ozone Hole, linked to the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES/COFECUB), Process No. 88887.130199/201701 and Instituto Nacional de Ciência e Tecnologia Antártico de Pesquisas Ambientais, Conselho Nacional de Desenvolvimento Científico e Tecnológico - CNPq process no. 574018/2008-5 and Fundação de Amparo à Pesquisa do Estado do Rio de Janeiro - FAPERJ process no. E-16/170.023/2008. They also thank NASA for the OMI satellite data, ECMWF/ERA-Interim for the daily ozone and PV data, and NOAA for the products generated through the HYSPLIT model.

REFERENCES

- ACKER JG & LEPTOUKH G. 2007. Online analysis enhances use of nasa earth science data. *Eos, Trans AGU* 88: 14-17.
- BITTENCOURT GD, BRESCIANI C, KIRSCH PINHEIRO D, BAGESTON JV, SCHUCH NJ, BENCHERIF H & VAZ PERES L. 2018. A major event of antarctic ozone hole influence in southern brazil in october 2016: an analysis of tropospheric and stratospheric dynamics. *Ann Geophys* 36(2): 415-424.
- BITTENCOURT GD, PINHEIRO DK, BAGESTON JV, BENCHERIF H, STEFFENEL LA & VAZ PERES L. 2019. Investigation of the behavior of the atmospheric dynamics during occurrences of the ozone hole's secondary effect in southern brazil. *Ann Geophys* 37: 1049-1061.
- BRESCIANI C, BITTENCOURT GD, BAGESTON JV, PINHEIRO DK, SCHUCH NJ, BENCHERIF H & PERES LV. 2018. Report of a large depletion in the ozone layer over southern brazil and uruguay by using multi-instrumental data. *Ann Geophys* 36(2): 405-413.
- CALBÓ J, PAGÈS D & GONZÁLEZ J-A. 2005. Empirical studies of cloud effects on UV radiation. *A review Rev Geophys* 43: 2002-2005.
- CEDE A, LUCCINI E, PIACENTINI RD, NUNEZ L & BLUMTHALER M. 2002. Monitoring of Erythemat Irradiance in the Argentine Ultraviolet Network. *J Geophys Res Atmos* 107:4165-4175.
- CORRÊA MP. 2015. Solar ultraviolet radiation: properties, characteristics and amounts observed in Brazil and South America. *An Bras Dermatol* 90(3): 297-313.
- CPTEC/INPE. 2021. Division of Satellites and Environmental Systems. Accessed 10 Mar. 2021. <http://satellite.cptec.inpe.br/acervo/goes.formulario.logic>.
- DAMIANI A, CORDERO RR, CABRERA S, LAURENZA M & RAFANELLI C. 2014. cloud cover and UV index estimates in Chile from satellite-derived and ground-based data. *Atmos Res* 138: 139-151. <https://doi.org/10.1016/j.atmosres.2013.11.006>.
- DAVIS SM ET AL. 2017. Assessment of upper tropospheric and stratospheric water vapor and ozone in reanalyses as part of S-RIP. *Atmos Chem Phys* 17: 12743-12778. <https://doi.org/10.5194/acp-17-12743-2017>.
- DEE DP ET AL. 2011. The era-interim reanalysis: configuration and performance of the data assimilation system. *Q J R Meteorol Soc* 137: 553-597.
- DESSLER AE. 2000. The chemistry and physics of stratospheric ozone, academic, San Diego, California 1: 214.
- DIAS NUNES M, MARIANO GL & ALONSO MF. 2020. Spatio-temporal variability of total ozone column and ultraviolet radiation: assessment of relationship in South America.

- Brazilian Journal of Physical Geography 13(05): 2053-2073. DOI: <https://doi.org/10.26848/rbgf.v13.5.p2053-2073>.
- DOS REIS MAG, PERES LV, BITTENCOURT GD, PINHEIRO DK, STEFFENEL LA, BENCHERIF H, SILVA R, NUNES MD & BAGESTON JV. 2021. Eventos de Influência do Buraco de Ozônio Antártico Ocorridos em 2016 Sobre o Sul do Brasil. Anuário do Instituto de Geociências 44: 35438. https://doi.org/10.11137/1982-3908_2021_44_36142.
- FARMAN JC, GARDINER BG & SHANKLIN JD. 1985. Large losses of total ozone in antarctica reveal seasonal clox/nox interaction. *Nature* 315: 207-210.
- FEISTER U, CABROL N & HÄDER D. 2015. UV Irradiance Enhancements by Scattering of Solar Radiation from Clouds. *Atmosphere* 6: 1211-1228.
- FIOLETOV V, KERR JB & FERGUSON. 2010. A. The UV Index: Definition, Distribution and Factors Affecting It. *Can J Public Health* 101: 15-19.
- GETTELMAN A, HOOR P, PAN LL, RANDEL WJ, HEGGLIN MI & BIRNER T. 2011. The extratropical upper troposphere and lower stratosphere. *Rev Geophys* 49: 267-268.
- GUARNIERI RA, PADILHA LF, GUARNIERI FL, ECHER E & MAKITA K, PINHEIRO DK, SCHUCH AMP, BOEIRA LS & SCHUCH NJ. 2004. Study of the anticorrelations between ozone and uv-b radiation using linear and exponential fits in southern Brazil. *Adv Space Res* 34: 764-768.
- GUHA A, MECHOSO CR, KONOR CS & HEIKES RP. 2016. Modeling Rossby Wave Breaking in the Southern Spring Stratosphere. *J Atmos Sci* 73: 393-406.
- HAUCHECORNE A, GODIN S, MARCHAND M, HEESE B & SOUPRAYEN C. 2002. Quantification of the transport of chemical constituents from the polar vortex to midlatitudes in the lower stratosphere using the high-resolution advection model mimosa and effective diffusivity. *J Geophys Res Atmos* 107: 1-13.
- HASSLER B, DANIEL JS, JOHNSON B J, SOLOMON S & OLTMANS SJ. 2011. An assessment of changing ozone loss rates at south pole: twenty-five years of ozonesonde measurements. *J Geophys Res-Atmos* 116: 1-12.
- HIO Y & YODEN S. 2004. Quasi-periodic variations of the polar vortex in the Southern Hemisphere stratosphere due to wave-wave interaction. *J Atmos Sci* 61: 2510-2527. doi:10.1175/JAS3257.1.
- ICNIRP - INTERNATIONAL COMMISSION OF NON-IONIZING RADIATION PROTECTION. 2004. Guidelines on limits of exposure to ultraviolet radiation of wavelength between 180 nm and 400 nm (incoherent optical radiation). *Health Physics* 87: 171-186.
- JESUS HS. 2015. Estudo teórico e observacional dos efeitos da nebulosidade na radiação ultravioleta. Dissertação de Mestrado. Instituto Nacional de Pesquisas Espaciais – INPE. (Unpublisehd).
- JOSEPH B & LEGRAS B. 2002. "Relation between kinematic boundaries, stirring, and barriers for the Antarctic polar vortex." *J Atmos Sci* 59(7): 1198-1212.
- KIESEWETTER G, SINNHUBER BM, VOUNTAS M, WEBER M & BURROWS JP. 2010. A long-term stratospheric ozone data set from assimilation of satellite observations: high-latitude ozone anomalies. *J Geophys Res Atmos* 115(D10).
- KIRCHHOFF VWJH, ECHER E, LEME NP & SILVA AA. 2000. A variação sazonal da radiação ultravioleta solar biologicamente ativa. *Rev Bras Geof* 18(1): 63-74.
- KIRCHHOFF VWJH, SCHUCH NJ, PINHEIRO DK & HARRIS JM. 1996. Evidence for an ozone hole perturbation at 30° south. *Atmos Environ* 33(9): 1481-1488.
- KOCH G, WERNLI H, STAEHELIN J & PETER T. 2002. A Lagrangian analysis of stratospheric ozone variability and long-term trends above Payerne (Switzerland) during 1970 – 2001. *J Geophys Res* 107: 2-14. doi:10.1029/2001JD001550.
- LEBLANC T, MCDERMID IS & HAUCHECORNE A. 2004. A study of ozone variability and its connection with meridional transport in the northern Pacific lower stratosphere during summer 2002. *J Geophys Res* 109: D11105. doi:10.1029/2003JD004027.
- LEVELT P, OORD GHJ, DOBBER M, MÄLKKI A, VISSER H, VRIES J, STAMMES P, LUNDELL J & SAARI H. 2006. The ozone monitoring instrument, *IEEE T. Geosci Remote Sens* 44: 1093-1101.
- LEVELT PF ET AL. 2018. The Ozone Monitoring Instrument: overview of 14 years in space. *Atmos Chem Phys* 18: 5699-5745. <https://doi.org/10.5194/acp-18-5699-2018>.
- LIU KN. 2002. An introduction to atmospheric radiation, *International Geophysics Series*, Academic Press 84: 583.
- LONDON J. 1985. Observed distribution of atmospheric ozone and its variations. In: Whitten RC & Prasad SS (Eds). *Ozone in the free atmosphere*. New York: Van Nostrand Reinhold. Cap. 1, p. 11-80.
- MADRONICH S, MCKENZIE RL, BJÖRN BLO & CALDWELL MM. 1998. Changes in biologically active ultraviolet radiation reaching the Earth's surface. *J Photochem Photobiol B, Biol* 46(1-3): 5-19.
- MARCHAND M, BEKKI S, PAZMINO A, LEFÈVRE F, GODIN-BEEKMANN S & HAUCHECORNE A. 2005. Model simulations of the impact of the 2002 antarctic ozone hole on the midlatitudes. *J Atmos Sci* 62: 871-884.

- MAYER B, KYLLING A, MADRONICH S & SECKMEYER G. 1998. Enhanced absorption of UV radiation due to multiple scattering in clouds: Experimental evidence and theoretical explanation. *J Geophys Res* 103: 31.241-31.254.
- MCINTYRE ME & PALMER TN. 1984. The surf zone in the stratosphere, *J Atmos Terr Phys* 9: 825-849.
- MCKENZIE RL, AUCAMP PJ, BAIAS AF, BJORN LO & ILYAS M. 2007. Changes in biologically-active ultraviolet radiation reaching the Earth's surface. *Photochem Photobiol Sci* 6: 218-231.
- MCKENZIE RL, LILEY JB & BJÖRN LO. 2009. UV radiation: balancing risks and benefits. *Photochem Photobiol* 85: 88-98.
- MIZUTA R & YODEN S. 2001. Chaotic mixing and transport barriers in an idealized stratospheric polar vortex. *J Atmos Sci* 58: 2616-2629. [https://doi.org/10.1175/1520-0469\(2001\)058<2616:CMATBI>2.0.CO;2](https://doi.org/10.1175/1520-0469(2001)058<2616:CMATBI>2.0.CO;2).
- NACK ML & GREEN AES. 1974. Influence of clouds, haze, and smog on the middle ultraviolet reaching the ground. *Applied Optics* 13: 2405-2415.
- NASH ER, NEWMAN PA, ROSENFELD JE & SCHOEBERL MR. 1996. An objective determination of the polar vortex using Ertel's potential vorticity. *J Geophys Res* 101: 9471-9478. <https://doi.org/10.1029/96JD00066>.
- NDARANA T, WAUGH DW, POLVANI LM, CORREA GJP & GERBER EP. 2012. Antarctic ozone depletion and trends in tropopause rossby wave breaking. *Atmos Sci Lett* 13(3): 164-168.
- OLIVEIRA MMF. 2014. Radiação ultravioleta/ índice ultravioleta e câncer de pele no brasil: condições ambientais e vulnerabilidades sociais. *Rev Bras Climatol [s.l.]* 13: 1-14. ISSN 2237-8642.
- PALANCAR GG & TOSELLI BM. 2004. Effects of meteorology on the annual and interannual cycle of the UVB and total radiation in Cordoba City, Argentina. *Atmos Environ* 38: 1073-1082.
- PERES LV ET AL. 2017. Measurements of the total ozone column using a brewer spectrophotometer and Toms and Omi satellite instruments over the southern space observatory in Brazil. *Ann Geophys* 35: 25-37.
- PERES LV, KALL E, CRESPO NM, FONTINELE JL, ANABOR V, PINHEIRO DK, SCHUCH NJ & LEME NMP. 2011. Caracterização sinótica do evento de efeito secundário do buraco de ozônio antártico sobre o sul do Brasil do dia 14/10/2008. *Ciência e Natura*, ed. Suplementar, Universidade Federal de Santa Maria, p. 323-326.
- PERES LV, PINHEIRO DK, STEFFENEL LA, MENDES D, BAGESTON JV, BITTENCOURT GD & BENCHERIF H. 2019. Monitoramento de longo prazo e climatologia de campos estratosféricos quando da ocorrência dos eventos de influência do buraco de ozônio antártico sobre o sul do Brasil. *Rev Bras Meteorol* 34(1): 151-163.
- PERES LV, REIS NCS DOS, SANTOS LDO DOS, BITTENCOURT GD, SCHUCH AP, ANABOR V, PINHEIRO DK, SCHUCH NJ & LEME NMP. 2014. Análise Atmosférica dos Eventos de Efeito Secundário do Buraco De Ozônio Antártico Sobre O Sul do Brasil Em 2012. Parte 2: Verificação Sinótica da Troposfera Durante Os Eventos. *Ciência e Natura* 36(2): 423-433. <https://doi.org/10.5902/2179460x13151>.
- PERES LV, REIS NCS, SANTOS LO, BITTENCOURT GD, SHUCH AP, ANABOR V, PINHEIRO DK, SCHUCH NJ & LEME NP. 2016. Análise atmosférica dos eventos de efeito secundário do buraco de ozônio antártico sobre o sul do Brasil em 2012. Parte 1: identificação dos eventos e análise da dinâmica da estratosfera. *Ciência e Natura* 38: 290-299.
- PINHEIRO DK, LEME NP, PERES LV & KALL E. 2011. Influence of the antarctic ozone hole over south of Brazil in 2008 and 2009. Annual active report 2010 - National Institute of Science and Technology Antarctic Environmental Research 1: 33-37.
- PINHEIRO DK, PERES LV, CRESPO NM, SCHUCH NJ & LEME NP. 2012. Influence of the antarctic ozone hole over south of Brazil in 2010 and 2011. Annual active report 2011 - National Institute of Science and Technology Antarctic Environmental Research 1: 34-38.
- PORTAFAIX T, MOREL B, BENCHERIF H, BALDY S, GODINBEEKMANN S & HAUCHECORNE A. 2003. Fine-scale study of a thick stratospheric ozone lamina at the edge of the southern subtropical barrier. *J Geophys Res* 108(d6): 4196.
- RODRIGUEZ J. 2017. Radiação solar ultravioleta e fotodermatoses em La Paz – Bolívia (Ultraviolet solar radiation and photodermatoses in La Paz – Bolívia). *Revista Brasileira de Geografia Física [S.l.]* 10(2): 371-380.
- ROLPH G, STEIN A & STUNDER B. 2017. Real-time environmental applications and display system: ready. *Environ Modell. Softw* 95: 210-228.
- SALBY ML. 1996. Fundamentals of atmospheric physics. International geophysics series, academic press, vol. 61.
- SÁNCHEZ-PÉREZ JF, VICENTE-AGULLO D, BARBERÁ M, CASTRO-RODRÍGUEZ E & CÁNOVAS M. 2019. Relação entre o índice ultravioleta (UVI) e queimaduras solares de primeiro, segundo e terceiro grau usando a metodologia Probit. *Sci Rep* 9: 733.

SCHOEBERL MR. 1988. Dynamics weaken the polar hole. *Nature* 336: 420-421.

SCHOEBERL MR & NEWMAN PA. 1995. A multiple-level trajectory analysis of vortex filaments. *J Geophys Res* 100: 25801-25816.

SCHUCH PA, SANTOS MB, LIPINSKI VM, PERES LV, SANTOS CP, CECHIN SZ, SCHUCH N J, PINHEIRO DK & LORETO ELS. 2015. Identification of influential events concerning the antarctic ozone hole over southern brazil and the biological effects induced by uvb and uva radiation in an endemic treefrog species. *Ecotoxicol Environ Saf* 118: 190-198.

SECKMEYER G, ERB R & ALBOLD A. 1996. Transmittance of a cloud is wavelength-dependent in the UV-range. *Geophys Re Lett* 23: 2753-2755.

SEMANE N, BENCHERIF H, MOREL B, HAUCHECORNE A & DIAB RD. 2006. An unusual stratospheric ozone decrease in southern hemisphere subtropics linked to isentropic air-mass transport as observed over irene (25.5° s, 28.1° e) in mid-may 2002. *Atmos Chem Phys* 6: 1927-1936.

SERRA M & HALLER G. 2017: Efficient computation of null geodesics with applications to coherent vortex detection. *Proc Roy Soc London*, 473A. 20160807. <https://doi.org/10.1098/rspa.2016.0807>.

SHEPHERD T. 2007: Transport in the middle atmosphere. *J Meteor Soc Japan* 85: 165-191. <https://doi.org/10.2151/jmsj.85B.165>.

SOLOMON S. 1999. Stratospheric ozone depletion: a review of concepts and history. *Rev Geophys* 37(3): 275-316.

STEFFENEL LA, PINHEIRO MK, PINHEIRO DK & PERES LV. 2016 Using a pervasive computing environment to identify secondary effects of the antarctic ozone hole. *Procedia Comput Sci* 83: 1007-1012.

UTRILLAS MP, MARÍN MJ, ESTEVE AR, SALAZAR G, SUÁREZ H, GANDÍA S & MARTÍNEZ-LOZANO JA. 2018. Relationship between erythemal UV and broadband solar irradiation at high altitude in Northwestern Argentina. *Energy* 162: 136-147.

WILKS DS. 2011. *Statistical Methods in the Atmospheric Sciences*. Second Edn., Vol. 91 (Academic Press, Inc., San Diego, 2011).

WHO. *Global Solar UV Index: A practical guide*. A joint recommendation of the World Health Organization, World Meteorological Organization, United Nations Environment Programme, and the International Commission on Non-Ionizing Radiation Protection. Geneva: WHO, 2002, 1-32.

WMO - WORLD METEOROLOGICAL ORGANIZATION. 2002. GENEVA, SWITZERLAND. 2002. Scientific assessment of ozone depletion: 2002, global ozone research and monitoring project-report no. 47.

YAMAMOTO ALC, CORRÊA MDP & CCOYLLO ORS. 2018. Validation and Analysis of UV Radiation Time Series Collected In Different Peruvian Sites. *Rev Bras Meteorol, Scielo Brasil* 33(2): 298-305. <https://doi.org/10.1590/0102-7786332011>.

ZHANG G & WANG Z. 2018. North Atlantic extratropical Rossby wave breaking during the warm season: Wave life cycle and role of diabatic heating. *Mon Weather Rev* 146(3): 695-712. <https://doi.org/10.1175/MWR-D-17-0204.1>.

How to cite

SOUZA AM ET AL. 2023. Impacts of the antarctic ozone hole influence events over southern Brazil in October 2015. *An Acad Bras Cienc* 95: e20210528. DOI 10.1590/0001-3765202320210528.

*Manuscript received on April 7, 2021;
accepted for publication on August 20, 2021*

ALANNA M. DE SOUZA¹

<https://orcid.org/0000-0002-0519-2123>

LUCAS V. PERES¹

<https://orcid.org/0000-0002-5612-5991>

GABRIELA D. BITTENCOURT²

<https://orcid.org/0000-0002-4572-119X>

DAMARIS K. PINHEIRO²

<https://orcid.org/0000-0001-6939-7091>

BIBIANA C. LOPES²

<https://orcid.org/0000-0002-6732-3096>

VAGNER ANABOR²

<https://orcid.org/0000-0002-7301-2075>

NEUSA M.P. LEME³

<https://orcid.org/0000-0002-3377-9917>

MARIA PAULETE P. MARTINS⁴

<https://orcid.org/0000-0002-3103-7944>

RODRIGO DA SILVA¹

<https://orcid.org/0000-0001-9222-5861>

GABRIELA C.G. DOS REIS¹

<https://orcid.org/0000-0001-9243-214X>

MARCO ANTÔNIO G. DOS REIS¹

<https://orcid.org/0000-0001-6364-1049>

JOSÉ V. BAGESTON⁵

<https://orcid.org/0000-0003-2931-8488>

HASSAN BENCHERIF⁶

<https://orcid.org/0000-0003-1815-0667>

¹Universidade Federal do Oeste do Pará, Instituto de Engenharia e Geociências, Rua Vera Paz, s/n, Salé, 68040-255 Santarém, PA, Brazil

²Programa de Pós-Graduação em Meteorologia, Universidade Federal de Santa Maria, Av. Roraima, 1000, Camobi, 97105-900 Santa Maria, RS, Brazil

³Coordenação Espacial do Nordeste, Instituto Nacional de Pesquisas Espaciais, Rua Carlos Serrano, 2073, Lagoa Nova, 59076-740 Natal, RN, Brazil

⁴Coordenação Geral de Engenharia, Tecnologia e Ciências Espaciais, Instituto Nacional de Pesquisas Espaciais, Av. Astronautas, 1758, Jardim da Granja, 12227-010 São José dos Campos, SP, Brazil

⁵Coordenação Espacial do Sul, Instituto Nacional de Pesquisas Espaciais, Av. Roraima, 1000, Camobi, 97105-340 Santa Maria, RS, Brazil

⁶Laboratoire de l'Atmosphère et des Cyclones - LACy, Université de La Réunion, UMR 8105, 97744, Reunion Island, France

Correspondence to: **Lucas Vaz Peres**

E-mail: lucas.peres@ufopa.edu.br

Author contributions

Conceptualization, A.M.S, L.V.P. and D.K.P; methodology and graphics plot, A.M.S, L.V.P, D.K.P, G.D.B., M.A.G.R, V.A. and B.C.L.; validation and data acquisition, A.M.S, L.V.P, and B.C.L.; original writing paper, A.M.S.; review and editing, L.V.P, D.K.P, G.D.B., M.A.G.R, V.A., R.S., G.C.G.R, N.M.P.L., M.P.P.M, H.B and J.V.B.; project administration and funding acquisition, H.B., D.K.P and N.M.P.L.

