



**HAL**  
open science

# Fungal-Bacterial Combinations in Plant Health under Stress: Physiological and Biochemical Characteristics of the Filamentous Fungus *Serendipita indica* and the Actinobacterium *Zhihengliuella* sp. ISTPL4 under In Vitro Arsenic Stress

Neha Sharma, Monika Koul, Naveen Chandra Joshi, Laurent Dufossé, Arti Mishra

## ► To cite this version:

Neha Sharma, Monika Koul, Naveen Chandra Joshi, Laurent Dufossé, Arti Mishra. Fungal-Bacterial Combinations in Plant Health under Stress: Physiological and Biochemical Characteristics of the Filamentous Fungus *Serendipita indica* and the Actinobacterium *Zhihengliuella* sp. ISTPL4 under In Vitro Arsenic Stress. *Microorganisms*, 2024, 12, 10.3390/microorganisms12020405 . hal-04493586

**HAL Id: hal-04493586**

<https://hal.univ-reunion.fr/hal-04493586v1>

Submitted on 7 Mar 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



## Article

# Fungal-Bacterial Combinations in Plant Health under Stress: Physiological and Biochemical Characteristics of the Filamentous Fungus *Serendipita indica* and the Actinobacterium *Zhihengliuella* sp. ISTPL4 under In Vitro Arsenic Stress

Neha Sharma <sup>1</sup>, Monika Koul <sup>2</sup> , Naveen Chandra Joshi <sup>1</sup>, Laurent Dufossé <sup>3,\*</sup> and Arti Mishra <sup>2,4,\*</sup>

<sup>1</sup> Amity Institute of Microbial Technology, Amity University, Noida 201313, India; nehasharma96500@gmail.com (N.S.); ncjoshi@amity.edu (N.C.J.)

<sup>2</sup> Department of Botany, Hansraj College, University of Delhi, Delhi 110007, India; drmkoul@gmail.com

<sup>3</sup> Chemistry and Biotechnology of Natural Products, CHEMBIOPRO, Université de La Réunion, ESIROI Agroalimentaire, 15 Avenue René Cassin, CS 92003, CEDEX 9, F-97744 Saint-Denis, France

<sup>4</sup> Umeå Plant Science Center, Department of Plant Physiology, Umeå University, 90187 Umeå, Sweden

\* Correspondence: laurent.dufosse@univ-reunion.fr (L.D.); artimishrahrc@gmail.com (A.M.)

**Abstract:** Fungal-bacterial combinations have a significant role in increasing and improving plant health under various stress conditions. Metabolites secreted by fungi and bacteria play an important role in this process. Our study emphasizes the significance of secondary metabolites secreted by the fungus *Serendipita indica* alone and by an actinobacterium *Zhihengliuella* sp. ISTPL4 under normal growth conditions and arsenic (As) stress condition. Here, we evaluated the arsenic tolerance ability of *S. indica* alone and in combination with *Z. sp. ISTPL4* under in vitro conditions. The growth of *S. indica* and *Z. sp. ISTPL4* was measured in varying concentrations of arsenic and the effect of arsenic on spore size and morphology of *S. indica* was determined using confocal microscopy and scanning electron microscopy. The metabolomics study indicated that *S. indica* alone in normal growth conditions and under As stress released pentadecanoic acid, glycerol tricaprylate, L-proline and cyclo(L-prolyl-L-valine). Similarly, d-Ribose, 2-deoxy-bis(thioheptyl)-dithioacetal were secreted by a combination of *S. indica* and *Z. sp. ISTPL4*. Confocal studies revealed that spore size of *S. indica* decreased by 18% at 1.9 mM and by 15% when in combination with *Z. sp. ISTPL4* at a 2.4 mM concentration of As. Arsenic above this concentration resulted in spore degeneration and hyphae fragmentation. Scanning electron microscopy (SEM) results indicated an increased spore size of *S. indica* in the presence of *Z. sp. ISTPL4* ( $18 \pm 0.75 \mu\text{m}$ ) compared to *S. indica* alone ( $14 \pm 0.24 \mu\text{m}$ ) under normal growth conditions. Our study concluded that the suggested combination of microbial consortium can be used to increase sustainable agriculture by combating biotic as well as abiotic stress. This is because the metabolites released by the microbial combination display antifungal and antibacterial properties. The metabolites, besides evading stress, also confer other survival strategies. Therefore, the choice of consortia and combination partners is important and can help in developing strategies for coping with As stress.

**Keywords:** *Serendipita indica*; arsenic; heavy metal stress; secondary metabolites; *Oryza sativa*



**Citation:** Sharma, N.; Koul, M.; Joshi, N.C.; Dufossé, L.; Mishra, A. Fungal-Bacterial Combinations in Plant Health under Stress: Physiological and Biochemical Characteristics of the Filamentous Fungus *Serendipita indica* and the Actinobacterium *Zhihengliuella* sp. ISTPL4 under In Vitro Arsenic Stress. *Microorganisms* **2024**, *12*, 405. <https://doi.org/10.3390/microorganisms12020405>

Academic Editor: Gary A. Strobel

Received: 12 January 2024

Revised: 1 February 2024

Accepted: 6 February 2024

Published: 17 February 2024



**Copyright:** © 2024 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

Various anthropogenic activities such as mining, modern agricultural practices, and industrialization have had long-term harmful impacts on the environment [1]. These factors are responsible for increasing the concentration of heavy metals in soil and water, thus leading to several environmental issues. Arsenic (As) is a highly toxic heavy metal that poses risk to millions of people worldwide [2]. Arsenic contamination in drinking water and groundwater used for irrigation is a global problem that not only affects agricultural productivity but also the ecosystem, as As uptake occurs in plant roots and is further

translocated to different plant tissues, including the edible parts, and therefore enters the ecosystem through the food chain [3]. Arsenite and arsenate are the two forms of arsenic, arsenite being more toxic. Roots are the prime site of arsenic exposure; thus root proliferation and extension are affected the most. The translocation of arsenic occurs from root to shoot, and the uptake in the aerial parts affects plant growth and reproduction by inhibiting cell division [4]. Therefore, it is essential to remove these heavy metals from the contaminated soil used for growing crop plants.

Microbes including fungal and bacterial species have the potential to cope with heavy metal stress. These microbes also aid in the growth and development of plants by increasing nutrient absorption and assimilation [5]. Microbes release various molecules including siderophores, extracellular polysaccharides, ammonia, and secondary metabolites that help plants to cope with biotic and abiotic stress [6]. Secondary metabolites are chemical substances that protect plants under stress conditions. They are not directly involved in plant growth and development but indirectly boost plant growth by acting as effective defense agents against various phytopathogens. They also help in scavenging free radicals generated during oxidative stress in plants [7].

Metabolites secreted by bacteria and fungi in the rhizosphere are not directly connected with plant growth and reproduction but they participate in rhizosphere ecological interactions [8]. Lipopeptides, phytohormones, their precursors, antimicrobial peptides, siderophores, metallothioneins, and volatile organic compounds are some examples of secondary metabolites released by the microbes [9]. Secondary metabolites, such as acyl-homoserine lactone (acyl-HSL), that act as autoinducer signaling molecules and help in cell-to-cell communication and crosstalk are also secreted by some microbes. Siderophores are some other secondary metabolites released by microbes that help in iron uptake from surroundings. Siderophores-releasing microbes help in iron uptake by the plants also [9]. Other metabolites including metallothioneins are also secreted by various microbes and plants. These are the chelating molecules that help in heavy metal detoxification [10]. These metallothioneins have tiny cysteine-rich proteins which bind with heavy metals thereby helping in metal storage and detoxification. These metallothioneins are mostly released in the presence of heavy metals such as  $\text{Cd}^{2+}$ ,  $\text{Hg}^{2+}$ ,  $\text{Pb}^{2+}$  and Ar.

In this study, we studied various secondary metabolites released by *S. indica* alone and with *Zhiehngliuella* sp. ISTPL4 under normal growth conditions and under arsenic stress. *S. indica* is an endophytic fungus that boosts plant growth by colonizing the roots of various plants and confers resistance against abiotic and biotic stress conditions [11–13]. *Zhiehngliuella* sp. ISTPL4 is an actinobacterium, isolated from Pangong Lake, Ladakh, Jammu and Kashmir, India. The interaction between *Zhiehngliuella* sp. ISTPL4 and *S. indica* has been previously reported in a systemic study [14]. The role of this microbial combination has also been deciphered in plant growth promotion. In the present study, we checked the significance of secondary metabolites secreted by a combination of *S. indica* and *Z. sp. ISTPL4* and found out which metabolites are released in response to combined microbial treatments.

## 2. Materials and Methods

### 2.1. Microbe Culture and Conditions

The fungal disc of *S. indica* was inoculated into Hill and Kaefer agar plates and incubated for 14 days at 28–30 °C (with an agitation rate of 120 rpm for broth). Similarly, *S. indica* growth was also assessed in combination with *Z. sp. ISTPL4*. Bacterial culture was streaked around the periphery after five days of fungal inoculation (5 dafi) [15].

### 2.2. Effect of Arsenic on Fungal Growth

The arsenic tolerance capability of *S. indica* was checked individually as well as in combination under in vitro conditions. The concentration of arsenic used was up to 2.4 mM [16]. Dry cell weight of *S. indica* alone and in combination was estimated (control and treated cultures) to check the effect of arsenic on the growth of *S. indica* [14].

### 2.3. Spore Morphology

To study spore morphology SEM studies were carried out for *S. indica* alone and in combination with *Z. sp. ISTPL4*. A section of fungal discs was fixed in 2.5% glutaraldehyde (control and interaction plates) followed by incubation at room temperature (RT) for 1 h. After this, samples were centrifuged at 5000 rpm for 5 min followed by washing with 0.1 M phosphate buffer (pH 7) followed by centrifugation. The resulting pellet was dissolved in 0.1% filter sterilized silver nitrate ( $\text{AgNO}_3$ ) solution and then incubated for an hour at RT. Samples were gradually dehydrated in a series of ethanol from 30–90% for 15 min. The final step was repeated three times in 100% ethanol for 5 min each. Resulting samples were dehydrated, air dried and placed in aluminum double adhesive carbon conductive tape gold-coated specimens in a Quorum Q150ES coater and then observed under a scanning electron microscope (model Zeiss EVO 18; Raipur, India) [14].

Confocal microscopy was also carried out to check spore size and spore morphology of *S. indica* alone and in combination with *Z. sp. ISTPL4* under normal growth conditions and under As stress. Spore isolation was done by adding 1 mL of a 0.02% autoclaved Tween 20 on freshly grown cultures of *S. indica* alone and with *Z. sp. ISTPL4* (control and As-exposed plates) followed by gentle scraping and centrifugation at 5000 rpm for 10–15 min. The pellet was mixed with autoclaved distilled water, a spore count was conducted, and their concentration was maintained at  $4.8 \times 10^5$  spores/mL. The resulting sample was analyzed under a Nikon confocal microscope (Model: Nikon A1) at 60 $\times$  magnification by using NIS Elements software version 4.6 (Nikon, Tokyo, Japan) (Amity Institute of Microbial Technology, Amity University, Noida, India) [17].

### 2.4. Biotransformation of Arsenic

The arsenic oxido-reduction potential of *S. indica* and *Z. sp. ISTPL4* was determined. For this, nutrient agar was supplemented with 500 ppm of As (V) and the fungal disc was inoculated followed by an incubation of 15 days at  $28 \pm 2$  °C. Spot inoculation of *Z. sp. ISTPL4* was conducted, and the bacterial culture was kept for incubation at 37 °C for 72 h. After fungal and bacterial growth, 100  $\mu\text{L}$   $\text{AgNO}_3$  (0.1 M) solution was flooded over the grown fungal and bacterial plates and a change in color was observed [18].

### 2.5. Gas Chromatography and Mass Spectroscopy Analysis (GC-MS)

GC-MS analysis was performed for the secondary metabolic profiling in *S. indica* alone and in combination with *Z. sp. ISTPL4* under normal growth conditions and in the presence of arsenic stress. The fungal disc was inoculated in H and K broth and then kept for incubation for 2 weeks. A similar protocol was followed for the combined growth of *S. indica* and *Z. sp. ISTPL4*. The bacterial culture was inoculated in H and K broth after 5 days of fungal inoculation (5 dafi) followed by incubation (2 weeks). A 100 mL sample was taken and then centrifuged at 9000 rpm for 10 min. The supernatant was collected and mixed with an equal volume of ethyl acetate. The resulting solution was shaken for 3 h. The sample was then concentrated on a rotatory evaporator. The final volume was mixed with methanol. This concentrated sample was used for GC-MS analysis for secondary metabolite detection.

GC-MS analysis was carried out in a Perkin-Elmer GC Clarus 500 system, which included an Elite-5MS (5% diphenyl/95% dimethyl poly siloxane) fused capillary column (30  $\times$  0.25  $\mu\text{m}$  ID  $\times$  0.25  $\mu\text{m}$  df) and an AOC-20i auto-sampler and Gas Chromatograph interfaced to a Mass Spectrometer (GC-MS). An electron ionization device with an ionization energy of 70 eV was used in electron impact mode for GC-MS detection. A split ratio of 10:1 was used with an injection volume of 2  $\mu\text{L}$  and helium gas (99.999%) was used as the carrier gas at a steady flow rate of 1 mL/min. The temperature of the ion source was 200 °C, the injector was kept at 250 °C, and the oven was set at 110 °C (isothermal for two minutes) which was further increased by 10 °C/min to 200 °C, and 5 °C/min to 280 °C, and stopped with a 9-min isothermal at 280 °C. Mass spectra with fragments from 45 to 450 Da and a scan interval of 0.5 s were obtained at 70 eV. The GC-MS was run for 36 min,

with a solvent delay of 0 to 2 min. By comparing the average peak area of each component to the total areas, the relative percentage amount was determined. The Turbo-Mass ver-5.2 software was utilized to handle mass spectra and chromatograms. Metabolite identification, retention time (R.T.) and mass to charge ratio ( $m/z$ ) was analyzed as listed in the NIST library [19–21].

### 2.6. Clustered Heatmap of Metabolites

A heatmap of common metabolites secreted by the individual culture of *S. indica* as well as in combination with *Z. sp. ISTPL4* under normal growth conditions and in the presence of As stress was prepared by versatile matrix visualization and analysis software (<https://software.broadinstitute.org/morpheus/> (accessed on 1 June 2022)).

### 2.7. Statistical Method

The experiments were repeated five times. Statistical analysis was conducted utilizing two-way ANOVA and student's *t* test ( $p < 0.05$ ) using online OPSTAT software (version 6.8). All the variables were subjected to statistical analysis.

## 3. Results

### 3.1. Microbial Conditions

The growth of *S. indica* was checked alone and with *Z. sp. ISTPL4* by measuring the hyphal radius and spore size was measured using scanning electron microscopy (Figure 1a–d). *S. indica* growth was assessed under varying concentrations of As and it was observed that *S. indica* was able to survive at 1.9 mM concentration (Figure 2). Growth of *S. indica* under normal conditions was ( $3.1 \pm 0.025$  cm) and it decreased up to ( $1.6 \pm 0.01$  cm) at 1.9 mM As concentration. *S. indica* growth was assessed in combination with *Z. sp. ISTPL4*. *S. indica* when grown in combination with *Z. sp. ISTPL4* was able to grow at 2.4 mM concentration. The hyphal radius of *S. indica* was ( $2.9 \pm 0.01$  cm) when grown with *Z. sp. ISTPL4* under normal growth conditions and it reduced ( $1.4 \pm 0.03$  cm) in the presence of As (Figure 2a–d).

The dry cell weight of *S. indica* alone as well as in combination with *Z. sp. ISTPL4* was measured in the control as well as in the presence of As. The dry cell weight of *S. indica* was 69% higher than dry cell weight of *S. indica* under As stress. Similarly, dry cell weight of combination of *S. indica* and *Z. sp. ISTPL4* was 51% more than the dry cell weight of the combination of *S. indica* and *Z. sp. ISTPL4* under arsenic stress (Figure 2e).


### 3.2. Spore Morphology

Increased spore size of *S. indica* was observed in the presence of *Z. sp. ISTPL4* ( $18 \pm 0.75$   $\mu\text{m}$ ) as compared to its individual culture ( $14 \pm 0.24$   $\mu\text{m}$ ) under normal growth conditions and confocal microscopy under normal growth conditions in scanning electron microscopy (Figure 1c–d). The spore size of *S. indica* decreased by 18% at 1.9 mM concentration and by 15% when used in combination with *Z. sp. ISTPL4* at 2.4 mM concentration of As as observed under a confocal microscope. Increase in concentration beyond this resulted in spore degeneration and hyphae fragmentation (Figure 3a–d). The spore count was also decreased at higher concentration of arsenic. It was  $4.59 \times 10^5$  spores/mL in *S. indica* under normal growth conditions and it reduced to  $3.68 \times 10^3$  spores/mL at 1.9 mM concentration of arsenic. Similarly, the spore count of *S. indica* was slightly higher  $4.65 \times 10^5$  spores/mL, when it was grown in presence of *Z. sp. ISTPL4* and at 2.4 mM concentration, it decreased to  $4 \times 10^3$  spores/mL.

### 3.3. Biotransformation of Arsenic

The biotransformation potential of *S. indica* and *Z. sp. ISTPL4* was checked on the basis of the color change of fungal and bacterial cultures. Color change to brown in fungal cultures indicated transformation ability of As (V) (arsenate) to As (III) (arsenite). The ability of the fungal strain to bring conversion was discovered after 72 h of incubating

fungus and bacterial plates. Contact of silver nitrate with As (V) (already present in media) produced a brownish precipitate, while the association with As (III) produced a pale yellow precipitate. The results of the arsenic transformation ability of *S. indica* indicated brown color formation which denotes that this fungus has certain genes which are involved in arsenic transformation. *Z. sp. ISTPL4* revealed no change in color which indicated no conversion of arsenic to other forms (Figure 4a–d).




**Figure 1.** Growth of (a) *Serendipita indica* (Fungus) alone (b) in the presence of *Zihhengliuella sp. ISTPL4* (Bacteria) on a Hill and Kaefer agar plate (c) morphology of *S. indica* spores visualized using scanning electron microscopy (SEM) at Magnification: 2910 $\times$ ; voltage: 10 Kv; scale: 10  $\mu$ m. (d) Spore morphology of *S. indica* in presence of *Z. sp. ISTPL4* at Magnification: 2000 $\times$ ; voltage: 10 Kv; scale: 10  $\mu$ m.


### 3.4. Production of Secondary Metabolites

Secondary metabolite production in *S. indica* alone as well as in combination with *Z. sp. ISTPL4* was checked under normal growth conditions and under As stress (Figure 5a,b). GC-MS studies revealed the presence of 69 metabolites secreted by *S. indica* under normal conditions and 47 metabolites under As stress. Among these metabolites, 23 metabolites are common including 2,4-Di-tert-butylphenol having a retention time of 12.910 and molecular weight 206, 1,2-Benzenedicarboxylic acid with a retention time of 17.125 and molecular weight of 278, respectively (Table 1). 2-Ethylbutyric acid, eicosyl ester, 2-hydroxy-1(hydroxymethyl)ethyl ester, Octocrylene 2-propenoic acid and Squalene e 2,6,10,14,18,22-Tetracosahaxaene are some of the metabolites secreted by culture of *S. indica* alone and under arsenic stress. A total of 67 metabolites were secreted by the combination of *S. indica* and *Z. sp. ISTPL4* under normal conditions, and 37 metabolites were secreted under arsenic stress (Figure 6a,b). Similarly, 16 metabolites were common including Cyclo(L-prolyl-L-


valine), hexahydro-3-(2-methylpropyl), Eicosane, 2,4-Di-tert-butylphenol, L-Proline and Heneicosane (Table 1). All these metabolites are involved in antibacterial, antifungal, and antioxidant activities while some of them have important roles in plant growth promotion under various environmental conditions. These metabolites are also represented in the form of venn diagram (Figure 7a,b). Figure 8 shows the clustered heatmap of the secondary metabolites. These metabolites include volatile organic compounds (VOCs), hydrocarbons, amino acids, fatty acids and vitamins. Hierarchical clustering analysis (HCA) shows the comparison of common metabolites based on the area percentage detected. A heatmap ranging from red (high) to blue (low) denoted the concentration of each metabolite detected. The numerical data of metabolites were normalized to allow for clustering and color scaling based on the concentration of compounds. In Figure 8a, compounds that appeared as red were considered highly abundant, and Figure 8b showed that an equal ratio of red and blue colored compounds appeared.




**Figure 2.** Growth of *S. indica* alone (a) under normal growth condition ( $2.9 \pm 0.01$  cm) (b) in As stress ( $1.6 \pm 0.01$  cm; 1.9 mM) (c) in presence of *Z. sp.* ISTPL4 under normal growth condition ( $2.5 \pm 0.05$  cm), (d) in presence of As stress ( $1.4 \pm 0.03$  cm; 2.4 mM) and (e) shows the difference in dry weight of *S. indica* alone, in the presence of arsenic stress, in combination with *Z. sp.* ISTPL4 and under arsenic stress after microbial inoculation. According to the student's *t*-test, asterisks showed significant differences: \**t*:  $p \leq 0.05$ ; \*\**t*:  $p \leq 0.01$ ).



**Figure 3.** Cont.




**Figure 3.** Confocal microscopy analysis shows the spore morphology and spore size of (a) *S. indica* alone (b) under arsenic stress (1.9 mM) (c) in combination with *Z. sp.* ISTPL4 and, (d) in combination with *Z. sp.* ISTPL4 under arsenic stress (2.4 mM).




**Figure 4.** Screening of arsenate transformation ability of *S. indica* and *Z. sp.* ISTPL4: Change in color from white to yellow indicates the reduction of arsenate to arsenite (a) *S. indica* (control): Arsenate plate without silver nitrate (b) *S. indica* (Test): Arsenate plate with silver nitrate (c) *Z. sp.* ISTPL4 (control): Arsenate plate without silver nitrate (d) *Z. sp.* ISTPL4 (test): Arsenate plate with silver nitrate.





(a)



(b)


**Figure 5.** GC-MS chromatogram of secondary metabolites secreted by (a) *S. indica* under normal growth conditions and, (b) in the presence of arsenic.

**Table 1.** List of common metabolites secreted by *S. indica* alone and in combination with *Z. sp. ISTPL4* under normal growth conditions and in presence of As stress.


S. No.	Metabolites ( <i>S. indica</i> / <i>S. indica</i> ; As)	Metabolites ( <i>S. indica</i> + <i>Z. sp. ISTPL4</i> / <i>S. indica</i> + <i>Z. sp. ISTPL4</i> ; As)
1	2,4-Di-tert-butyl-phenol) phosphate	2,4-Di-tert-butyl-phenol) phosphate
2	Cyclo(L-prolyl-L-valine)	Cyclo(L-prolyl-L-valine)
3	1,2-Benzenedicarboxylic acid, bis (2-methyl propyl) ester	1,2-Benzenedicarboxylic acid, bis (2-methyl propyl) ester
4	7,9-Di-tert-butyl-1-oxaspiro (4,5) deca-a-6,9-diene-2,8-dione	7,9-Di-tert-butyl-1-oxaspiro (4,5) deca-a-6,9-diene-2,8-dione
5	Hexadecenoic acid, methyl ester	Hexadecenoic acid, methyl ester

Table 1. Cont.


S. No.	Metabolites ( <i>S. indica</i> / <i>S. indica</i> ; As)	Metabolites ( <i>S. indica</i> + <i>Z. sp. ISTPL4</i> / <i>S. indica</i> + <i>Z. sp. ISTPL4</i> ; As)
6	Pyrrole[1,2-a]pyrazine-1,4-dione, hexahydro-3-(2-methyl propyl)	octadecanoic acid, 3-oxo-, ethyl ester
7	1,2-Benzenedicarboxylic acid, butyl octyl ester	Olean-18-ene
8	Methyl stearate	L-Proline
9	2-Ethylbutyric acid, eicosyl ester	Glycerol 1-palmitate
10	2-(4-Hydroxy-4-methyl-tetrahydro-pyran-3-ylamino)-3-(1H-indol-2-yl)-propionic acid	Hexadecanoic acid, 2-hydroxy-1-(hydroxymethyl)ethyl ester
11	Lycopene	Octadecanoic acid, 2,3-dihydroxypropyl ester
12	Hexadecanoic acid, 2-hydroxy-1-(hydroxymethyl) ethyl ester	Methyl stearate
13	Bis(2-ethylhexyl) phthalate	Pyrrolo[1,2-a]pyrazine-1,4-dione, hexahydro-3-(2-methylpropyl)-
14	Octocrylene 2-Propenoic acid,	Tetradecane
15	Octadecanoic acid, 2,3-dihydroxypropyl ester	Eicosane
16	Squalene e 2,6,10,14,18,22-Tetracosahexaene,	Heneicosane
17	Phenol, 2,4-bis (1,1-dimethyl ethyl)-, phosphite	
18	Tris (2,4-di-tert-butyl phenyl) phosphate	
19	Octahydro-2H-pyrido(1,2-a)pyrimidin-2-one	
20	Olean-18-ene	
21	l-Leucine	
22	L-Proline	
23	Ergotaman-3',6',18-trione	



**Figure 6.** GC-MS chromatogram of secondary metabolites secreted by (a) a combination of *S. indica* and *Z. sp. ISTPL4* under normal growth conditions and, (b) in the presence of arsenic.



**Figure 7.** (a) Venn diagram representing different and common metabolites released by *S. indica* under normal growth conditions and in the presence of arsenic with common metabolites mentioned in box. (b) Venn diagram representing different and common metabolites released by combination of *S. indica* and *Z. sp. ISTPL4* under normal growth conditions and in the presence of arsenic with common metabolites mentioned in box.



**Figure 8.** Gas chromatography-mass spectrometry (GC-MS) spectrum comparison (a) clustered heatmap of common metabolites secreted by *S. indica* alone under normal growth conditions and in presence of As stress (b) heatmap showing common metabolites secreted by combination of *S. indica* and *Z. sp. ISTPL4* under normal growth conditions and in presence of As stress; heatmap ranging from red to blue shows high to low abundance of metabolites detected.

#### 4. Discussion

Microorganisms release various secondary metabolites to cope with abiotic stress resulting in better plant growth performance, increased photosynthesis, and the production of antioxidants [21]. Besides bacteria, fungal species also confer abiotic stress tolerance in plants by helping them to adapt to a variety of conditions, such as heat, salinity, cold, drought, and toxic metals [22]. *Fusarium culmorum*, *Azotobacter*, *Pseudomonas*, *Curvularia protuberata*, *Piriformospora indica*, *Neotyphodium lolii*, and *Trichoderma* have all been documented to confer abiotic stress tolerance [23]. Among these, *S. indica* is a plant-growth-promoting root endophytic fungus. It plays an important role in defending plants from adverse biotic and abiotic factors [24,25]. The role of various secondary metabolites secreted by *S. indica*

alone and in combination with *Z. sp. ISTPL4* was studied in the present study. *S. indica* growth was also checked in the presence of arsenic individually and in combination with *Z. sp. ISTPL4*. It was reported that *S. indica* was able to grow up to 1.9 mM and up to 2.4 mM in combination with *Z. sp. ISTPL4* in the culture conditions. Mohd et al. (2017) studied *P. indica* growth under arsenite (III) and arsenate (V) stress and observed a reduction in the growth of fungus at higher As concentrations [16]. The spore morphology and size of *S. indica* alone and in combination was also assessed under control conditions and in As stress. It was observed that the hyphae and spores were intact at 1.9 mM As concentration and hyphae started fragmenting and spores degenerated beyond 1.9 mM concentration of As in *S. indica* alone. Similarly, spore size and morphology of *S. indica* in combination with *Z. sp. ISTPL4* was observed in up to 2.4 mM concentration of arsenic. The morphology of *S. indica* spores was also checked in the presence of Cd by Dabral et al. 2019. It was observed that spore germination inhibited beyond a 0.1 mM concentration of Cd [26].


An Increase of 69% in dry cell weight was observed in *S. indica* when compared with dry cell weight of *S. indica* under arsenic stress. Similarly, the dry cell weight of the combined culture of *S. indica* and *Z. sp. ISTPL4* was 51% more than a combined culture of *S. indica* and *Z. sp. ISTPL4* under arsenic stress. The spore morphology of *S. indica* was also checked under normal conditions and in arsenic stress by confocal microscopy. It was observed that the spore degenerated beyond a 1.9 mM concentration of arsenic and beyond 2.4 mM in the microbial combination of *S. indica* and *Z. sp. ISTPL4*. The results of the spore count of *S. indica* under arsenic stress was measured which indicated a reduction in spore count to  $3.68 \times 10^3$  spores/mL at 1.9 mM concentration compared to the control ( $4.59 \times 10^5$  spores/mL). The spore count of *S. indica* in the presence of *Z. sp. ISTPL4* was also measured, which indicated an increased spore count ( $4.65 \times 10^5$  spores/mL) under normal growth conditions, and it was reduced to  $4 \times 10^3$  spores/mL. The spore germination and spore size of *S. indica* alone as well as in the presence of *Z. sp. ISTPL4* was also determined under normal growth conditions which indicated an increased spore size and germination of *S. indica* in the presence of *Z. sp. ISTPL4* than its respective control (*S. indica*) [14].

The arsenic biotransformation ability of *S. indica* and *Z. sp. ISTPL4* was confirmed. The results of arsenic biotransformation in *S. indica* indicated the formation of a brown-colored precipitate (Figure 4), which indicated the conversion of arsenate (v) to arsenite (III). Mohd et al. (2017) also reported the bioaccumulation and biotransformation ability of *S. indica* under arsenic stress [16]. In the case of *Z. sp. ISTPL4*, there was no change in color observed but the findings of whole genome sequencing indicated the presence of arsenic-resistant genes in *Z. sp. ISTPL4* [18,27].

In this study, we reported the significance of various metabolites secreted by *S. indica* alone and in combination with *Z. sp. ISTPL4* under normal and in arsenic stress. These metabolites regulate plant growth and development under stress conditions by acting as antioxidants which scavenge free radicals generated in plants during oxidative stress [28,29]. A total of 69 metabolites were produced in *S. indica* under normal conditions and 47 metabolites were produced in As stress (Supplementary File S3). Among all the metabolites analyzed, 23 metabolites were common (Table 1). These metabolites include 1,2-Benzenedicarboxylic acid, 2-ethylbutyric acid, eicosyl ester, Octocrylene 2-propenoic acid, squalene 2,6,10,14,18,22-Tetracosahaxaene, phenol and, 2,4-bis(1,1-dimethyl-ethyl)-phosphite. The chromatogram of each metabolite has been shown in Supplementary Files S1 and S4. Similarly, 67 metabolites were secreted by a combination of *S. indica* and *Z. sp. ISTPL4* under normal conditions and 37 metabolites were secreted by a combination of *S. indica* and *Z. sp. ISTPL4* under arsenic stress (Supplementary File S4). Among them, 16 metabolites were common including Cyclo(L-prolyl-L-valine), hexa hydro-3-(2-methylpropyl), Eicosane, 2,4-Di-tert-butylphenol, methyl ester, Methyl stearate and L-Proline (Table 1). These metabolites have different functions such as antifungal activities, antimicrobial activities, and antioxidant activities [28–30]. Some of these metabolite such as glycine, Quinoline-4-carboxamide 2-phenyl-N-n-octyl are antimicrobial and reduces

stress in plants [31,32]. Other metabolites including Dodecane, 4,6-dimethyl, Tetradecane, I-Hexadecanol, Nonadecane and, Heneicosane were also identified in the individual culture of *S. indica*. Similarly, 5-Azacytosine, *N,N,O*-trimethyl, 5-Nitroso-2,4,6-triaminopyrimidine, d-Ribose, 2-deoxy-bis(thioheptyl)-dithioacetal, 5-Nitroso-2,4,6-triaminopyrimidine were secreted by combination of *S. indica* and *Z. sp. ISTPL4*. These metabolites have antimicrobial, antioxidant and nematocidal activities. I-hexadecanol is a major volatile compound produced by molds and acts as an antifeedant in different species of aphids [23,25,32,33]. Heneicosane has antimicrobial activities against *Streptococcus pneumonia* and *Aspergillus fumigatus* and Cyclo(L-propyl-L-valine) is a class of diketopiperazines (DKPs). These metabolites are mainly involved in interactions between mixed microbial communities [33,34]. The secretion of this metabolite by microbes can be a possible reason for the positive interaction of *S. indica* and *Z. sp. ISTPL4*. Other metabolites including l-(+)-Ascorbic acid 2,6-dihexadecanoate, Docosanoic acid, ethyl ester, Isopropyl palmitate, 2-Methyltetracosane, n-Hexane, 13-Docosenamide (Z) have been reported. These metabolites play a crucial role in antiallergic, antibacterial, antioxidant, termiticide and antiviral activity, antimicrobial and free radical scavenging activity [35,36]. Tetratriacontyl heptafluorobutyrate, myristic acid, glycidyl ester, tetradecanoic acid, 2-Propenoic acid, 3-(4-methoxyphenyl) are the metabolites released by *S. indica* in the presence of As. These metabolites have antifungal and antimicrobial activity [6,34,37]. Metabolites including 2,3-dihydroxypropyl ester Stearin, Eicosane, 3-Indol-1-yl-propionic acid, methyl ester and, Glycerol tricaprilate, have antimicrobial, antifungal, probiotics and antioxidant activities [38–40]. L-Proline, glycine, squalene, heptadecane, 7,9 di-tert butyl-1-oxaspiro(4,5) deca-a-6,9-diene-2,8-dione, Dodecane,4,6-dimethyl, Tridecanoic acid, 12-methyl-, methyl ester, 2-methyl tetracosane and quercetin play an essential role in plant growth and development under abiotic stress [41–45]. These metabolites protect plants under stress by acting as antioxidants, promoting signaling cascade in plants for enhancing plant growth and scavenging free radicals generated during oxidative stress in plants. Quercetin is a polyphenolic compound with antioxidant properties secreted by microbial combinations.

The study suggests that an active efflux system operates in the removal of arsenic by *S. indica* and *Z. sp. ISTPL4*. *S. indica* and *Z. sp. ISTPL4* uptakes arsenate through phosphate transporters (Ars C) and converts it into arsenite by arsenate reductase. The converted form of arsenite (III) is then shunted out. The presence of the ArsC gene is also reported in *Z. sp. ISTPL4* which also confers arsenic tolerance ability of this actinobacterium. This combination of microbes can be used to mitigate the toxic effects of arsenic in the contaminated environments and increase sustainable agriculture in the soils that are contaminated (Figure 9).



**Figure 9.** Schematic representation of arsenic uptake and efflux by *S. indica* and *Z. sp. ISTPL4*.

## 5. Conclusions

Our study concluded the significance of various metabolites secreted by *S. indica* alone and in combination with *Z. sp. ISTPL4* under normal environmental conditions and in the presence of arsenic stress. The secretion of Cyclo(L-propyl-L-valine) by both *S. indica* and *Z. sp. ISTPL4* indicated the possible reason for their interaction. Also, the arsenic tolerance ability of *S. indica* was checked individually and in combination with *Z. sp. ISTPL4* in this study. Results of the biotransformation experiment also indicated the arsenic tolerance ability of *S. indica* and *Z. sp. ISTPL4*. However, these microbes can be used to mitigate arsenic stress. The presence of various metabolites also indicated the capability of *S. indica* and *Z. sp. ISTPL4* in plant growth and stress alleviation under various biotic and abiotic factors. Our data are largely confirmed by other scientific literature, which has shown that these microbial strains can efficiently remove arsenic from contaminated environments.

**Supplementary Materials:** The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/microorganisms12020405/s1>, Supplementary File S1: Mass fragmentation pattern and structural elucidations of common metabolites secreted by *Serendipita indica* under normal growth conditions and in presence of arsenic stress; Supplementary File S2: Mass fragmentation patterns and structural elucidations of common secondary metabolites secreted by *Serendipita indica* and *Zhihengliuella sp. ISTPL4* under normal growth conditions and in the presence of arsenic stress; Supplementary File S3: Secondary metabolites production by *Serendipita indica* and *Zhihengliuella sp. ISTPL4* under normal growth conditions and in presence of As stress; Supplementary file S4: Secondary metabolites production by *Serendipita indica* and *Zhihengliuella sp. ISTPL4* under normal growth conditions and in presence of As stress. Refs. [40–154] can be found in Supplementary Materials.

**Author Contributions:** Conceptualization, A.M. and L.D.; methodology, N.S.; software, M.K.; validation, A.M.; investigation, N.C.J.; resources, N.C.J.; data curation, A.M. and L.D.; writing—original draft preparation, N.S.; writing—review and editing, N.S.; supervision, N.C.J.; project administration, N.C.J.; funding acquisition, N.C.J. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research received no external funding.

**Data Availability Statement:** Data are contained within the article.

**Acknowledgments:** All the authors are thankful to the Amity Institute of Microbial Technology, Amity University, Noida for providing all the facilities and DST-FIST for the Confocal facility. Laurent Dufossé is deeply thankful to the Conseil Régional de La Réunion and the Conseil Régional de Bretagne for their continuous support of microbial biotechnology activities.

**Conflicts of Interest:** The authors declare no conflicts of interest.

## References

1. Zaman, N.R.; Chowdhury, U.F.; Reza, R.N.; Chowdhury, F.T.; Sarker, M.; Hossain, M.M.; Akbor, M.A.; Amin, A.; Islam, M.R.; Khan, H. Plant Growth Promoting Endophyte Burkholderia Contaminans NZ Antagonizes Phytopathogen Macrophomina Phaseolina through Melanin Synthesis and Pyrrolnitrin Inhibition. *PLoS ONE* **2021**, *16*, e0257863. [CrossRef] [PubMed]
2. Genchi, G.; Lauria, G.; Catalano, A.; Carocci, A.; Sinicropi, M.S. Arsenic: A Review on a Great Health Issue Worldwide. *Appl. Sci.* **2022**, *12*, 6184. [CrossRef]
3. Bianucci, E.; Peralta, J.M.; Furlan, A.; Hernández, L.E. *Arsenic in Drinking Water and Food*; Springer: Berlin/Heidelberg, Germany, 2020. [CrossRef]
4. Chandrakar, V.; Pandey, N.; Keshavkant, S. Plant Responses to Arsenic Toxicity: Morphology and Physiology. In *Mechanisms of Arsenic Toxicity and Tolerance in Plants*; Springer: Singapore, 2018.
5. Pérez-Montaña, F.; Alías-Villegas, C.; Bellogín, R.A.; Del Cerro, P.; Espuny, M.R.; Jiménez-Guerrero, I.; López-Baena, F.J.; Ollero, F.J.; Cubo, T. Plant Growth Promotion in Cereal and Leguminous Agricultural Important Plants: From Microorganism Capacities to Crop Production. *Microbiol. Res.* **2014**, *169*, 325–336. [CrossRef] [PubMed]
6. Mitra, D.; Mondal, R.; Khoshru, B.; Senapati, A.; Radha, T.K.; Mahakur, B.; Uniyal, N.; Myo, E.M.; Boutaj, H.; Sierra, B.E.G.; et al. Actinobacteria-Enhanced Plant Growth, Nutrient Acquisition, and Crop Protection: Advances in Soil, Plant, and Microbial Multifactorial Interactions. *Pedosphere* **2022**, *32*, 149–170. [CrossRef]

7. Koza, N.A.; Adayo, A.A.; Babalola, O.O.; Kappo, A.P. Microorganisms in Plant Growth and Development: Roles in Abiotic Stress Tolerance and Secondary Metabolites Secretion. *Microorganisms* **2022**, *10*, 1528. [[CrossRef](#)]
8. Pang, Z.; Chen, J.; Wang, T.; Gao, C.; Li, Z.; Guo, L.; Xu, J.; Cheng, Y. Linking Plant Secondary Metabolites and Plant Microbiomes: A Review. *Front. Plant Sci.* **2021**, *12*, 621276. [[CrossRef](#)]
9. Barra Caracciolo, A.; Terenzi, V. Rhizosphere Microbial Communities and Heavy Metals. *Microorganisms* **2021**, *9*, 1462. [[CrossRef](#)]
10. Raychaudhuri, S.S.; Pramanick, P.; Talukder, P.; Basak, A. Polyamines, Metallothioneins, and Phytochelatins—Natural Defense of Plants to Mitigate Heavy Metals. *Stud. Nat. Prod. Chem.* **2021**, *69*, 227–261.
11. Varma, A.; Sowjanya Sree, K.; Arora, M.; Bajaj, R.; Prasad, R.; Kharkwal, A.C. Functions of Novel Symbiotic Fungus—*Piriformospora indica*. *Proc. Indian Natl. Sci. Acad.* **2014**, *80*, 429–441. [[CrossRef](#)]
12. Jangir, P.; Shekhawat, P.K.; Bishnoi, A.; Ram, H.; Soni, P. Role of *Serendipita indica* in Enhancing Drought Tolerance in Crops. *Physiol. Mol. Plant Pathol.* **2021**, *116*, 101691. [[CrossRef](#)]
13. Yang, G.; Ahmad, F.; Zhou, Q.; Guo, M.; Liang, S.; Gaal, H.A.; Mo, J. Investigation of Physicochemical Indices and Microbial Communities in Termite Fungus-Combs. *Front. Microbiol.* **2021**, *11*, 581219. [[CrossRef](#)] [[PubMed](#)]
14. Sharma, N.; Dabral, S.; Tyagi, J.; Yadav, G.; Aggarwal, H.; Joshi, N.C.; Varma, A.; Koul, M.; Choudhary, D.K.; Mishra, A. Interaction Studies of *Serendipita indica* and *Zhihengliuella* sp. ISTPL4 and Their Synergistic Role in Growth Promotion in Rice. *Front. Plant Sci.* **2023**, *14*, 1155715. [[CrossRef](#)] [[PubMed](#)]
15. Hill, T.W.; Kafer, E. Improved Protocols for *Aspergillus minimal* Medium: Trace Element and Minimal Medium Salt Stock Solutions. *Fungal Genet. Rep.* **2001**, *48*, 20–21. [[CrossRef](#)]
16. Mohd, S.; Shukla, J.; Kushwaha, A.S.; Mandrah, K.; Shankar, J.; Arjaria, N.; Saxena, P.N.; Narayan, R.; Roy, S.K.; Kumar, M. Endophytic Fungi *Piriformospora indica* Mediated Protection of Host from Arsenic Toxicity. *Front. Microbiol.* **2017**, *8*, 754. [[CrossRef](#)]
17. Dabral, S.; Saxena, S.C.; Choudhary, D.K.; Bandyopadhyay, P.; Sahoo, R.K.; Tuteja, N.; Nath, M. Synergistic Inoculation of *Azotobacter Vinelandii* and *Serendipita indica* Augmented Rice Growth. *Symbiosis* **2020**, *81*, 139–148. [[CrossRef](#)]
18. Bhati, R.; Sreedharan, S.M.; Rizvi, A.; Khan, M.S.; Singh, R. An Insight into Efflux-Mediated Arsenic Resistance and Biotransformation Potential of *Enterobacter cloacae* RSC3 from Arsenic Polluted Area. *Indian J. Microbiol.* **2022**, *62*, 456–467. [[CrossRef](#)]
19. Gengmao, Z.; Quanmei, S.; Yu, H.; Shihui, L.; Changhai, W. The Physiological and Biochemical Responses of a Medicinal Plant (*Salvia miltiorrhiza* L.) to Stress Caused by Various Concentrations of NaCl. *PLoS ONE* **2014**, *9*, e89624. [[CrossRef](#)] [[PubMed](#)]
20. Tripathi, A.; Singh, Y.; Verma, D.K.; Ranjan, M.R.; Srivastava, S.K. Bioremediation of Hazardous Azo Dye Methyl Red by a Newly Isolated *Bacillus megaterium* ITBHU01: Process Improvement through ANN-GA Based Synergistic Approach. *Indian J. Biochem. Biophys.* **2016**, *53*, 112–125.
21. Anantha, P.S.; Deventhiran, M.; Saravanan, P.; Anand, D.; Rajarajan, S. A Comparative GC-MS Analysis of Bacterial Secondary Metabolites of *Pseudomonas* Species. *Pharma Innov.* **2016**, *5*, 84–89.
22. Poveda, J.; Baptista, P.; Sacristán, S.; Velasco, P. Editorial: Beneficial Effects of Fungal Endophytes in Major Agricultural Crops. *Front. Plant Sci.* **2022**, *13*, 1061112. [[CrossRef](#)]
23. Okon, O.; Matrood, A.; Okon, J. Na<sup>+</sup> Exclusion and Na<sup>+</sup>/K<sup>+</sup> Ratio Adjustment by Mycorrhiza Enhances Macro/Micro Nutrients Uptake in Two Members of Cucurbitaceae Family under Salt Stress. *Int. J. Bot. Stud.* **2020**, *5*, 108–114.
24. Fa, A.N. Endophytic Fungi for Sustainable Agriculture. *Microb. Biosyst.* **2019**, *4*, 31–44. [[CrossRef](#)]
25. Rabiey, M.; Ullah, I.; Shaw, L.J.; Shaw, M.W. Potential Ecological Effects of *Piriformospora indica*, a Possible Biocontrol Agent, in UK Agricultural Systems. *Biol. Control* **2017**, *104*, 1–9. [[CrossRef](#)]
26. Dabral, S.; Yashaswee; Varma, A.; Choudhary, D.K.; Bahuguna, R.N.; Nath, M. Biopriming with *Piriformospora indica* Ameliorates Cadmium Stress in Rice by Lowering Oxidative Stress and Cell Death in Root Cells. *Ecotoxicol. Environ. Saf.* **2019**, *186*, 109741. [[CrossRef](#)] [[PubMed](#)]
27. Mishra, A.; Jha, G.; Thakur, I.S. Draft Genome Sequence of *Zhihengliuella* sp. Strain ISTPL4, a Psychrotolerant and Halotolerant Bacterium Isolated from Pangong Lake, India. *Genome Announc.* **2018**, *6*, e01533-17. [[CrossRef](#)] [[PubMed](#)]
28. Bernhard, A.E.; Sheffer, R.; Giblin, A.E.; Marton, J.M.; Roberts, B.J. Population Dynamics and Community Composition of Ammonia Oxidizers in Salt Marshes after the Deepwater Horizon Oil Spill. *Front. Microbiol.* **2016**, *7*, 854. [[CrossRef](#)]
29. Vanitha, V.; Vijayakumar, S.; Nilavukkarasi, M.; Punitha, V.; Vidhya, E.; Praseetha, P. Heneicosane—A Novel Microbicidal Bioactive Alkane Identified from *Plumbago Zeylanica* L. *Ind. Crops Prod.* **2020**, *154*, 112748. [[CrossRef](#)]
30. Wani, S.H.; Kumar, V.; Shriram, V.; Sah, S.K. Phytohormones and Their Metabolic Engineering for Abiotic Stress Tolerance in Crop Plants. *Crop J.* **2016**, *4*, 162–176. [[CrossRef](#)]
31. Singh, R.; Kumar, M.; Mittal, A.; Mehta, P.K. Microbial Metabolites in Nutrition, Healthcare and Agriculture. *3 Biotech* **2017**, *7*, 15. [[CrossRef](#)]
32. Gadhi, A.A.A.; El-Sherbiny, M.M.O.; Al-Sofyani, A.M.A.; Ba-Akdah, M.A.; Satheesh, S. Antibiofilm Activities of Extracts of the Macroalga *Halimeda* sp. from the Red Sea. *J. Mar. Sci. Technol.* **2018**, *26*, 838–846. [[CrossRef](#)]
33. Khattab, A.I.; Babiker, E.H.; Saeed, H.A. Streptomyces: Isolation, Optimization of Culture Conditions and Extraction of Secondary Metabolites. *Int. Curr. Pharm. J.* **2016**, *5*, 27–32. [[CrossRef](#)]
34. Shaaban, M.T.; Ghaly, M.F.; Fahmi, S.M. Antibacterial Activities of Hexadecanoic Acid Methyl Ester and Green-Synthesized Silver Nanoparticles against Multidrug-Resistant Bacteria. *J. Basic Microbiol.* **2021**, *61*, 557–568. [[CrossRef](#)]
35. Ferdosi, M.F.H.; Javaid, A.; Khan, I.H. Phytochemical Profile of N-Hexane Flower Extract of *Cassia fistula* L. *Bangladesh J. Bot.* **2022**, *51*, 393–399. [[CrossRef](#)]

36. Chenniappan, J.; Sankaranarayanan, A.; Arjunan, S. Evaluation of Antimicrobial Activity of *Cissus quadrangularis* L. Stem Extracts against Avian Pathogens and Determination of Its Bioactive Constituents Using GC-MS. *J. Sci. Res.* **2020**, *64*, 90–96. [[CrossRef](#)]
37. Chandrasekaran, M.; Senthilkumar, A.; Venkatesalu, V. Antibacterial and Antifungal Efficacy of Fatty Acid Methyl Esters from the Leaves of *Sesuvium portulacastrum* L. *Eur. Rev. Med. Pharmacol. Sci.* **2011**, *15*, 775–780.
38. Naim, M.J.; Alam, O.; Alam, J.; Bano, F.; Alam, P.; Shrivastava, N. Recent Review on Indole: A privileged scaffold structure. *Int. J. Pharma Sci. Res.* **2016**, *7*, 51–62.
39. Chen, S.; Liu, J.; Gong, H.; Yang, D. Identification and Antibacterial Activity of Secondary Metabolites from *Taxus* Endophytic Fungus. *Chin. J. Biotechnol.* **2009**, *25*, 368–374. [[PubMed](#)]
40. Kumari, N.; Menghani, E.; Mithal, R. GCMS Analysis of Compounds Extracted from Actinomycetes AIA6 Isolates and Study of Its Antimicrobial Efficacy. *Indian J. Chem. Technol.* **2019**, *26*, 362–370.
41. Ramya, B.; Malarvili, T.; Velavan, S. GC-MS Analysis of Bioactive Compounds in Bryonopsis Laciniosa Fruit Extract. *Int. J. Pharm. Sci. Res.* **2015**, *6*, 3375–3379.
42. Mooney, E.C.; Holden, S.E.; Xia, X.J.; Li, Y.; Jiang, M.; Banson, C.N.; Zhu, B.; Sahingur, S.E. Quercetin Preserves Oral Cavity Health by Mitigating Inflammation and Microbial Dysbiosis. *Front. Immunol.* **2021**, *12*, 774273. [[CrossRef](#)] [[PubMed](#)]
43. Babar, A.G. Antifungal Activity and Investigation of Bioactive Compounds of Marine Intertidal Bivalve *Gafrarium Divaricatum* from West Coast of India. *Int. J. Pure Appl. Biosci.* **2016**, *4*, 211–217. [[CrossRef](#)]
44. Nakkeeran, S.; Priyanka, R.; Rajamanickam, S.; Sivakumar, U. *Bacillus amyloliquefaciens* Alters the Diversity of Volatile and Non-Volatile Metabolites and Induces the Expression of Defence Genes for the Management of Botrytis Leaf Blight of Liliium under Protected Conditions. *J. Plant Pathol.* **2020**, *102*, 1179–1189. [[CrossRef](#)]
45. Younis, A.A.; Saleh, H.A. Phytochemical Screening and Assessment of Antioxidant and Antimicrobial Potentialities of Two Egyptian Medicinal Plants. *Egypt. J. Pure Appl. Sci.* **2021**, *59*, 49–57. [[CrossRef](#)]
46. Ahmad Mustafa, F.N. Green Synthesis of Isopropyl Palmitate Using Immobilized *Candida Antarctica* Lipase: Process Optimization Using Response Surface Methodology. *Clean. Eng. Technol.* **2022**, *8*, 100516. [[CrossRef](#)]
47. Kumari, N.; Menghani, E.; Mithal, R. Bioactive Compounds Characterization and Antibacterial Potentials of Actinomycetes Isolated from Rhizospheric Soil. *J. Sci. Ind. Res.* **2019**, *78*, 793–798.
48. Parthasarathy, R. Ethnic Quotas as Term-Limits: Caste and Distributive Politics in South India. *Comp. Political Stud.* **2017**, *50*, 1735–1767. [[CrossRef](#)]
49. Arya, A.; Kumar, S.; Kain, D.; Ahanger, A.M.; Suryavanshi, A.; Vandana. Unfolding the Chemical Composition, Antioxidant and Antibacterial Activities of *Drymaria cordata* (Linn.) Willd. against Chloramphenicol-Resistant *Bacillus subtilis* and  $\beta$ -Lactams-Resistant *Pseudomonas aeruginosa*. *Vegetos* **2023**, 1–10. [[CrossRef](#)]
50. Babatunde, D.E.; Otusemade, G.O.; Efeovbokhan, V.E.; Ojewumi, M.E.; Bolade, O.P.; Owioye, T.F. Chemical Composition of Steam and Solvent Crude Oil Extracts from *Azadirachta indica* Leaves. *Chem. Data Collect.* **2019**, *20*, 100208. [[CrossRef](#)]
51. Kalaba, M.H.; Sultan, M.H.; Elbahnasawy, M.A.; El-Didamony, S.E.; Bakary, N.M.E.; Sharaf, M.H. First Report on Isolation of *Mucor Bainieri* from Honeybees, *Apis mellifera*: Characterization and Biological Activities. *Biotechnol. Rep.* **2022**, *36*, e00770. [[CrossRef](#)] [[PubMed](#)]
52. Selvamani, K.; Tadigiri, S.; Das, D. Isolation and Characterization of Chemical Constituents from *B. amyloliquefaciens* and Their Nematicidal Activity. *J. Entomol. Zool. Stud.* **2020**, *8*, 2062–2066.
53. Zhu, F.; Zhu, C.; Doyle, E.; Gao, J. Fate of Di (2-Ethylhexyl) Phthalate in Different Soils and Associated Bacterial Community Changes. *Sci. Total Environ.* **2018**, *637*, 460–469. [[CrossRef](#)]
54. Saboon, Bibi, Y.; Ayoubi, S.A.; Afzal, T.; Gilani, S.; Malik, K.; Qayyum, A.; Hussain, M.; Kumar, S. Antioxidant-Activity-Guided Purification and Separation of Octocrylene from *Saussurea heteromalla*. *Separations* **2023**, *10*, 107. [[CrossRef](#)]
55. Stevenson, J.; Luu, W.; Kristiana, I.; Brown, A.J. Squalene Mono-Oxygenase, a Key Enzyme in Cholesterol Synthesis, Is Stabilized by Unsaturated Fatty Acids. *Biochem. J.* **2014**, *461*, 435–442. [[CrossRef](#)] [[PubMed](#)]
56. Eramma, N.; Patil, S.J. Exploration of the Biomolecules in roots of *Flacourtia indica* (Burm. F) Merr. Methanol Extract by Chromatography Approach. *Lett. Appl. NanoBioSci.* **2023**, *12*, 166–177.
57. Tyagi, S.; Singh, R.K.; Tiwari, S.P. Anti-Enterococcal and Anti-Oxidative Potential of a Thermophilic Cyanobacterium, *Leptolyngbya* sp. HNBGU 003. *Saudi J. Biol. Sci.* **2021**, *28*, 4022–4028. [[CrossRef](#)]
58. Saleh, A.E.; Ul-Hassan, Z.; Zeidan, R.; Al-Shamary, N.; Al-Yafei, T.; Alnaimi, H.; Higazy, N.S.; Migheli, Q.; Jaoua, S. Biocontrol Activity of *Bacillus megaterium* BM344-1 against Toxicogenic Fungi. *ACS Omega* **2021**, *6*, 10984–10990. [[CrossRef](#)]
59. Guo, C.; Chen, Y.; Wu, D.; Du, Y.; Wang, M.; Liu, C.; Chu, J.; Yao, X. Transcriptome Analysis Reveals an Essential Role of Exogenous Brassinolide on the Alkaloid Biosynthesis Pathway in *Pinellia Ternata*. *Int. J. Mol. Sci.* **2022**, *23*, 10898. [[CrossRef](#)] [[PubMed](#)]
60. Chakraborty, B.; Suresh, R.; Almansour, A.I.; Gunasekaran, P.; Nayaka, S. Saudi Journal of Biological Sciences Bioprospection and Secondary Metabolites Profiling of Marine *Streptomyces levis* Strain KS46. *Saudi J. Biol. Sci.* **2022**, *29*, 667–679. [[CrossRef](#)]
61. Uyan, A.; Turan, C.; Erdogan-Eliuz, E.A.; Sangun, M.K. Antimicrobial Properties of Bioactive Compounds Isolated from Epidermal Mucus in Two Ray Species (*Dasyatis marmorata* and *Gymnura altavela*). *Trop. J. Pharm. Res.* **2020**, *19*, 2115–2121. [[CrossRef](#)]
62. Siddharthan, N.; Balagurunathan, R.; Venkatesan, S.; Hemalatha, N. Bio-Efficacy of *Geobacillus thermodenitrificans* PS41 against Larvicidal, Fungicidal, and Plant Growth-Promoting Activities. *Environ. Sci. Pollut. Res.* **2022**, *30*, 42596–42607. [[CrossRef](#)]



63. Kanbak, S.A.; Koksall, M.; Sahin, F.; Erol, D.D. Synthesis of New 4(1H)-Pyridinone Derivatives and Their Antibacterial Activity. *Rev. Chim.* **2009**, *60*, 888–892.
64. Łukowska-Chojnacka, E.; Kowalkowska, A.; Gizińska, M.; Koronkiewicz, M.; Staniszevska, M. Synthesis of Tetrazole Derivatives Bearing Pyrrolidine Scaffold and Evaluation of Their Antifungal Activity against *Candida albicans*. *Eur. J. Med. Chem.* **2019**, *164*, 106–120. [[CrossRef](#)]
65. Dutta, J.; Thakur, D. Evaluation of Antagonistic and Plant Growth Promoting Potential of *Streptomyces* sp. TT3 Isolated from Tea (*Camellia Sinensis*) Rhizosphere Soil. *Curr. Microbiol.* **2020**, *77*, 1829–1838. [[CrossRef](#)]
66. Sakthiselvan, P.; Madhumathi, R.; Meenambiga, S.S. Moronic Acid: An Antiviral for Herpes Simplex Virus. In *A Century of Valuable Plant Bioactives*; Academic Press: Berlin/Heidelberg, Germany, 2021; pp. 143–158. [[CrossRef](#)]
67. Díaz-Pérez, A.L.; Díaz-Pérez, C.; Campos-García, J. Bacterial L-Leucine Catabolism as a Source of Secondary Metabolites. *Rev. Environ. Sci. Bio/Technol.* **2016**, *15*, 1–29. [[CrossRef](#)]
68. Jung, H.; Hilger, D.; Raba1, M. The Na<sup>+</sup>/L-Proline Transporter PutP. *Front. Biosci.* **2012**, *17*, 745–759. [[CrossRef](#)] [[PubMed](#)]
69. Shwaish, T.; Al-Imarah, F.J.M.; Jasim, F.A. Wound Healing Capacity, Antibacterial Activity, and GC-MS Analysis of *Bienertia sinuspersici* Leaves Extract. *J. Phys. Conf. Ser.* **2019**, *1294*, 062056. [[CrossRef](#)]
70. Rhetso, T.; Shubharani, R.; Roopa, M.S.; Sivaram, V. Chemical Constituents, Antioxidant, and Antimicrobial Activity of *Allium chinense* G. Don. *Future J. Pharm. Sci.* **2020**, *6*, 102. [[CrossRef](#)]
71. Touria Ould Bellahcen, M.C. Chemical Composition and Antibacterial Activity of the Essential Oil of *Spirulina platensis* from Morocco. *J. Essent. Oil Bear. Plants* **2019**, *22*, 1265–1276. [[CrossRef](#)]
72. Singh, P.; Singh, J.; Ray, S.; Vaishnav, A.; Jha, P.; Singh, R.K.; Singh, H. Microbial Volatiles (MVOs) Induce Tomato Plant Growth and Disease Resistance Against Wilt Pathogen *Fusarium oxysporum* f.sp. *lycopersici*. *J. Plant Growth Regul.* **2023**, 1–14. [[CrossRef](#)]
73. Ladkau, N.; Assmann, M.; Schrewe, M.; Julsing, M.K.; Schmid, A.; Bühler, B. Efficient Production of the Nylon 12 Monomer ω-Aminododecanoic Acid Methyl Ester from Renewable Dodecanoic Acid Methyl Ester with Engineered *Escherichia coli*. *Metab. Eng.* **2016**, *36*, 1–9. [[CrossRef](#)]
74. Ashraf, M.F.M.R.; Foolad, M.R. Roles of Glycine Betaine and Proline in Improving Plant Abiotic Stress Resistance. *Environ. Exp. Bot.* **2007**, *59*(2), 206–216. [[CrossRef](#)]
75. He, K.; Zhang, T.; Zhang, S.; Sun, Z.; Zhang, Y.; Yuan, Y.; Jia, X. Tunable Functionalization of Saturated C–C and C–H Bonds of *N,N'*-Diarylpiperazines Enabled by *tert*-Butyl Nitrite (TBN) and NaNO<sub>2</sub> Systems. *Org. Lett.* **2019**, *21*, 5030–5034. [[CrossRef](#)]
76. Vorobjeva, L.I.; Khodjaev, E.Y.; Vorobjeva, N.V. Propionic Acid Bacteria as Probiotics. *Microb. Ecol. Health Dis.* **2008**, *20*, 109–112. [[CrossRef](#)]
77. Rathod, S.V.; Shinde, K.W.; Kharkar, P.S.; Shah, C.P.; Aruna, K.; Raut, D.A. Synthesis, Molecular Docking and Biological Evaluation of New Quinoline Analogues as Potent Anti-Breast Cancer and Antibacterial Agents. *Indian J. Chem. Sect. B Org. Med. Chem.* **2021**, *60B*, 1215–1222. [[CrossRef](#)]
78. Monjil, M.S.; Takemoto, D.; Kawakita, K. Defense Induced by a Bis-Aryl Methanone Compound Leads to Resistance in Potato against *Phytophthora infestans*. *J. Gen. Plant Pathol.* **2014**, *80*, 38–49. [[CrossRef](#)]
79. Singh, J.; Sharma, S.; Kaur, A.; Vikal, Y.; Cheema, A.K.; Bains, B.K.; Kaur, N.; Gill, G.K.; Malhotra, P.K.; Kumar, A.; et al. Marker-Assisted Pyramiding of Lycopene-ε-Cyclase, β-Carotene Hydroxylase1 and Opaque2 Genes for Development of Biofortified Maize Hybrids. *Sci. Rep.* **2021**, *11*, 12642. [[CrossRef](#)] [[PubMed](#)]
80. Daood, U.; Matinlinna, J.P.; Pichika, M.R.; Mak, K.K.; Nagendrababu, V.; Fawzy, A.S. A Quaternary Ammonium Silane Antimicrobial Triggers Bacterial Membrane and Biofilm Destruction. *Sci. Rep.* **2020**, *10*, 10970. [[CrossRef](#)] [[PubMed](#)]
81. Das, P.; Devi, N.; Gaur, N.; Goswami, S.; Dutta, D.; Dubey, R.; Puzari, A. Acrylonitrile Adducts: Design, Synthesis and Biological Evaluation as Antimicrobial, Haemolytic and Thrombolytic Agent. *Sci. Rep.* **2023**, *13*, 6209. [[CrossRef](#)] [[PubMed](#)]
82. Yadav, B.; Jogawat, A.; Rahman, M.S.; Narayan, O.P. Secondary Metabolites in the Drought Stress Tolerance of Crop Plants: A Review. *Gene Rep.* **2021**, *23*, 101040. [[CrossRef](#)]
83. Harikrishnan, S.; Parivallal, M.; Alsalhi, M.S.; Sudarshan, S.; Jayaraman, N.; Devanesan, S.; Rajasekar, A.; Jayalakshmi, S. Characterization of Active Lead Molecules from *Lissocarinus orbicularis* with Potential Antimicrobial Resistance Inhibition Properties. *J. Infect. Public Health* **2021**, *14*, 1903–1910. [[CrossRef](#)] [[PubMed](#)]
84. Kumaresan, S.; Senthilkumar, V.; Stephen, A.; Balakumar, B.S. GC-MS analysis and pass-assisted prediction of biological activity spectra of extract of *Phomopsis* sp. isolated from *Andrographis paniculata*. *World J. Pharm. Res.* **2015**, *4*, 1035–1053.
85. Trivedi, D.K.; Sinclair, E.; Xu, Y.; Sarkar, D.; Walton-Doyle, C.; Liscio, C.; Banks, P.; Milne, J.; Silverdale, M.; Kunath, T.; et al. Discovery of Volatile Biomarkers of Parkinson’s Disease from Sebum. *ACS Cent. Sci.* **2019**, *5*, 599–606. [[CrossRef](#)]
86. Varsha, K.K.; Devendra, L.; Shilpa, G.; Priya, S.; Pandey, A.; Nampootheri, K.M. 2,4-Di-*tert*-butyl Phenol as the Antifungal, Antioxidant Bioactive Purified from a Newly Isolated *Lactococcus* sp. *Int. J. Food Microbiol.* **2015**, *211*, 44–50. [[CrossRef](#)]
87. Abdullah, R. Insecticidal Activity of Secondary Metabolites of Locally Isolated Fungal Strains against Some Cotton Insect Pests. *J. Plant Prot. Pathol.* **2019**, *10*, 647–653. [[CrossRef](#)]
88. Rani, A.; Saini, K.C.; Bast, F.; Mehariya, S.; Bhatia, S.K.; Lavecchia, R.; Zuurro, A. Microorganisms: A Potential Source of Bioactive Molecules for Antioxidant Applications. *Molecules* **2021**, *26*, 1142. [[CrossRef](#)]
89. Sharma, S.; Chandra, D.; Sharma, A.K. Rhizosphere Plant–Microbe Interactions Under Abiotic Stress. In *Rhizosphere Biology: Interactions between Microbes and Plants*; Springer: Singapore, 2021; pp. 195–216. [[CrossRef](#)]

90. Javed, S.; Javaid, A.; Al-Taie, A.H.; Zahid, M. Identification of Antimicrobial Compounds from *n*-Hexane Stem Extract of *Kochia indica* by GC-MS Analysis. *Mycopath* **2018**, *16*, 51–55.
91. Ahmad, R.; Ahmad, T.; Hussain, B. *Proceedings of International Conference on Recent Advances in Space Technologies (IEEE Cat. No.03EX743)*; IEEE: New York, NY, USA, 2011; ISBN 9788195663415.
92. Rachmawaty; Mu’Nisa, A.; Hasri; Pagarra, H.; Hartati; Qureshi, M.Z. Active Compounds Extraction of Cocoa Pod Husk (*Thebroma cacao* L.) and Potential as Fungicides. *J. Phys. Conf. Ser.* **2018**, *1028*, 012013. [[CrossRef](#)]
93. Duraisamy, M.; Raja, S. Analysis of Bioactive Compounds by Gas Chromatography-Mass Spectrum and Anti-Bacterial Activity of *Zonaria crenata*. *Aegaeum J.* **2020**, *8*, 829–844. [[CrossRef](#)]
94. Younus, I.; Khan, S.J.; Maqbool, S.; Begum, Z. Anti-Fungal Therapy via Incorporation of Nanostructures: A Systematic Review for New Dimensions. *Phys. Scr.* **2022**, *97*, 012001. [[CrossRef](#)]
95. Ser, H.L.; Palanisamy, U.D.; Yin, W.F.; Abd Malek, S.N.; Chan, K.G.; Goh, B.H.; Lee, L.H. Presence of Antioxidative Agent, Pyrrolo[1,2-*a*]pyrazine-1,4-dione, Hexahydro- in Newly Isolated *Streptomyces mangrovisoli* sp. nov. *Front. Microbiol.* **2015**, *6*, 854. [[CrossRef](#)] [[PubMed](#)]
96. Komansilan, A.; Abadi, A.L.; Yanuwadi, B.; Kaligis, D.A. Isolation and Identification of Biolarvicide from Soursop (*Annona muricata* Linn) Seeds to Mosquito (*Aedes aegypti*) Larvae. *Int. J. Eng. Technol.* **2012**, *12*, 28–32.
97. Momodu, I.B.; Okungbowa, E.S.; Agoreyo, B.O.; Maliki, M.M. Gas Chromatography—Mass Spectrometry Identification of Bioactive Compounds in Methanol and Aqueous Seed Extracts of *Azanza garckeana* Fruits. *Niger. J. Biotechnol.* **2022**, *38*, 25–38. [[CrossRef](#)]
98. Ansari, M.W.; Gill, S.S.; Tuteja, N. *Piriformospora indica* a Powerful Tool for Crop Improvement. *Proc. Indian Natl. Sci. Acad.* **2014**, *80*, 317–324. [[CrossRef](#)]
99. Sahu, P.K.; Jayalakshmi, K.; Tilgam, J.; Gupta, A.; Nagaraju, Y.; Kumar, A.; Rajput, V.D.; Vikram, M.; Rajawat, S. ROS Generated from Biotic Stress: Effects on Plants and Alleviation by Endophytic Microbes. *Front. Plant Sci.* **2022**, *13*, 1042936. [[CrossRef](#)] [[PubMed](#)]
100. Rani, S.V.G.; Murugaiah, K. GC-MS Determination of Bioactive Compounds in *Azima tetraacantha* Leaves. *World J. Pharm. Res.* **2015**, *4*, 2225–2231.
101. Suharti, W.S.; Tini, E.W.; Istiqomah, D. Antimicrobial Activity of *Kaempferia galanga* against Plant Pathogen on Rice. *Biodiversitas* **2023**, *24*, 1320–1326.
102. Saravanan, R.; Pemiah, B.; Narayanan, M.; Ramalingam, S. In Vitro Cytotoxic and Gas Chromatography-Mass Spectrometry Studies on *Orthosiphon stamineus* Benth. (Leaf) against MCF-7 Cell Lines. *Asian J. Pharm. Clin. Res.* **2017**, *10*, 129–135. [[CrossRef](#)]
103. Okoro, E.E.; Osoniyi, O.R.; Jabeen, A.; Shams, S.; Choudhary, M.I.; Onajobi, F.D. Anti-Proliferative and Immunomodulatory Activities of Fractions from Methanol Root Extract of *Abrus precatorius* L. *Clin. Phytosci.* **2019**, *5*, 45. [[CrossRef](#)]
104. Adnan, M.; Nazim Uddin Chy, M.; Mostafa Kamal, A.T.M.; Azad, M.O.K.; Paul, A.; Uddin, S.B.; Barlow, J.W.; Faruque, M.O.; Park, C.H.; Cho, D.H. Investigation of the Biological Activities and Characterization of Bioactive Constituents of *Ophiorrhiza rugosa* var. *prostrata* (D.Don) & Mondal Leaves through in Vivo, in Vitro, and in Silico Approaches. *Molecules* **2019**, *24*, 1367. [[CrossRef](#)]
105. Ponnamma, S.U.; Manjunath, K. GC-MS Analysis of Phytocomponents in the Methanolic Extract of *Justicia wynaadensis* (NEES) T. Anders. *Int. J. Pharma Bio Sci.* **2012**, *3*, 570–576.
106. Chrzęszcz, M.; Miazga-Karska, M.; Klimek, K.; Dybowski, M.P.; Typek, R.; Tchórzewska, D.; Dos Santos Szewczyk, K. The Anti-Acne Potential and Chemical Composition of *Knautia drymeia* Heuff. and *Knautia macedonica* Griseb Extracts. *Int. J. Mol. Sci.* **2023**, *24*, 9188. [[CrossRef](#)]
107. Kandasamy, S.; Sahu, S.K.; Kandasamy, K. In Silico Studies on Fungal Metabolite against Skin Cancer Protein (4,5-Diarylisoxazole HSP90 Chaperone). *Int. Sch. Res. Not.* **2012**, *2012*, 626214. [[CrossRef](#)]
108. Oviya, R.; Thiruvudainambi, S.; Ramamoorthy, V.; Vendan, R.T.; Vellaikumar, S. Gas Chromatography Mass Spectrometry (GCMS) Analysis of the Antagonistic Potential of *Trichoderma hamatum* against *Fusarium oxysporum* f. sp. *cepae* Causing Basal Rot Disease of Onion. *J. Biol. Control* **2022**, *36*, 17–30. [[CrossRef](#)]
109. Sen, S.; Dehingia, M.; Talukdar, N.C.; Khan, M. Chemometric Analysis Reveals Links in the Formation of Fragrant Bio-Molecules during Agarwood (*Aquilaria malaccensis*) and Fungal Interactions. *Sci. Rep.* **2017**, *7*, 44406. [[CrossRef](#)] [[PubMed](#)]
110. Ain, Q.U.; Asad, S.; Jamal, A.; Arif, M.; Mahmood-Ul-hassan, M. Phytochemical Analysis and Antifungal Activity of Gymnosperm against *Fusarium* Wilt of Banana. *Appl. Ecol. Environ. Res.* **2021**, *19*, 2477–2493. [[CrossRef](#)]
111. O’Leary, J.; Hiscox, J.; Eastwood, D.C.; Savoury, M.; Langley, A.; McDowell, S.W.; Rogers, H.J.; Boddy, L.; Müller, C.T. The Whiff of Decay: Linking Volatile Production and Extracellular Enzymes to Outcomes of Fungal Interactions at Different Temperatures. *Fungal Ecol.* **2019**, *39*, 336–348. [[CrossRef](#)]
112. Abu Zeid, I.M.; Al-Thobaiti, S.A.; EL Hag, G.A.; Alghamdi, S.A.; Umar, A.; Abdalla Ahmed Hamdi, O. Phytochemical and GC-MS Analysis of Bioactive Compounds from *Balanites aegyptiaca*. *Acta Sci. Pharm. Sci.* **2019**, *3*, 129–134. [[CrossRef](#)]
113. Jiménez-Teja, D.; Daoubi, M.; Collado, I.G.; Hernández-Galán, R. Lipase-Catalyzed Resolution of 5-Acetoxy-1,2-dihydroxy-1,2,3,4-tetrahydronaphthalene. Application to the Synthesis of (+)-(3*R*,4*S*)-*cis*-4-Hydroxy-6-deoxyscytalone, a Metabolite Isolated from *Colletotrichum acutatum*. *Tetrahedron* **2009**, *65*, 3392–3396. [[CrossRef](#)]
114. Zhang, Y.; Berman, A.; Shani, E. Plant Hormone Transport and Localization: Signaling Molecules on the Move. *Annu. Rev. Plant Biol.* **2023**, *74*, 453–479. [[CrossRef](#)]

115. Yang, H.; Luo, P. Changes in Photosynthesis Could Provide Important Insight into the Interaction between Wheat and Fungal Pathogens. *Int. J. Mol. Sci.* **2021**, *22*, 8865. [[CrossRef](#)]
116. Sebastian, J.; Hegde, K.; Kumar, P.; Rouissi, T.; Brar, S.K. Bioproduction of Fumaric Acid: An Insight into Microbial Strain Improvement Strategies. *Crit. Rev. Biotechnol.* **2019**, *39*, 817–834. [[CrossRef](#)]
117. Vijay, K.; Hassan, S.; Govarthanam, M.; Kavitha, T. Biosurfactant Lipopeptides and Polyketide Biosynthetic Gene in *Rhizobacterium Achromobacter kerstersii* to Induce Systemic Resistance in Tomatoes. *Rhizosphere* **2022**, *23*, 100558. [[CrossRef](#)]
118. Palani, V.; Shanmugasundaram, M.; Maluventhen, V.; Chinnaraj, S.; Liu, W.; Balasubramanian, B.; Arumugam, M. Phytoconstituents and Their Potential Antimicrobial, Antioxidant and Mosquito Larvicidal Activities of *Goniothalamus wightii* Hook. F. & Thomson. *Arab. J. Sci. Eng.* **2020**, *45*, 4541–4555. [[CrossRef](#)]
119. Ratheesh, M.; Sunil, S.; Sheethal, S.; Jose, S.P.; Sandya, S.; Ghosh, O.S.N.; Rajan, S.; Jagmag, T.; Tilwani, J. Anti-Inflammatory and Anti-COVID-19 Effect of a Novel Polyherbal Formulation (Imusil) via Modulating Oxidative Stress, Inflammatory Mediators and Cytokine Storm. *Inflammopharmacology* **2022**, *30*, 173–184. [[CrossRef](#)]
120. Aziz, S.D.A.; Jafarah, N.F.; Yusof, Z.N.B. Phytol-Containing Seaweed Extracts as Control for *Ganoderma boninense*. *J. Oil Palm Res.* **2019**, *31*, 238–247. [[CrossRef](#)]
121. Park, H.B.; Lee, B.; Kloepper, J.W.; Ryu, C.M. One Shot-Two Pathogens Blocked: Exposure of *Arabidopsis* to Hexadecane, a Long Chain Volatile Organic Compound, Confers Induced Resistance against Both *Pectobacterium carotovorum* and *Pseudomonas syringae*. *Plant Signal. Behav.* **2013**, *8*, e24619. [[CrossRef](#)]
122. Parmagnani, A.S.; Kanchiswamy, C.N.; Paponov, I.A.; Bossi, S.; Malnoy, M.; Maffei, M.E. Bacterial Volatiles (MVOC) Emitted by the Phytopathogen *Erwinia amylovora* Promote *Arabidopsis thaliana* Growth and Oxidative Stress. *Antioxidants* **2023**, *12*, 600. [[CrossRef](#)]
123. Majrashi, T.A.; Sabt, A.; Abd El Salam, H.A.; Al-Ansary, G.H.; Hamissa, M.F.; Eldehna, W.M. An Updated Review of Fatty Acid Residue-Tethered Heterocyclic Compounds: Synthetic Strategies and Biological Significance. *RSC Adv.* **2023**, *13*, 13655–13682. [[CrossRef](#)]
124. Zheng, L.; Yan, X.; Xu, J.; Chen, H.; Lin, W. Hymeniacidin Perleve Associated Bioactive Bacterium *Pseudomonas* sp. NJ6-3-1. *Appl. Biochem. Microbiol.* **2005**, *41*, 29–33. [[CrossRef](#)]
125. Taghavi, N. Enhancement in Biodegradation of Thermoplastics Using UV-Pretreatment and Biosurfactant. Ph.D. Thesis, The University of Auckland, Auckland, New Zealand, 2021.
126. Sehim, A.E.; Amin, B.H.; Yosri, M.; Salama, H.M.; Alkhalifah, D.H.; Alwaili, M.A.; Elghaffar, R.Y.A. GC-MS Analysis, Antibacterial, and Anticancer Activities of *Hibiscus sabdariffa* L. Methanolic Extract: In Vitro and In Silico Studies. *Microorganisms* **2023**, *11*, 1601. [[CrossRef](#)]
127. Zhang, X.; Li, Q.; Xu, W.; Zhao, H.; Guo, F.; Wang, P.; Wang, Y.; Ni, D.; Wang, M.; Wei, C. Identification of MTP Gene Family in Tea Plant (*Camellia sinensis* L.) and Characterization of CsMTP8.2 in Manganese Toxicity. *Ecotoxicol. Environ. Saf.* **2020**, *202*, 110904. [[CrossRef](#)]
128. Adesanwo, J.K.; Ajayi, I.S.; Ajayi, O.S.; Igbeneghu, O.A.; McDonald, A.G. Identification of Chemical Constituents and Evaluation of the Antibacterial Activity of Methanol Extract and Fractions of the Leaf of *Melanthera scandens* (Schum. et Thonn.) Roberty. *J. Explor. Res. Pharmacol.* **2019**, *4*, 31–40. [[CrossRef](#)]
129. Hawar, S.N.; Taha, Z.K.; Hamied, A.S.; Al-Shmgani, H.S.; Sulaiman, G.M.; Elsilk, S.E. Antifungal Activity of Bioactive Compounds Produced by the Endophytic Fungus *Paecilomyces* sp. (JN227071.1) against *Rhizoctonia solani*. *Int. J. Biomater.* **2023**, *2023*, 2411555. [[CrossRef](#)]
130. Idris, N. Potential of Hexadecanoic Acid as Antimicrobials in Bacteria and Fungi That Cause Decay in Mustard Greens *Brassica juncea* L. *Int. J. Appl. Biol.* **2022**, *6*, 36–42.
131. Phukan, H.; Brahma, R.; Mitra, P.K. An Endophytic Fungus Associated with *Kayea assamica* (King & Prain): A Study on Its Molecular Phylogenetics and Natural Products. *S. Afr. J. Bot.* **2020**, *134*, 314–321. [[CrossRef](#)]
132. Jiao, S.; Chen, W.; Wang, E.; Wang, J.; Liu, Z.; Li, Y.; Wei, G. Microbial Succession in Response to Pollutants in Batch-Enrichment Culture. *Sci. Rep.* **2016**, *6*, 21791. [[CrossRef](#)]
133. Boadu, A.; Karpoormath, R.; Nlooto, M. Spondias Mombin: Biosafety and GC-MS Analysis of Anti-Viral Compounds from Crude Leaf Extracts. *Adv. Tradit. Med.* **2023**, *1*–24. [[CrossRef](#)]
134. Firdaus, M.; Kartikaningsih, H.; Sulifah, U. *Sargassum* spp Extract Inhibits the Growth of Foodborne Illness Bacteria. *AIP Conf. Proc.* **2019**, *2202*, 020083. [[CrossRef](#)]
135. Williams, A.; Langridge, H.; Straathof, A.L.; Muhamadali, H.; Hollywood, K.A.; Goodacre, R.; de Vries, F.T. Root Functional Traits Explain Root Exudation Rate and Composition across a Range of Grassland Species. *J. Ecol.* **2022**, *110*, 21–33. [[CrossRef](#)]
136. Shahbaz, E.; Ali, M.; Shafiq, M.; Atiq, M.; Hussain, M.; Balal, R.M.; Sarkhosh, A.; Alferez, F.; Sadiq, S.; Shahid, M.A. Citrus Canker Pathogen, Its Mechanism of Infection, Eradication, and Impacts. *Plants* **2023**, *12*, 123. [[CrossRef](#)]
137. Chuah, X.; Okechukwu, P.; Amini, F.; Teo, S. Eicosane, Pentadecane and Palmitic Acid: The Effects in In Vitro Wound Healing Studies. *Asian Pac. J. Trop. Biomed.* **2018**, *8*, 490–499. [[CrossRef](#)]
138. Ghorbani, A.; Tafteh, M.; Roudbari, N.; Pishkar, L.; Zhang, W.; Wu, C. *Piriformospora indica* Augments Arsenic Tolerance in Rice (*Oryza sativa*) by Immobilizing Arsenic in Roots and Improving Iron Translocation to Shoots. *Ecotoxicol. Environ. Saf.* **2021**, *209*, 111793. [[CrossRef](#)]

139. Laport, M.; Santos, O.; Muricy, G. Marine Sponges: Potential Sources of New Antimicrobial Drugs. *Curr. Pharm. Biotechnol.* **2009**, *10*, 86–105. [[CrossRef](#)]
140. Dimmock, J.; Turner, W.; Baker, H. Antimicrobial Evaluation of Diastereoisomeric 2-Dimethylaminomethyl-6-phenylcyclohexanols and Related Esters. *J. Pharm. Sci. Res.* **1975**, *64*, 995–997. [[CrossRef](#)]
141. Hostettmann–Kaldas, M.; Nakanishi, K. Moronic Acid, a Simple Triterpenoid Keto Acid with Antimicrobial Activity Isolated from *Ozoroa mucronata*. *Planta Medica* **1979**, *37*, 358–360. [[CrossRef](#)]
142. Huang, Y.; Sun, C.; Guan, X.; Lian, S.; Li, B.; Wang, C. Butylated Hydroxytoluene Induced Resistance Against *Botryosphaeria dothidea* in Apple Fruit. *Front. Microbiol.* **2021**, *11*, 599062. [[CrossRef](#)]
143. Qian, F.; Huang, X.; Su, X.; Bao, Y. Responses of Microbial Communities and Metabolic Profiles to the Rhizosphere of *Tamarix ramosissima* in Soils Contaminated by Multiple Heavy Metals. *J. Hazard. Mater.* **2022**, *438*, 129469. [[CrossRef](#)]
144. Fathi, M.; Ghane, M.; Pishkar, L. Phytochemical Composition, Antibacterial, and Antibiofilm Activity of *Malva sylvestris* Against Human Pathogenic Bacteria. *Jundishapur J. Nat. Pharm. Prod.* **2022**, *17*, e114164. [[CrossRef](#)]
145. Nautiyal, V.; Pradhan, S.; Dubey, R.C. In Silico Molecular Docking Analysis and ADME Prediction of Cow Urine Derived 1-Heneicosanol against Some Bacterial Proteins. *Vegetos* **2023**, *38*, 1–7. [[CrossRef](#)]
146. Salim, S.A. Identification of Active Pharmaceutical Ingredients in *Thevetia nerifolia* Juss Leaf Callus Using Analysis of GC-MS. *Indian J. Public Health Res. Dev.* **2018**, *9*, 1019–1023.
147. Li, J.; Wang, Y.; Zheng, L.; Li, Y.; Zhou, X.; Li, J.; Gu, D.; Xu, E.; Lu, Y.; Chen, X.; et al. The Intracellular Transporter AtNRAMP6 Is Involved in Fe Homeostasis in Arabidopsis. *Front. Plant Sci.* **2019**, *10*, 1124. [[CrossRef](#)]
148. Angeli, A.; Petrou, A.; Kartsev, V.; Lichitsky, B.; Komogortsev, A.; Capasso, C.; Geronikaki, A.; Supuran, C.T. Synthesis, Biological and In Silico Studies of Griseofulvin and Usnic Acid Sulfonamide Derivatives as Fungal, Bacterial and Human Carbonic Anhydrase Inhibitors. *Int. J. Mol. Sci.* **2023**, *24*, 2802. [[CrossRef](#)]
149. Xuan, T.D.; Roni, Y.; Andriana, Y.; Khanh, T.D.; Anh, T.T.T.; Kakar, K.; Haqani, M.I. Chemical Profile, Antioxidant Activities and Allelopathic Potential of Liquid Waste from Germinated Brown Rice. *Allelopath. J.* **2018**, *45*, 89–100. [[CrossRef](#)]
150. Ahmad, K.S. Environmental Contaminant 2-Chloro-N-(2,6-diethylphenyl)-N-(methoxymethyl)acetamide Remediation via *Xanthomonas axonopodis* and *Aspergillus niger*. *Environ. Res.* **2020**, *182*, 109117. [[CrossRef](#)]
151. Vázquez-Fuentes, S.; Pelagio-Flores, R.; López-Bucio, J.; Torres-Gavilán, A.; Campos-García, J.; de la Cruz, H.R.; López-Bucio, J.S. N-Vanillyl-Octanamide Represses Growth of Fungal Phytopathogens In Vitro and Confers Postharvest Protection in Tomato and Avocado Fruits against Fungal-Induced Decay. *Protoplasma* **2021**, *258*, 729–741. [[CrossRef](#)]
152. Minerdi, D.; Maggini, V.; Fani, R. Volatile Organic Compounds: From Figurants to Leading Actors in Fungal Symbiosis. *FEMS Microbiol. Ecol.* **2021**, *97*, fiab067. [[CrossRef](#)]
153. Kawle, P.R. Synthesis and Antituberculosis Action of Symmetric Acridin-9-yl-bisbenzothiazol-2-yl-amines. *Indian J. Chem. IJC* **2023**, *62*, 691–695. [[CrossRef](#)]
154. Brodzka, A.; Kowalczyk, P.; Trzepizur, D.; Koszelewski, D.; Kramkowski, K.; Szymczak, M.; Wypych, A.; Lizut, R.; Ostaszewski, R. The Synthesis and Evaluation of Diethyl Benzylphosphonates as Potential Antimicrobial Agents. *Molecules* **2022**, *27*, 6865. [[CrossRef](#)]

**Disclaimer/Publisher’s Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.