



HAL
open science

How do climate models represent wind resources in Southern Africa and predict their future changes?

Chao Tang, Béatrice Morel

► **To cite this version:**

Chao Tang, Béatrice Morel. How do climate models represent wind resources in Southern Africa and predict their future changes?. EGU24, Apr 2024, Vienna, Austria, Austria. 10.5194/egusphere-egu24-8855 . hal-04555247

HAL Id: hal-04555247

<https://hal.univ-reunion.fr/hal-04555247>

Submitted on 22 Apr 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

EGU24-8855, updated on 22 Apr 2024

<https://doi.org/10.5194/egusphere-egu24-8855>

EGU General Assembly 2024

© Author(s) 2024. This work is distributed under the Creative Commons Attribution 4.0 License.



How do climate models represent wind resources in Southern Africa and predict their future changes?

Chao Tang and Béatrice Morel

ENERGY-Lab, University of Reunion, 15 avenue René Cassin, CS 92003, 97744 Saint-Denis Cedex 9, Reunion, France

As intermittent meteorological resources become increasingly vital in our energy system, it is crucial to better understand how a changing climate may affect weather variability and, in turn, influence future wind energy production. This study evaluates the performance of climate models in reproducing wind resources over Southern Africa (SA), and in assessing potential impacts of climate change. An ensemble of climate simulations is evaluated over SA, which includes simulations from three Regional Climate Models (RCM; i.e., CCLM4, RegCM4, and REMO2009) that participated in the Coordinated Regional Downscaling Experiment program over Africa (CORDEX-Africa) at a horizontal resolution of approximately 25 km, and the coarser-resolution ones from their four driving General Circulation Models (GCMs; i.e., HadGEM2-ES, MPI-ESM-LR, MPI-ESM-MR, and NorESM1-M) from the Coupled Model Intercomparison Project Phase 5 (CMIP5). The simulated wind is first compared to the reference datasets, derived from ground-based measurements and reanalyses, during 2000-2023 at 3-hour intervals, covering both surface and hub-height (100 m) levels. The performances of both RCMs and GCMs are quantified and compared in terms of their representation of wind resource characteristics, including the mean wind speed and its spatio-temporal variability at hourly-to-annual timescales. Then wind energy potential and capacity factors are also derived for the wind resource assessment in SA, where direct observational data is limited. Finally, the potential impact of climate change on the 100-meter wind energy potential in SA is assessed based on this ensemble of climate projections, up to 2099, under the RCP2.6 and RCP8.5 scenarios. This study helps to understand the performance difference between regional and global models in simulating wind resources and provides insight into how changing climate conditions might affect wind energy production over the long term in SA.